

A seven-state RNCCA that is not isomorphic to any septenary RNCCA

Barbara Wolnik,^{*} Anna Nenca,[†] Adam Dzedzej,[‡] and Bernard De Baets[§]

We consider two-dimensional cellular automata with the von Neumann neighborhood that satisfy two properties of interest from a modelling viewpoint: rotation symmetry (*i.e.*, the local rule is invariant under rotation of the neighborhood by 90 degrees) and number conservation (*i.e.*, the sum of all the cell states is conserved upon every update). It is known that if the number of states k is smaller than or equal to six, then each rotation-symmetric number-conserving cellular automaton is isomorphic to some k -ary one, *i.e.*, one with state set $\{0, 1, \dots, k-1\}$. In this paper, we exhibit an example of a seven-state rotation-symmetric number-conserving cellular automaton that is not isomorphic to any septenary one. This example strongly supports our plea that research into multi-state cellular automata should not only focus on those that have $\{0, 1, \dots, k-1\}$ as state set.

I. INTRODUCTION

Cellular Automata (CAs) are discrete dynamical systems that evolve according to a simple local rule, but can nevertheless produce very complex dynamics. For this reason, such systems are willingly used to build models of various phenomena, in particular physical ones.

The definition of a CA requires the choice of a cellular space \mathcal{C} , a neighborhood \mathcal{N} of every cell, a state set Q and a local rule f . Although the specification of the quadruple $(\mathcal{C}, \mathcal{N}, Q, f)$ has a great influence on the resulting model, usually little attention is paid to what elements Q contains, but only to how many there are. Therefore, theoretical research — if it does not consider some general hypothetical state set Q — mostly concerns the so-called k -ary CAs (binary, ternary, quaternary, quinary, and so on), for which the state set equals $\{0, 1, \dots, k-1\}$ for some integer $k \geq 2$ (see, for instance, [1–4]).

The above approach seems justified for the following reason. Suppose that \mathcal{C} and \mathcal{N} are fixed and that two finite equinumerous sets Q and \tilde{Q} are given. Then for each CA $A = (\mathcal{C}, \mathcal{N}, Q, f)$, using any bijection $\phi : Q \rightarrow \tilde{Q}$, one can define an isomorphic CA $\tilde{A} = (\mathcal{C}, \mathcal{N}, \tilde{Q}, \tilde{f})$, which means that when identifying the states q and $\phi(q)$ (for any $q \in Q$), the CAs A and \tilde{A} are indistinguishable (for the strict meaning, see Definition 4). For example, in the one-dimensional case, where the dynamics of a given cellular automaton is usually represented by a space-time diagram, the fact that two CAs $A = (\mathcal{C}, \mathcal{N}, Q, f)$ and $\tilde{A} = (\mathcal{C}, \mathcal{N}, \tilde{Q}, \tilde{f})$ are isomorphic simply means that if we

choose some collection of colors to represent the states of \tilde{A} and then for every state $q \in Q$ we use the same color as for the state $\phi(q)$, then the corresponding space-time diagrams for A and \tilde{A} will be exactly the same. Perhaps the most common situation is when considering binary CAs and the set of states can be named $\{0, 1\}$ or $\{\text{white}, \text{black}\}$ or $\{\text{dead}, \text{alive}\}$ or $\{\text{healthy}, \text{sick}\}$ or $\{\text{empty}, \text{occupied}\}$, to name but a few.

However, in many situations, only CAs having some additional property P are considered (for instance, resulting from the properties of the model being built). Since isomorphy does not need to preserve P , for two equinumerous sets Q and \tilde{Q} , the family of all Q -state CAs satisfying P may differ significantly from the family of all \tilde{Q} -state CAs satisfying P . For example, when modeling various kinds of physical phenomena, very often some additional condition is imposed on the local rule f related to the symmetries of the neighborhood \mathcal{N} (for instance, rotation invariance) or related to some conservation laws (of mass, energy, and so on). While in the former case we still do not have to worry about the choice of Q , in the latter case, it is known that selecting Q may be important.

We are encountering such a situation in the case of two-dimensional rotation-symmetric number-conserving CAs with the von Neumann neighborhood (RNCCAs) (see Section II for definitions). Tanimoto *et al.* [5] proved that such CAs with at most four states are all trivial, regardless of Q , *i.e.*, if just $|Q| \leq 4$, then the identity is the only RNCCA with such a state set. Later, Imai *et al.* [6] showed that the case of five states is a bit more complicated: if the elements of a given five-element set Q form an arithmetic progression, then there are exactly three RNCCAs with this state set Q ; if the elements of Q do not form an arithmetic progression, then there is only one RNCCA with this state set (the identity). Thus the choice of Q has a big influence on the number of such CAs. However, it was also proved that the set $\{0, 1, 2, 3, 4\}$ can be treated as a basic set of states in the sense that any five-state RNCCA is isomorphic to some quinary RNCCA, *i.e.*, with state set $\{0, 1, 2, 3, 4\}$ (see [6]). An analogous result was proved in [7] for the more complicated case of six states: the au-

^{*} Institute of Mathematics, Faculty of Mathematics, Physics and Informatics, University of Gdańsk, 80-308 Gdańsk, Poland; KERMIT, Department of Data Analysis and Mathematical Modelling, Faculty of Bioscience Engineering, Ghent University, Coupure links 653, B-9000 Gent, Belgium

[†] Institute of Informatics, Faculty of Mathematics, Physics and Informatics, University of Gdańsk, 80-308 Gdańsk, Poland

[‡] Institute of Mathematics, Faculty of Mathematics, Physics and Informatics, University of Gdańsk, 80-308 Gdańsk, Poland

[§] KERMIT, Department of Data Analysis and Mathematical Modelling, Faculty of Bioscience Engineering, Ghent University, Coupure links 653, B-9000 Gent, Belgium

thors showed that the set $\{0, 1, 2, 3, 4, 5\}$ can be treated as a basic set for all six-state RNCCAs. Indeed, there are exactly 116 senary RNCCAs (with the state set $\{0, 1, 2, 3, 4, 5\}$) and any six-state RNCCA is isomorphic to some of those 116 RNCCAs. Thus, knowing the dynamics of 116 senary RNCCAs, we know the dynamics of any six-state RNCCA.

The results of [5–7] lead to the open question whether or not the following general statement holds.

Question 1. *Is every RNCCA with k states isomorphic to some k -ary RNCCA?*

Although the proof technique used in [7] does not easily generalize to larger state sets, it was believed that the answer to Question 1 would be positive.

Unfortunately, reality turns out to be different. In this paper, we demonstrate that the answer to the above question is negative. More precisely, we provide an example of a seven-state RNCCA that is not isomorphic to any septenary RNCCA (*i.e.*, with the state set $\{0, 1, 2, 3, 4, 5, 6\}$).

II. PRELIMINARIES

As cellular space we consider the two-dimensional infinite grid \mathbb{Z}^2 . At every discrete time step, every cell (i, j) in \mathbb{Z}^2 is assigned a state from a given set Q . We assume the state set Q to be a finite set of real numbers containing 0.

A configuration is any mapping from the grid \mathbb{Z}^2 to Q . The set of all possible configurations $Q^{\mathbb{Z}^2}$ is denoted as X . For a given configuration $\mathbf{x} \in X$, the value of cell (i, j) is denoted by $x_{i,j}$. By X_{fin} we denote the set of all *finite configurations*, *i.e.*, those configurations that are almost everywhere equal to zero:

$$X_{\text{fin}} = \{\mathbf{x} \in X \mid \{(i, j) \in \mathbb{Z}^2 \mid x_{i,j} \neq 0\} \text{ is finite}\}.$$

A configuration $\mathbf{x} \in X$ is called *totally spatially periodic* if there exist positive integers p_1 and p_2 such that for each $(i, j) \in \mathbb{Z}^2$ it holds that

$$x_{i,j} = x_{i+p_1,j} = x_{i,j+p_2}$$

(see, for instance, [8]). In other words, a totally spatially periodic configuration consists of some rectangular pattern (with sizes p_1 and p_2) that is periodically repeated in both dimensions of \mathbb{Z}^2 . The couple (p_1, p_2) is referred as the period of \mathbf{x} . The set of all totally spatially periodic configurations is denoted as X_{per} .

Generally, a neighborhood is understood as an m -tuple (v_1, v_2, \dots, v_m) of distinct vectors of \mathbb{Z}^2 that specify for a given cell the relative positions of its neighbor cells (w.r.t. that cell). In this paper, we consider the so-called von Neumann neighborhood (with radius one), for a given cell consisting of the cell itself and its four adjacent cells; hence, we use $\mathcal{N} =$

$\{(0, -1), (-1, 0), (0, 0), (1, 0), (0, 1)\}$. For the sake of convenience, for a given configuration $\mathbf{x} \in X$ we list the states of the cells in the neighborhood of a cell (i, j) as

$$\begin{array}{ccc} & x_{i,j-1} & \\ x_{i-1,j} & x_{i,j} & x_{i+1,j} \\ & x_{i,j+1} & \end{array}.$$

The new state of a cell (i, j) depends only on the states of the cells in its neighborhood at the previous time step through a local rule f . Thus, in our case, the domain of f is given by

$$\mathcal{D} = \left\{ \begin{array}{c} u \\ l \ c \ r \\ d \end{array} \mid u, l, c, r, d \in Q \right\}.$$

A given local rule $f : \mathcal{D} \rightarrow Q$ generates a global rule $F : X \rightarrow X$ in the usual way: for any $\mathbf{x} \in X$, the configuration $F(\mathbf{x})$ is given by

$$F(\mathbf{x})_{i,j} = f \left(\begin{array}{ccc} & x_{i,j-1} & \\ x_{i-1,j} & x_{i,j} & x_{i+1,j} \\ & x_{i,j+1} & \end{array} \right).$$

By a CA we mean the quadruple $(\mathbb{Z}^2, \mathcal{N}, Q, f)$.

In our investigation, we focus on CAs that have two properties that are important from the point of view of applications. The first one is rotation symmetry, which means that the local rule of the CA is invariant under rotation of the neighborhood by 90 degrees.

Definition 1. *A CA is called rotation-symmetric if its local rule f satisfies for any $u, l, c, r, d \in Q$*

$$f \left(\begin{array}{c} u \\ l \ c \ r \\ d \end{array} \right) = f \left(\begin{array}{c} r \\ u \ c \ d \\ l \end{array} \right). \quad (1)$$

The second property of CAs we are interested in is number conservation, which means that the sum of all the states in any configuration remains constant throughout the evolution of the automaton. Since the cellular space \mathbb{Z}^2 is infinite, it is convenient to consider number conservation in the sense of *finite* or *periodic* number conservation.

Definition 2. *Let F be the global rule of a CA. We say that this CA is finite number-conserving if for any $\mathbf{x} \in X_{\text{fin}}$, it holds that*

$$\sum_{(i,j) \in \mathbb{Z}^2} F(\mathbf{x})_{i,j} = \sum_{(i,j) \in \mathbb{Z}^2} x_{i,j}.$$

Definition 3. *Let F be the global rule of a CA. We say that this CA is periodic number-conserving if for any $\mathbf{x} \in X_{\text{per}}$, it holds that*

$$\sum_{0 \leq i < p_1, 0 \leq j < p_2} F(\mathbf{x})_{i,j} = \sum_{0 \leq i < p_1, 0 \leq j < p_2} x_{i,j},$$

where (p_1, p_2) is the period of \mathbf{x} .

Durand *et al.* [9] proved that Definitions 2 and 3 are equivalent.

Theorem 1. *A cellular automaton $(\mathbb{Z}^2, \mathcal{N}, Q, f)$ is finite number-conserving if and only if it is periodic number-conserving.*

The above theorem allows us to use the term “number-conserving” having in mind any of these two definitions. If a CA is number-conserving, then we also call its local and global rules number-conserving. In this paper, a two-dimensional CA with the von Neumann neighborhood that is both rotation-symmetric and number-conserving will be abbreviated as an RNCCA.

A detailed characterization of all RNCCAs with up to five states, as mentioned in the introduction, can be found in [5, 6]. The case of six states was considered in [7] (on finite grids only, but this is the same as considering the infinite grid using Definition 3), where a very tedious and technical approach allowed to pick out all senary RNCCAs from the huge set of all two-dimensional rotation-symmetric CAs with the state set $\{0, 1, 2, 3, 4, 5\}$ and the von Neumann neighborhood (note that the cardinality of this set is greater than $6^{6^{5/4}}$, a number with 1513 digits). It appeared that there are only 116 such RNCCAs. Moreover, the authors gave a full characterization of the set of all RNCCAs for any six-element state set, from which it resulted, *inter alia*, that any six-state RNCCA is isomorphic to some senary RNCCA (see [7] for details). When stating that two CAs are isomorphic (sometimes the term *equivalent* is also used), we have the following definition in mind.

Definition 4. *We say that a CA $A = (\mathbb{Z}^2, \mathcal{N}, Q, f)$ is isomorphic to a CA $\tilde{A} = (\mathbb{Z}^2, \mathcal{N}, \tilde{Q}, \tilde{f})$ if there exists a bijection $\phi : Q \rightarrow \tilde{Q}$ such that for any $u, l, c, r, d, x \in Q$*

$$f \begin{pmatrix} u \\ l \ c \ r \\ d \end{pmatrix} = x \quad \Leftrightarrow \quad \tilde{f} \begin{pmatrix} \phi(u) \\ \phi(l) \ \phi(c) \ \phi(r) \\ \phi(d) \end{pmatrix} = \phi(x). \quad (2)$$

The results of [5–7] can be summarized as follows.

Theorem 2. *Let $k \leq 6$. Then every RNCCA with k states is isomorphic to some k -ary RNCCA.*

In the next section, we present a counterexample showing that Theorem 2 cannot be generalized, even for $k = 7$.

III. COUNTEREXAMPLE

The method described in [7] allows to enumerate all septenary RNCCAs with the help of a computer. One can also use a much more general method introduced in [10] based on the characterization of number-conserving CAs with the von Neumann neighborhood. Either of these two ways results in a complete list of all 30144 septenary RNCCAs, which can be found in [11] (this dataset contains the full list, before reduction according to conjugation or reflection dependency).

Now, let us define the following RNCCA $A_* = (\mathbb{Z}^2, \mathcal{N}, Q, f_*)$, by setting $Q = \{0, 2, 3, 4, 5, 6, 8\}$ and

$$f_* \begin{pmatrix} u \\ l \ c \ r \\ d \end{pmatrix} = \begin{cases} 2 \cdot N_8(u, r, d, l) & , \text{ if } c = 0 \\ 8 - 2 \cdot N_0(u, r, d, l) & , \text{ if } c = 8 \\ N_6(u, r, d, l) & , \text{ if } c = 2 \\ 6 - N_2(u, r, d, l) & , \text{ if } c = 6 \\ c & , \text{ otherwise,} \end{cases}$$

where $N_b(u, r, d, l)$ denotes the number of occurrences of b among u, r, d, l . The local rule f_* thus has a very simple interpretation in the language of particle representation: every cell with 8 particles gives 2 particles to each empty cell in its neighborhood (therefore also every empty cell gets 2 particles from each cell in its neighborhood having 8 particles) and every cell with 6 particles gives 1 particle to each cell in its neighborhood having 2 particles (therefore also every cell with 2 particles gets 1 particle from each cell in its neighborhood having 6 particles). The remaining cells do not change their states. Since the local rule f_* can be interpreted in such a simple way in terms of motion of particles, one can see that it is rotation-symmetric and number-conserving. We will argue that the RNCCA $A_* = (\mathbb{Z}^2, \mathcal{N}, Q, f_*)$ is not isomorphic to any septenary RNCCA.

Since there are as many as $7!$ possible bijections between two seven-element sets, a brute force approach checking for all septenary RNCCAs whether Eq. (2) holds or not requires $30144 \times 7! \times 7^5 = 2\,553\,416\,248\,320$ operations, which looks rather daunting. Instead, we propose to use the method of invariants: for each RNCCA A we define some simple numerical parameter $\gamma(A)$, which proves invariant under isomorphy. Then we calculate the value of this parameter for all septenary RNCCAs. If no value turns out to be equal to the value $\gamma(A_*)$, then this proves that A_* is not isomorphic to any septenary RNCCAs.

Let $\frac{a|b}{c|d}$ denote the configuration $\mathbf{x} \in X_{\text{per}}$ with period $(2, 2)$, for which $x_{0,0} = a$, $x_{1,0} = b$, $x_{0,1} = c$ and $x_{1,1} = d$. For a given $A = (\mathbb{Z}^2, \mathcal{N}, Q, f)$ and $q \in Q$, let $\gamma(A, q)$ denote the number of configurations $\mathbf{x} \in X_{\text{per}}$ with period $(2, 2)$ such that $F(\mathbf{x})$ is the homogeneous configuration in which all cells have state q , *i.e.*,

$$\gamma(A, q) = \left| \left\{ \frac{a|b}{c|d} \in X_{\text{per}} \mid F\left(\frac{a|b}{c|d}\right) = \frac{q|q}{q|q} \right\} \right|$$

and let $\gamma(A) = \max_{q \in Q} \gamma(A, q)$.

It is easy to see that if $A = (\mathbb{Z}^2, \mathcal{N}, Q, f)$ and $\tilde{A} = (\mathbb{Z}^2, \mathcal{N}, \tilde{Q}, \tilde{f})$ are isomorphic by a bijection $\phi : Q \rightarrow \tilde{Q}$, then for any $q \in Q$ it holds that $\gamma(A, q) = \gamma(\tilde{A}, \phi(q))$, thus also $\gamma(A) = \gamma(\tilde{A})$, which means that the parameter γ is invariant under isomorphy.

For the example local rule f_* , one can calculate that $\gamma(A_*) = \gamma(A_*, 4) = 41$. It is shown in Figure 1 which configurations are transformed into the homogeneous configuration with 4's. To keep the schema readable, we grouped these configurations according to significantly

different patterns, ignoring rotations and axial symmetries. For example, all eight configurations $\frac{6}{2}|\frac{4}{4}$, $\frac{2}{4}|\frac{6}{4}$, $\frac{4}{4}|\frac{2}{6}$, $\frac{4}{6}|\frac{4}{2}$, $\frac{2}{6}|\frac{4}{4}$, $\frac{4}{2}|\frac{4}{6}$, $\frac{4}{4}|\frac{6}{2}$, $\frac{6}{4}|\frac{2}{4}$ are transformed into $\frac{4}{4}|\frac{4}{4}$, but we present only the first one and give the information that it represents a group of eight configurations ($\times 8$). We have calculated $\gamma(A)$ for all septenary RNC-




FIG. 1. Configurations with period $(2, 2)$ transformed by A_* into the homogeneous configuration with 4's.

CAs and only the following values were obtained: 1, 5, 9, 13 and 17.

TABLE I. Number of septenary RNCCAs with a given value of $\gamma(A)$.

$\gamma(A)$	1	5	9	13	17
number of septenary RNCCAs	48	696	6048	9328	14024

Table I presents the numbers of septenary RNCCAs with a given $\gamma(A)$. As the largest possible value of $\gamma(A)$ for any septenary RNCCA is 17, we conclude that A_* is not isomorphic to any septenary RNCCA.

IV. CONCLUSIONS

There was a common belief among CA researchers that it suffices to consider CAs with the state set $\{0, 1, \dots, k-1\}$. The authors of [7] were asked by an anonymous reviewer why they wrote down a complicated proof for the seemingly obvious fact that every RNCCA with six states is isomorphic to some senary RNCCA. The example in this paper shows that such effort is necessary, since Theorem 2 does not generalize to higher values of k . For general CAs, it is merely a matter of renaming states, so that one can always use the state set $\{0, 1, \dots, k-1\}$. However, number conservation (for instance) imposes some additional additive structure on the state set, implying that some natural arguments regarding the renaming of states become invalid. One indeed has to be very careful when performing such operation.

The local rule f_* was inspired by some models in which particles carrying different amounts of energy coexist. These models are quite natural to consider in the context of number-conserving CAs and our example shows that limiting the attention to the most popular k -ary world is not always sufficient.

-
- [1] F. G. Pazzona, G. Pireddu, and P. Demontis, Quasiequilibrium multistate cellular automata, *Physical Review E* **105**, 014116 (2022).
 - [2] E. L. Ruivo, P. P. de Oliveira, F. Lobos, and E. Goles, Shift-equivalence of k -ary, one-dimensional cellular automata rules, *Communications in Nonlinear Science and Numerical Simulation* **63**, 280 (2018).
 - [3] K. Bhattacharjee and S. Das, Reversibility of d -state finite cellular automata., *Journal of Cellular Automata* **11** (2016).
 - [4] N. Boccara and H. Fuk s, Number-conserving cellular automaton rules, *Fundamenta Informaticae* **52**, 1 (2002).
 - [5] N. Tanimoto, K. Imai, C. Iwamoto, and K. Morita, On the non-existence of rotation-symmetric von neumann neighbor number-conserving cellular automata of which the state number is less than four, *IEICE Trans. Inf. Syst.* **92-D**, 255 (2009).
 - [6] K. Imai, H. Ishizaka, and V. Poupet, 5-state rotation-symmetric number-conserving cellular automata are not strongly universal, in *Cellular Automata and Discrete Complex Systems: 20th International Workshop, AUTOMATA 2014, Himeji, Japan, July 7-9, 2014, Revised Selected Papers*, edited by T. Isokawa, K. Imai, N. Matsui, F. Peper, and H. Umeo (Springer International Publishing, Cham, 2015) pp. 31–43.
 - [7] A. Dzedzej, B. Wolnik, A. Nenca, J. M. Baetens, and B. De Baets, Two-dimensional rotation-symmetric number-conserving cellular automata, *Information Sciences* **577**, 599 (2021).
 - [8] A. Dennunzio, E. Formenti, and M. Weiss, Multidimensional cellular automata: closing property, quasi-expansivity, and (un)decidability issues, *Theoretical Computer Science* **519**, 40 (2014).
 - [9] B. Durand, E. Formenti, and Z. R ka, Number-conserving cellular automata I: decidability, *Theoretical Computer Science* **299**, 523 (2003).
 - [10] B. Wolnik, A. Nenca, J. M. Baetens, and B. De Baets, A split-and-perturb decomposition of number-conserving cellular automata, *Physica D: Nonlinear Phenomena* **413**, 132645 (2020).
 - [11] A. Nenca, A. Dzedzej, and B. Wolnik, The complete list of two-dimensional rotation-symmetric number-conserving septenary cellular automata, <https://doi.org/10.34808/jr67-r637> (2022).