Pesticide concentration in three selected fish species and human health risk in the Lake Tana sub-basin, Ethiopia

This Accepted Manuscript (AM) is a PDF file of the manuscript accepted for publication after peer review, when applicable, but does not reflect post-acceptance improvements, or any corrections. Use of this AM is subject to the publisher's embargo period and AM terms of use. Under no circumstances may this AM be shared or distributed under a Creative Commons or other form of open access license, nor may it be reformatted or enhanced, whether by the Author or third parties. By using this AM (for example, by accessing or downloading) you agree to abide by Springer Nature's terms of use for AM versions of subscription articles: https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms

The Version of Record (VOR) of this article, as published and maintained by the publisher, is available online at: <u>https://doi.org/10.1007/s10661-023-11594-y</u>. The VOR is the version of the article after copy-editing and typesetting, and connected to open research data, open protocols, and open code where available. Any supplementary information can be found on the journal website, connected to the VOR.

For research integrity purposes it is best practice to cite the published Version of Record (VOR), where available (for example, see ICMJE's guidelines on overlapping publications). Where users do not have access to the VOR, any citation must clearly indicate that the reference is to an Accepted Manuscript (AM) version.

1 Pesticide concentration in three selected fish species and human health risk in the Lake Tana Sub-basin, Ethiopia

- 2 Wondie Zelalem^{1, 2, *}, Wassie Anteneh³, Minwyelet Mingist², Mulugeta Kibret⁴, Enyew Adgo⁵, Felegush Erarto², Jasmine
- 3 De Rop⁶, Andrée De Cock⁷, Pieter Spanoghe⁶, Peter L.M Goethals⁷, Jan Nyssen⁸, Elie Verleyen⁹, Alain De Vocht^{1, 10}
- 4 1 UHasselt, Centre for Environmental Sciences, Agoralaan, 3590 Diepenbeek, Belgium
- ² Department of Fisheries and Aquatic Sciences, Bahir Dar University, Bahir Dar, Ethiopia
- 6 ³ Intergovernmental Authority on Development (IGAD), Djibouti
- ⁴ Department of Biology, Bahir Dar University, Bahir Dar, Ethiopia
- 8 ⁵ Department of Natural Resource Management, Bahir Dar University, Bahir Dar, Ethiopia
- 9 ⁶ Department of Plants and Crops, Ghent University, Ghent, Belgium
- 10 ⁷ Department of Animal Sciences and Aquatic Ecology, Ghent University, Ghent, Belgium
- 11 ⁸ Department of Geography, Ghent University, Ghent, Belgium
- ⁹ Department of Biology, Ghent University, Ghent, Belgium
- ¹⁰ PXL University College, Hasselt, Belgium
 *Correspondence author; Email: wondiehm@gmail.com, ORCID ID: <u>0000-0001-5322-7100</u>

14 Abstract

- 15 Pesticide use has increased in the Lake Tana sub-basin due to increased agricultural activity, potentially endangering 16 nontargeted organisms. To assess its potential impact on fish health and fish-consuming human populations, pesticide 17 concentrations in the fillet and liver tissue of three fish species, namely Labeobarbus megastoma, L. tsanensis, and 18 Oreochromis niloticus, were investigated in Lake Tana. Fish samples were taken from the lake near the rivers of Ribb and 19 Gumara, which flow through agricultural areas where considerable amounts of pesticides have been applied. A total of 96 20 fish samples were collected. Liquid chromatography coupled with tandem mass spectrometry (LC-MS/MS) revealed the 21 presence of ten pesticides. Pyrimethanil was frequently detected in 96 % of liver and 65 % of fillet samples at a median 22 concentration of 33.9 μ g kg⁻¹ and 19.7 μ g kg⁻¹, respectively. The highest concentration of pyrimethanil was found in L. 23 megastoma (1850.0 μ g kg⁻¹). Labeobarbus megastoma also had the highest concentration of oxamyl (507.0 μ g kg⁻¹) and 24 flazasulfuron ($\frac{60.1}{\mu g}$ kg⁻¹) detected in the liver tissue. The highest concentration of carbaryl ($\frac{56.5}{\mu g}$ kg⁻¹) was found in 25 the liver tissue of O. niloticus. Fish tissue samples from the two study sites contained pyrimethanil, oxamyl, carbaryl, and 26 flazasulfuron. Only pyrimethanil showed a statistically significant difference between the two sites and the species L. 27 *megastoma* and *L. tsanensis*. The amounts of pesticides found in the fish species pose no direct risk to the health of fish 28 consumers human population. However, the results show that the lake ecosystem needs immediate attention and regular
- 29 monitoring of the rising pesticide usage in the lake watershed.
- 30 Keywords: Lake Tana, pesticide, *Labeobarbus*, *Oreochromis*, health risk, toxicity

31 Introduction

32 Pesticides have been used in agriculture for a long time to protect crops from weeds, pests, and diseases while increasing 33 crop yields (Lengai et al., 2020; Sabzevari & Hofman, 2022). They have also been used to reduce crop deterioration during 34 storage and extend shelf life, ensuring food security for the world's ever-growing population (Sharma et al., 2019). About 35 4.2 million tons of pesticides, including herbicides, insecticides, and fungicides, are utilized worldwide annually (FAO, 36 2021). In developing countries such as Ethiopia, the government has encouraged farmers to use pesticides and fertilizers 37 in smallholder farming (WHO & FAO, 2019). However, because farmers and agricultural workers frequently lack proper 38 personal protection equipment and may not understand the labels with safety instructions, human health and environmental 39 hazards associated with pesticide use are commonly noticed in those nations (Sarkar et al., 2021).

- 40 Although pesticides improve crop yield, their extensive use produces residual effects on food products. Moreover, they 41 could develop problems associated with chemical buildup in aquatic organisms, including fish, and negatively impact 42 aquatic life and human health (Amenyogbe et al., 2021). They potentially affect the growth and reproductive efficiency of 43 non-targeted organisms due to their bio-magnification and persistent nature (Deribe et al., 2013). Some pesticides, such as 44 organochlorine pesticides (OCPs), can last for many years in the environment, are deposited in sediments (MacKay & 45 Fraser, 2000; Deribe et al., 2013), and concentrated in the top predators of food chains (Di et al., 2017). As a result, most 46 pesticide-related ecosystem studies focused on those OCPs (Marchand et al., 2010; Sifakis et al., 2017; Yohannes et al., 47 2017; Worku et al., 2022). Organophosphate pesticides (OPPs) and other synthetic pesticides, which are less persistent 48 than OCPs, are also widely used by farmers worldwide (Manuelmolina-Ruiz et al., 2014). Most of them are water-soluble 49 and taken up by organisms through skin contact, inhalation, or ingestion as food particles. As a result, these pesticides are 50 hazardous to aquatic species, and their residues might affect the phosphorylation of the acetylcholinesterase enzyme 51 (AChE) at nerve terminals in non-targeted animals, causing malfunctioning (Gultekin et al., 2000; Kushwaha et al., 2016). 52 Pesticides can limit the food sources available to fish through their toxicity effect on invertebrate prey, potentially reducing 53 fish growth and survival. At high concentrations in water bodies, pesticides can affect fish by changing their growth rates, 54 progeny survival, and behavior. This influences fish population stability and structure (Holden, 1972). Toxicity effects on 55 fish can also lead to histological and hormonal changes in females, which may reduce the number of oocytes produced 56 (Forsgren et al., 2013).
- 57 The fish species of the genus *Labeobarbus* are the most numerous and endemic to Lake Tana. Two of the five *Labeobarbus* 58 species most frequently caught for commercial fishing are *L. tsanensis* and *L. megastoma* (Gebremedhin et al., 2019). 59 *Labeobarbus megastoma* is a piscivores fish that dominantly feeds (up to 67%) on fish, while *L. tsanensis* is feeding 60 dominantly on benthos 43%, 18% mollusks, and 14% detritus/substratum (Sibbing & Nagelkerke, 2001). Similarly, 61 *Oreochromis niloticus*, a highly commercially demanded species in the lake (Nagelkerke et al., 1995; Vijverberg et al., 62 2013), is an omnivorous fish that feeds on plankton and aquatic plants (Tesfahun & Alebachew, 2023).

63 The Lake Tana sub-basin, one of Ethiopia's most productive places, is suitable for growing a variety of staple food crops, 64 including Teff (*Eragrostis tef*), Finger millet (*Eluesine coracana*), rice (*Oryza sativa*), maize (*Zea mays*), grass pea 65 (*Lathyrus sativus*) and chickpea (*Cicer arietinum*), as well as cash crops like Khat (*Catha edulis*) and other vegetables 66 (tomato, cabbage, onion, garlic, potato, pepper, etc.) (Abera, 2017). Recently, pesticide consumption has increased due to 67 increased agricultural activity in the area, particularly in irrigation-based dry season cash crops and vegetable cultivations 68 (Agmas & Adugna, 2020; Abera et al., 2022). Farmers spray various pesticides on those crops and vegetables in the field

69 and at their home storage, but they are usually not fully informed or aware of the dangers of the pesticides they use (Agmas 70 & Adugna, 2020). Improper spraying techniques, dropping empty containers into water bodies, careless disposal, or 71 unintentional spills of residual solutions have all been noted as contamination routes in the catchment (Agmas & Adugna, 72 2020). Additionally, the lake has a large catchment area (approximately 16,750 km²), and the process of pesticides leaching 73 through soil erosion and surface runoff increases environmental hazards and may have detrimental effects on the lake's 74 biodiversity. For example, due to brain, endocrine, behavioral, and genetic abnormalities, as well as histopathological and 75 hematological changes, fish species may not be able to grow and reproduce normally (Kumari, 2020; Srivastava et al., 76 2016). Furthermore, pesticide residues may pose a health risk to human communities that consume fish (Agmas & Adugna, 77 2020; Abera et al., 2022). Despite the increasing use of pesticides in the lake catchment, no research has been done yet to 78 determine the concentrations and the possible harmful impacts on non-target organisms except a recent work by Abera et 79 al. (2022), and to the best of our knowledge, there was no work done on the concentration and effect of pesticide residues, 80 particularly on the economically important and endangered fish species in the lake. Therefore, this study aimed to 1) 81 determine the concentration of pesticides routinely used in the fillet and liver tissue of three commercially important fish 82 species in the study area, 2) assess any potential impact of pesticide residues on fish health and 3) assess the effect of the 83 measured pesticides on the health of fish-consuming communities.

84 Materials and Methods

85 Study area

Lake Tana and its tributaries, such as Ribb and Gumara (Fig. 1), provide important ecosystem services to people living in
the catchment. The two river watersheds are the most important agricultural areas for most staple crops through smallholder
irrigation farming. These water bodies supply drinking water for humans and animals. They are also important for fisheries. *Labeobarbus* species, *Oreochromis niloticus*, and *Clarias garipinus* are commercially important and frequently consumed
fish species in the research area. People living near the lake and along the river use rainwater to grow crops and vegetables
during the rainy season. While in the dry season, farmers use the lake and river waters through furrow and pump irrigation.





Fig. 1 Map of the study locations in Lake Tana and sampling points (Red circles) at the Ribb River Mouth (RRM) andGumara River Mouth (GRM) sites.

95 Fish sampling and preparations

96 Fish samples were collected from Lake Tana's shore near the mouths of the Gumara and Ribb Rivers. Sampling was 97 executed between September and October 2021 using multifilament gillnets (10, 12, and 14 cm mesh sizes) and catches 98 from fishers at the sampling points. Specimens of three fish species, Labeobarbus tsanensis, L. megastoma, and 99 Oreochromis niloticus, with a weight of more than 200 grams and with a total length of at least 23 cm were identified and 100 selected (Table 1). Tissue samples were collected based on the approach stated by Rosseland et al. (2001). Each fish sample 101 was extracted with stainless steel disposable scalpel blades with a handle, yielding around 50 g (wet weight) of the fillet 102 (white meat taken from the dorsal muscle) and 2 to 5g of the liver. Samples were wrapped in aluminum foil, frozen (-20° C), 103 and transported to Belgium in an ice-filled cooling cabinet. The pesticide analysis was done in the laboratory for 104 phytopharmaceuticals at Ghent University.

- 105 Each fillet and the liver sample were homogenized for 10 minutes with an Ultra-Turrax (T 25 ultra Turrax IKA). The
- 106 homogenizer was cleaned with acetone after each homogenization to prevent cross-contamination. Based on the analyses'
- 107 weight requirements, samples were weighed and stored in falcon tubes.

Table 1 Biometric data and feeding mode of the three fish species sampled from Lake Tana between September andOctober 2021.

		Gumara river mouth			Ribb river mouth		
Fish species	Feeding	Total Length	Weight		Total Length	Weight (g)*	
	mode	(cm)*	(g)*	n	(cm)*		n
L. megastoma	Piscivore	29.7 - 42.5	205 - 615	8	31.7 - 43.5	285 - 700	8
L. tsanensis	Insectivore	30.5 - 39.5	270 - 670	8	30.1 - 35.2	250 - 445	8
O. niloticus	Herbivore	23.2 - 26.4	220 - 295	8	23.5 - 26.5	220 - 290	8

* minimum-maximum values, n: number of samples analyzed.

111 Pesticide Analysis

112 Based on data from farmers and district (locally called woreda) agricultural experts on local pesticide use in 2019 and

113 2020, a list of 109 relevant pesticide products was prepared for the multi-residue analysis before the pesticide study. Liquid 114 chromatography coupled with tandem mass spectrometry (LC-MS/MS) with the setting defined in Table S2 (supplementary

material) was used to detect and quantify all of the screening pesticides (n = 109) (Table S1 (supplementary material)).

116 The fish tissue samples were extracted using a modified quick, easy, cheap, effective, rugged, and safe (QuEChERS)

117 approach. This approach has a small number of steps, is highly reliable, and yields excellent recoveries for a wide range of

110

118 pesticides from various chemical families (Kim et al., 2019).

119 In a QuEChERS tube, 10 ml of acetonitrile was added to 10 g of fillet sample and agitated for one minute. Subsequently, 120 a salt solution was added (6 g magnesium sulfate, 0.75 g citrate sesquihydrate, 1.5 g citrate dihydrate, 1.5 sodium chloride), 121 and the tube was vigorously shaken. The sample was then homogenized for 2 minutes in an ultra-turax mixer before being 122 centrifuged for 5 minutes at 3,000 rpm. In a 10 ml flask, 1 ml of the supernatant was diluted in 9 ml ultrapure/miliQ/ water, 123 and 1.5 ml of the solution was then transferred to a vial for LC-MS/MS analysis. The liver samples were extracted by adding 10 ml of acetonitrile to 2 - 5 g of material and then adding water to a total mass of 10 g. The falcon tubes were 124 125 shaken for 1 minute before adding the salt mixture. The remaining stages were the same as the fillet, except for a clean-up 126 phase. As a result, 7 ml of the supernatant was transferred to an SPE tube, which was agitated for 5 minutes before being 127 centrifuged at 3000 rpm for 5 minutes. In a 10 ml flask, 1 ml of the supernatant was diluted in 9 ml ultrapure (miliQ) water. 128 Finally, 1.5 ml of the sample was subsampled into a vial for pesticide determination.

129 Method validation

The minimal concentration at which the analyte was identified and established by comparing measured signals from samples with known low amounts of the analyte with those of blank samples. The present method was developed and validated using the ICH Q2(R1) guideline. The pesticide residues in the fish tissue samples were validated and quantified using a calibration curve from samples spiked with the multi-compound stock standard solution. The linear range of the calibration curve was developed using five concentration levels between 0.001 and 0.1 mg l⁻¹. The spike-placebo recovery method was used to determine the validation of the analyte's lowest detectable concentration in a sample (limit of detection or LOD) and the lowest confirmed level with adequate precision and recovery (limit of quantification or LOQ). A blank

sample was spiked and analyzed under the same conditions in four replicates. The LOD and LOQ values for the pesticide
analysis in LC-MS/MS were set as 0.0003 mg l⁻¹ and 0.00086 mg l⁻¹, respectively.

139 Human health risk assessment

- 140 A human health risk assessment was conducted to evaluate if the pesticides found in the lake threatened those who included
- fish in their diet. The Acute Reference Dose (ARfD) and Acceptable Daily Intake (ADI) were utilized as expected, with no impact levels for acute and chronic pesticide exposure. The Estimated Daily Intake (EDI) for a given pesticide was
- 143 calculated using the following formula (Mahmood & Malik, 2014; Hamid et al., 2017)

144 EDI =
$$\frac{C_p \times F_c}{bw}$$
 (1)

145 Cp is the maximum pesticide residue concentration in white fish meat in μ g/kg wet weight. Fc is the amount of fish food 146 consumed by a person per day in kg/day. bw is the fish-consuming adult in kg.

- 147 The hazard quotient (HQ) for acute and chronic estimations was performed using the following formula (Hamid et al.,148 2017)
- $HQ_{acute} = \frac{EDI}{ARfD}$ (2)
- 150 $HQ_{chronic} = \frac{EDI}{ADI}$ (3)

For the Ethiopian community who live around the water bodies and have high access to fish in their diet, the average daily intake rate was taken as 0.03 kg/day (Yohannes et al., 2014) and assumed the average body weight of 60 kg for an adult Ethiopian for the calculation (Teklu et al., 2015). As liver tissue is not consumed in the area, the pesticide concentration in the fillet was only considered for the risk assessment.

- 155 The Hazard Index (HI) was taken as the sum of the chronic or acute hazard quotients separately of all pesticides detected 156 in the fish fillet. Pesticide residues could potentially produce acute/chronic health hazards if the percentage of HI is greater 157 than 100 (Lozowicka, 2015).
- 158 Data analysis

159 The Kruskal-Wallis ANOVA test was used to examine differences in pesticide residue concentrations in the tissue among 160 species, and the Mann-Whitney U test between tissue types using the R package version 4.2.0. The Dunn test was used to 161 evaluate any significant differences between the groups. As a result of an uneven distribution of the pesticide concentration 162 data in the fish tissue, the median values were utilized to represent the pesticide concentrations in the different fish species 163 and tissue types. For these statistical comparisons, a pesticide concentration below the LOQ was taken to be half of the 164 corresponding assigned limiting value. Hence the number of > LOQ values recorded among the pesticides detected was 165 very small (< 60%); the deterministic approach was used to determine the exposure of the fish consumers and the hazard 166 quotients by taking an average pesticide concentration for values that had two or greater than two > LOO values. For the 167 risk assessment, the LOQ cutoff value, 0.00086 mg/kg, was used for fish fillet samples with none > LOQ value or those 168 detected pesticides with < LOQ values as a maximum pesticide concentration, considering the worst-case scenario. The 169 exposure analysis was performed using Microsoft Excel.

170 Results

171 Pesticide concentrations in fish samples

172 Ten pesticides were found in the fish fillet and liver tissue samples of the three fish species under investigation. The 173 pesticide types insecticides and fungicides were found in equal proportion (40% of each) from the detected pesticides, 174 while herbicides constitute 20%. The level of pesticides quantified ranges from below LOQ to 1850.0 μ g kg⁻¹ wet weight. 175 The pesticide most commonly found and had the highest concentration was pyrimethanil. The difference between the two 176 research sites was statistically significant (p < 0.001) due to the higher median concentration of pyrimethanil (49.7 µg kg⁻ 177 ¹) at the site near the mouth of the Gumara River (GRM) compared to 2.1 µg kg⁻¹ at the site near the mouth of the Ribb 178 **River (RRM)** (Table 2). Pyrimthanil was detected in 65% of fillet samples with a mean concentration of 96.6 μ g kg⁻¹ and 179 96% of liver tissue samples with a mean concentration of 104.0 μ g kg⁻¹ for the three fish species. A statistically significant 180 difference in pyrimethanil concentration between species (p = 0.043) was observed, with the highest concentration in L. 181 megastoma and the lowest in L. tsanensis (Fig. 2 and Table 3). The mean detected concentration of pyrimethanil was much 182 higher in the liver (258.8 µg kg⁻¹) than in the fillet (55.4 µg kg⁻¹) of L. megastoma. However, pyrimethanil was detected in 183 higher concentration in the fillet of O. niloticus (mean = 359.8 μ g kg⁻¹) and L. tsanensis (mean = 99.1 μ g kg⁻¹) than in the 184 liver tissue of those species (Table 4). Like pyrimethanil, carbaryl was quantified in the fillet and liver tissue of the three 185 species and found in both studied sites. The highest mean concentration of carbaryl (10.4 µg kg⁻¹) was recorded in O. 186 niloticus. However, variations among species were not statistically different.



Fig. 2 Boxplot showing pyrimethanil concentration in the tissue of the three studied fish species in the studied sites inLake Tana.

189 Table 2 Mean concentrations and range of detected pesticides in Lake Tana fish species sampled between September and190 October 2021 at the GRM and RRM sampling site.

		GRM			RRM
	Pesticide	Median	Range	Median	Range
Pesticide	type	$(\mu g \ kg^{-1})$	(µg kg-1)	$(\mu g \ kg^{-1})$	(µg kg ⁻¹)
Benalaxyl	Fungicide	2.2*	< LOD – 2.2	< LOD	-
Butachlor	Herbicide	< LOD	-	1.1	< LOD – 1.1
Carbaryl	Insecticide	<mark>0.4</mark>	0.2 - 56.5	0.4	0.2 - 2.2
Difenoconazole	Fungicide	0.2*	< LOD - 2.2	< LOD	
Flazasulfuron	Herbicide	2.2	2.2 - 60.1	2.2	< LOD – 2.2
Imidacloprid	Insecticide	< LOD	-	0.9	0.4 - 2.2
Oxamyl	Insecticide	216.5	32.0 - 507.0	24.4	2.2 - 136.9
Pirimiphos-methyl	Insecticide	< LOD	-	0.4*	< LOD – 0.4
Pyrimethanil	Fungicide	<mark>49.7</mark>	2.2 - 1850.0	2.2	0.4 - 171.7
Tebuconazole	Fungicide	0.2*	< LOD – 0.2	<lod< td=""><td>-</td></lod<>	-

191 *quantified only in a single sample; $LOD = 0.0003 \ \mu g \ kg^{-1}$; $LOQ = 0.00086 \ \mu g \ kg^{-1}$

192 Oxamyl and flazasulfuron were detected in both study sites (Table 2). However, oxamyl was found in the tissues of the 193 two Labeoburbus species but at a level below LOD in O. niloticus (Table 3). Oxamyl was found in L. megastoma and L. 194 *tsanensis*, and the highest concentration was detected in the liver tissue of L. megastoma (median = 361.7 μ g kg⁻¹) (Tables 195 3 & 4). Though its concentration was small compared to liver tissue, oxamyl was also detected in the fillet of L. megastoma, 196 while it was below LOD in L. tsanensis and O. niloticus. Imidacloprid and flazasulfuron were detected in different 197 quantities in the tissues of the three fish species under investigation at site RRM (Table 5.4), whereas imidacloprid was 198 below LOD at site GRM (Table 3). Imidacloprid was only detected in the liver tissue of L. tsanensis; however, it was 199 detected in the fillets of L. megastoma and O. niloticus at the RRM site (Table 4). However, the concentrations of oxamyl, imidacloprid, and flazasulfuron in neither species nor the tissue types were statistically different. 200

sampling sites in Lake Tana and variations of pesticide types among species is presented in the *p* value.

Sampling	Pesticide	Pesticide concentrations in fish species (µg kg ⁻¹)				
site		L. megastoma	L. tsanensis	O. niloticus	<mark>p value</mark>	
GRM	Carbaryl	0.2	0.4	<mark>4.6</mark>	<mark>0.041</mark>	
	Flazasulfuron	2.2	<mark>2.2*</mark>	<mark>2.2</mark>	<mark>0.446</mark>	
	Imidacloprid	<lod< th=""><th><lod< th=""><th><lod< th=""><th>•</th></lod<></th></lod<></th></lod<>	<lod< th=""><th><lod< th=""><th>•</th></lod<></th></lod<>	<lod< th=""><th>•</th></lod<>	•	
	<mark>Oxamyl</mark>	<mark>361.7</mark>	<lod< th=""><th>< LOD</th><th>•</th></lod<>	< LOD	•	
	Pyrimethanil	<mark>82.6</mark>	30.1	<mark>55.1</mark>	<mark>0.074</mark>	
RRM	Carbaryl	<mark>0.4</mark>	•	<mark>1.3</mark>	<mark>0.739</mark>	
	Flazasulfuron	2.2	2.2	<mark>2.2</mark>	•	
	Imidacloprid	<mark>0.4</mark>	2.2	<mark>0.4</mark>	0.223	
	<mark>Oxamyl</mark>	<lod< th=""><th><mark>24.4</mark></th><th><lod< th=""><th>•</th></lod<></th></lod<>	<mark>24.4</mark>	<lod< th=""><th>•</th></lod<>	•	

²⁰¹ Table 3. The median pesticide concentration detected in the tissues (fillet + liver) of the three fish species at the two

		Pyrimethanil	<mark>22.8</mark>	<mark>2.2</mark>	<mark>2.2</mark>	0.010	
203	* quantifie	d only in a single sar	nple				
204	Benalaxyl, a p	besticide that was rar	ely discovered, was	found in the l	iver tissue of L.	megastoma at the GRM site (m	iedian
205	= 2.2 μg kg ⁻¹); nevertheless, dife	noconazole was fou	nd in the fille	et tissue (mediar	$n = 0.22 \ \mu g \ kg^{-1}$) of <i>L. megas</i>	toma.
206	Similarly, but	achlor and pirimipho	os-methyl were deter	cted only in R	RM. Butachlor	was detected in the liver tissue	e of L.
207	<i>tsanensis</i> (me	dian = 1.1 μg kg ⁻¹),	and pirimiphos-met	hyl was found	d in the fillet of	<i>O. niloticus</i> (median = $0.4 \mu g$	kg-1).
208	Tebuconazole	was also detected in	the fillet of L. mega	<i>stoma</i> in the O	GRM site (Table	<u>4).</u>	

209 Table 4 Detected pesticide concentration in the tissues of the studied fish species sampled between September and October

		Mean pes	sticide concent	ration in µg kg ⁻¹ w	vet weight	Õ
Pesticide	L. me	egastoma	L. ts	anensis	<i>O. n</i>	iloticus
-	liver	fillet	liver	fillet	Liver	fillet
Benalaxyl	2.2*	< LOD	< LOD	< LOD	< LOD	< LOD
Butachlor	< LOD	< LOD	1.1	< LOD	< LOD	< LOD
Carbaryl	2.2*	0.2	2.2*	0.4	29.3	3.1
Difenoconazole	< LOD	0.2*	< LOD	< LOD	< LOD	< LOD
Flazasulfuron	2.2	< LOD	2.2	<lod< td=""><td>2.2</td><td>< LOD</td></lod<>	2.2	< LOD
Imidacloprid	< LOD	0.2	2.2*	< LOD	< LOD	0.4*
Oxamyl	361.7	32.0*	24.4	< LOD	< LOD	< LOD
Pirimiphos-methyl	< LOD	< LOD	< LOD	< LOD	< LOD	0.4*
Pyrimethanil	15.0	<mark>54.1</mark>	<mark>7.9</mark>	<mark>35.6</mark>	<mark>25.4</mark>	<mark>94.9</mark>
Tebuconazole	< LOD	0.2	< LOD	< LOD	< LOD	< LOD

- 211 * quantified only in a single sample; $LOD = 0.0003 \ \mu g \ kg^{-1}$; $LOQ = 0.00086 \ \mu g \ kg^{-1}$
- 212 Human health risk assessment
- People eating fish from Lake Tana might be exposed to the seven pesticides detected in the fillet of the three fish species
 (Table 4). The risk assessment estimation considers all values found in the fillet, including < LOQ values. The highest
 daily exposure of consumers was recorded for oxamyl (0.016 μg/kg bw/d) (Table 5). The calculated values for HQ acute
- and HQ chronic were low, with the highest value of 1.60% found for oxamyl. Similarly, the HI values for both HI acute
- and HI chronic (Table 5) showed a lower value than the threshold value of 100, indicating that adverse non-cancer health
- 218 effects are not likely to occur or the health risk for the fish meat-consuming community is very low.

Table 5 Adult human exposure to selected quantified (> LOQ) pesticides in the fillet of fish species in the Lake Tana sub-

basin. Maximum detected quantity (Cp), Acceptable daily intake (ADI), Acceptable reference dose (ARfD), Estimated

221 Daily Intake (EDI), and the percentage of hazard quotient for acute (HQ acute) and chronic (HQ chronic) cases.

Pesticide	ADI	ARfD	Max	EDI	HQ acute (%)	HQ chronic (%)
	$(\mu g / kg bw/d)$	(µg/kg bw)	$(\mu g \ kg^{-1})$	($\mu g/kg bw/d$)		
Carbaryl	7.5	10.0	14.0	0.007	0.07	0.10
Oxamyl	1.0	1.0	32.0	0.016	1.60	1.60
Pyrimethanil	170	1000	1206.6	0.603	0.06	0.35
				HI (%)	1.73	2.06

222 Discussion

223 Detected pesticides and the possible effect on fish species

The development, survival, and reproduction of aquatic animals are known to be impacted when surface waters are contaminated by high pesticide concentrations (Vonesh & Kraus, 2009; Akhter, 2019; Kumari, 2020). It has also been shown that several pesticides in water systems have affected primary and secondary production, including fish species in the food chain (Seeland et al., 2012; Müller et al., 2019). The effect, however, potentially depends on the physicochemical environment and may be ecoregion-specific, long-lasting, or delayed, yet most likely exacerbated by indirect effects resulting from interspecific interactions (Araújo et al., 2012; Müller et al., 2019).

230 The highest concentration and the frequent occurrence of pyrimethanil in the three fish species in Lake Tana indicate that 231 it is intensively used in the watershed. Pyrimethanil is one of the most widely used fungicides worldwide and has been 232 found in numerous aquatic environments (Lozowicka, 2015; De Cock et al., 2021). It is used mainly on fruits and vegetables 233 and was manufactured to treat resistant fungus strains such as Botrytis spp, which affects onions and other vegetables and 234 fruits (Tournas, 2005; Abo-Elyousr et al., 2020). Pyrimethinil is also applied for seed treatment on cereals (EFSA, 2011). 235 It affects algae, zooplankton, and macroinvertebrates which are potential food sources for fish species. For example, for 236 the model organism Scenedemus acutus in a 48h chronic test and Daphnia magna in a 48-h acute test, the average LC50 was 23.2 mg l^{-1} and 3.6 mg l^{-1} , respectively (Araújo et al., 2012). They also reported that the reproduction of adult D. 237 238 magna was adversely affected when exposed to 0.5 mg 1⁻¹ of pyrimethanil over many generations under variable 239 temperatures between 20 and 27 °C. Although the pesticide concentration was not directly measured from the water in 240 Lake Tana, the maximum concentration (up to 1.85 mg kg-1) in the fish tissue suggests that the concentration was 241 potentially high. This high concentration of pyrimethanil could affect the normal development of fish organs. According 242 to Bernab et al. (2017), doses up to 50 µgl⁻¹ cause a histological alteration in tree frogs (Hyla intermedia). The effect of 243 this pesticide also elevated with increasing temperature. For example, Araújo et al. (2012) confirmed that a rise in ambient 244 temperature exacerbated the fungicide's toxicological effects on D. magna. Thus, with the relatively high dry season 245 temperature, the effect of pyrimethanil on aquatic invertebrates in Lake Tana might be higher, which could indirectly affect 246 fish species.

Metals and pesticides can be absorbed by aquatic macrophytes and algae (Dosnon-Olette et al., 2010, 2009), and organisms
 that directly consume those primary sources may have higher concentrations in their tissue. The higher mean concentration

of pyrimethanil in the tissue of *O. niloticus* (Table 3) might reflect the assimilation of the fungicide through the food chain.
Similarly, the highest concentration in the piscivore species *L. megastoma* (Table 4) could also result from its feeding habit,

despite pyrimethanil having less potential in bioaccumulation (Araújo et al., 2015; Lewis et al., 2016).

- Flazasulfuron, belonging to the sulfonylurea herbicide family, was detected in 29% of tissue samples from the three fish species with a maximum concentration of 0.06 mg kg⁻¹ in *L. megastoma*. This herbicide is known to prevent protein synthesis in plants by blocking branched-chain amino acids protein synthesis (Olette et al., 2008). It has also been shown to impact the chlorophyll pigments and leaf gas exchange (Frankart et al., 2003). While it is not toxic to aquatic animals (US EPA, 2007), it can indirectly have an impact because it might reduce plant and nonvascular aquatic plant productivity
- 257 (US EPA, 2007). For instance, flazasulfuron reduces the photosynthetic capacity of duckweed (*Lemna minor*) by 16% after
- 258 seven days of exposure to a 100 μ g l⁻¹ concentration (Olette et al., 2008).
- 259 The other frequently found pesticide in the three sampled fish species and the two study sites in Lake Tana was carbaryl 260 (Tables 2 & 3, Fig. S2 (supplementary material)). This is a carbamate insecticide that has been reported to have an impact 261 on several fish species. For example, Patnaik and Patra (2006) observed its impact on the blood cells in Clarias batrachus 262 following a 96-hour acute exposure, with a cell shape modified at 12.6 mg l⁻¹ and an LC50 at 15.6 mg l⁻¹. In addition, under a controlled investigation, Boran et al. (2010) found that rainbow trout had an LC50 at 2.5 mg l⁻¹ and 1.4 mg l⁻¹ over 24 263 264 and 96 hours of exposure time, respectively. Matos et al. (2007) also report the effect of carbaryl as it resulted in necrosis 265 of the liver tissue of *O. niloticus* when exposed to a concentration of 0.25 mg l⁻¹ for seven days. The pH and temperature 266 of the aquatic environment significantly impact the persistence of carbamate insecticides in natural waters (Aly & El-Dib, 267 1971). Carbaryl generally has a low water persistence (US EPA, 2003). However, the amount recorded in the fish tissue, 268 up to 56.5 µg kg⁻¹ in this study, indicates its excessive pesticide use. The detected amount may not impose risk in the short 269 term. However, prolonged exposure possibly imposes environmental and health impacts, including oxidative stress in the 270 fish species, as observed in several animal groups (Ribera et al., 2001; Matos et al., 2007). In addition, the effect of 271 pesticides is more pronounced on juveniles than on adults of the same species (Lin et al., 2007), so any prolonged and 272 extensive use could affect fish recruitment and population growth.
- 273 The other carbamate insecticide that was detected in the two Labeobarbus species of Lake Tana was oxamyl (Table 3, Fig. 274 S1 (supplementary material)), which is relatively more toxic than carbaryl for aquatic organisms (Lewis et al., 2016; 275 Alvarez et al., 2022). Oxamyl is applied as an insecticide on field crops, vegetables, fruits, and ornamental plants (Alvarez 276 et al., 2022). It is moderately toxic to fish, and its LC50 after 96 h exposure for species such as bluegill sunfish, goldfish, 277 and rainbow trout were found to be 5.6 mg l⁻¹, 27.5 mg l⁻¹, and 42 mg l⁻¹, respectively (Smith, 1982). Less persistent 278 pesticides such as oxamyl are rarely found in fish tissue. For example, Polat et al. (2018) found a 45 µg kg⁻¹ concentration 279 in the fillet of sardine (Sardine pilchardus) among nine studied fish species in the Iskenderun Bay of Turkey. However, 280 in this investigation, oxamyl was detected in 7% of the fish tissue sampled from the lake (Table 3), with the highest 281 concentration (507 µg kg⁻¹) in the liver tissue of *L. megastoma*, indicating its excessive use in the catchment. Although 282 additional research on fish sensitivity to pesticides in Lake Tana is needed, the amount of oxamyl in this study showed the 283 necessity for strict regulation and monitoring.
- 284 The fungicides, benalaxyl, difenoconazole and tebuconazole were found < LOQ in fish sampled from GRM site. Albeit
- their concentrations were low, the presence of these pesticides in this site indicates that they have been applied in the
- 286 Gumara River catchment. Benalaxyl and difenoconazole, however, were not listed in Abera et al. (2022)'s list of pesticides
- used in the lake watershed, indicating that there are still other pesticides available in the local market that may come through

288 a different marketing route and circulate through an informal supply network as reported in Mengistie et al. (2016). 289 Benalaxyl is a widely used fungicide on tomatoes, onions, and potatoes in the field, while difenoconazole and tebuconazole 290 are applied on cereals and other field crops (Lewis et al., 2016). Although these fungicides had a high $\log K_{ow}$ value used 291 (Lewis et al., 2016) compared to other pesticides found in Lake Tana, they were only found in one sample of the fillet of 292 L. megastoma. This suggests that they are probably applied infrequently in the catchment or, due to their lipophilic nature 293 (Houbraken et al., 2016; Rasool et al., 2022), adsorbed in organic wastes in sediment. Studies showed that tebuconazole 294 could cause a decrease in hemoglobin and the number of red blood cells in cyprinids when applied with a 2.5 mg l^{-1} 295 concentration for a 96-h exposure (Cilingir Yeltekin et al., 2020). The LC50 of tebuconazole for fish is 4.4 mg l⁻¹ (Lewis 296 et al., 2016). Similarly, the broad-spectrum triazole fungicide difenoconazole is extensively used in agriculture (Jiang et 297 al., 2022). It influences immune-related pathways and energy, lipid, and amino acid metabolism in the early stages of 298 zebrafish development (Teng et al., 2018). Although the examined fish species' susceptibility to those pesticides was not 299 assessed, the observed concentration, which was < LOQ, suggests they very likely do not cause any risk to fish at the 300 current application level in the lake catchment.

301 The other commonly used insecticides, imidacloprid and pirimiphos-methyl, are frequently detected in many aquatic 302 environments (Vignet et al., 2019). Both insecticides are moderately toxic to fish (Lewis et al., 2016). Imidacloprid was 303 detected in the three studied fish species sampled from RRM. According to Mhadhbi and Beiras (2012), when imidacloprid 304 concentrations in the water were above 0.04 mg 1-1, it posed pericardial edema and spinal deformities, and it also reduced 305 hatching success when its concentration reached above 0.4 mg l⁻¹ in turbot (*Psetta maxima*). In the Ribb and Gumara 306 catchments, crops like rice and other cereals, including maize, are grown and commonly treated with pirimiphos-methyl 307 and imidacloprid. However, the fact that both pirimiphos-methyl and imidacloprid were found in the fish samples taken 308 from the RRM site suggests that farmers in the Ribb River catchment may have used these insecticides in greater quantities 309 than in Gumara.

310 In general, the number of herbicides, fungicides, and insecticides found in the tissue of the three fish species in Lake Tana 311 indicates how much pesticide use has polluted the lake water. Pesticides used by farmers in the Gumara and Ribb rivers to 312 reduce crop loss and increase the quality of their crops, vegetables, and fruits are indispensable. However, reducing agricultural losses by using too many of these agrochemicals should not come at the expense of aquatic life. Increasing 313 314 production using intensive agrochemicals could not be sustainable because a rise in pesticide use seriously harms an 315 ecosystem's health and natural resource. The best way to recommend sustainable production and an environmentally 316 friendly approach is using integrated pest management, which weighs natural alternatives to chemical pesticides and only 317 uses them as a last resort (Barzman et al., 2015). Using less toxic and less persistent pesticides is also better for creating 318 more resilient, sustainable food systems. Monitoring the type and quantity of pesticides used and assessing their impact on 319 biodiversity should be part of the management plan of the lake and its catchment, even though the concentrations currently 320 seem to have no adverse toxicological effect on fish species at the current application level.

321 Pesticides risk to human health

Fish ingest pesticide residues mainly through the epidermis of the gills and ingested food (Braunbeck et al., 1998). When a fish is consumed, the pesticides accumulated in its edible tissue can be transferred to humans. Liver tissue is the detoxification center and stores toxicants in many studied fish species (Braunbeck et al., 1998; Macirella et al., 2022). In this study, however, some pesticides, e.g., pyrimethanil, were found in higher concentrations in the fillet tissue of *L*.

326 tsanensis and O. niloticus (Table 4). Consequently, humans who consume fish meat from the lake are exposed to these 327 pollutants. Two to three different pesticides were found in some fillet samples, and their impact on the health of fish and 328 its consumers is likely greater than that of just a single pesticide due to their combined effect (Kortenkamp, 2007; 329 Hernández et al., 2013). No clear maximum residual limit (MRL) for the three pesticides was available for the risk 330 assessment (Table 5). However, from the metadata for most of the animal products in Great Britain, a maximum value of 331 50 µg kg⁻¹ for carbaryl and pyrimethanil and 10 µg kg⁻¹ for oxamyl are used (Lewis et al., 2016). Considering those 332 conservative MRL values, the detected concentration of pyrimethanil in 12 samples and one sample for oxamyl was higher 333 than the limit. The risk assessment, however, might reflect the worst-case scenario because we did not consider the impact 334 of washing, soaking with solutions, cooking, and freezing procedures that could lower the concentration of pesticides in 335 the tissue (Witczak, 2009; Alaboudi et al., 2021; Islam et al., 2022). Overall, the HQ and HI values showed that, at the 336 current application practices, pesticides pose no known harm to human health or consumers of fish collected from the lake. 337 However, the detection of ten pesticides in the fish tissue in this study shows that the use of pesticides along the river 338 catchments needs urgent attention and close monitoring to safeguard the public health of local communities and the 339 population of both culturally endangered and commercially important fish species.

340 Conclusion

341 The contamination of ten pesticide residues (2 herbicides, 4 fungicides, and 4 insecticides) was detected in the three commercially important fish species. Variations in pesticide concentrations among fish species and between tissue types 342 343 were not statistically significant, except for pyrimethanil, which revealed a significant difference between L. megastoma 344 and L. tsanensis. L. megastoma was found to have the highest level of pesticide contamination (eight out of ten) among the 345 three studied fish species, which may be related to its' piscivorous feeding behavior. Pyrimethnil > carbaryl > oxamyl > 346 flazasulfuron were the most frequently occurring pesticides in the studied fish species with > LOQ values. Moreover, most 347 pesticides were found in higher concentrations in the fish's liver than in the fillet. For all of the dominantly detected 348 pesticides, the health risk calculations revealed that the health risk index was below the cutoff value, indicating that, given 349 current pesticide use, there is likely no harm to human health from pesticide exposure in fish. Due to the lack of samples 350 from the water and sediment, the study is limited in its ability to evaluate the effects of combined exposure to toxins and 351 their influence on the lake's biota. Additionally, the small sample size results in an uneven distribution of data about the 352 presence of pesticides in fish tissue. However, although the study was conducted on a limited sample size, only three 353 species and a single developmental stage, the results could serve as a preliminary step for further research, and the 354 information will be used to generate a management strategy for the fish populations in Lake Tana. Further investigations 355 are recommended, including a wide variety of fish species, a sizable sample at various stages of development, and 356 measurements of pesticide concentrations in the water and sediment for a more in-depth ecotoxicological assessment.

357 Ethics statement

358 The Ethical Committee of Bahir Dar University, College of Agriculture and Environmental Sciences, approved the study.

359 The care and use of animals followed all applicable international, national, and institutional guidelines.

360 Acknowledgment

- 361 The authors thank the Belgian Development Cooperation (VLIR-UOS) for financially supporting this research through the
- 362 Inter-University Cooperation of Flemish and Bahir Dar Universities (IUC-BDU). The VLIR-UOS project staff at Bahir
- 363 Dar University are acknowledged for the support and logistics provided. We thank Mr. Ferede Tsegaw and Mr. Kassahun
- **364** Erkyehun for their help in fish specimen collections.

365 Author contribution

- 366 Wondie Zelalem: conceptualization, investigation, methodology, formal analysis, writing-original draft; Wassie Anteneh,
- 367 Minwyelet Mingist: conceptualization, investigation, editing; Alain De Vocht: conceptualization, project administration,
- 368 methodology, investigation, data curation, editing; Jasmine De Rop, Andrée De Cock, Pieter Spanoghe, Peter L.M
- 369 Goethals: methodology, data curation, formal analysis, editing; Mulugeta Kibret, Enyew Adgo, Jan Nyssen, Elie Verleyen:
- 370 investigation, project administration, editing. Felegush Erarto: investigation, editing.

371 Funding

372 No funding was obtained for this study.

373 Data Availability

374 The corresponding author will provide the data upon reasonable request.

375 Competing interests

376 The authors declare no competing interests.

377 References

- Abera, B., Van Echelpoel, W. Van, De Cock, A. De, Tytgat, B., Kibret, M., Spanoghe, P., Mengistu, D., Adgo, E., Nyssen,
 J., Goethals, P. L. M., & Verleyen, E. (2022). Environmental and Human Health Risks of Pesticide Presence in the
 Lake Tana Basin (Ethiopia). *Sustainability*, *14*(21), 14008. https://doi.org/10.3390/su142114008
- Abera, M. (2017). Agriculture in the Lake Tana Sub-basin of Ethiopia. In A. S. and G. G. Stave K. (Ed.), *Social and Ecological System Dynamics* (pp. 375–397). Springer. https://doi.org/10.1007/978-3-319-45755-0_23
- Abo-Elyousr, K. A. M., Alamri, S. A. M., Hussein, M. M. A., Hassan, M. A. H., Abd El-Fatah, B. E. S., & Hashem, M.
 (2020). Molecular disparities among Botrytis species involved in onion umbel blight disease and its management
 using Bacillus subtilis PHYS7. *Egyptian Journal of Biological Pest Control*, 30(1), 1–12.
 https://doi.org/10.1186/s41938-020-0205-x

- Agmas, B., & Adugna, M. (2020). Attitudes and practices of farmers with regard to pesticide use in NorthWest Ethiopia.
 Cogent Environmental Science, 6(1), 1791462. https://doi.org/10.1080/23311843.2020.1791462
- 389 Akhter, Z. (2019). Pesticide Contamination in Aquatic Ecosystems. *International Journal of Scientific Research and* 390 *Review*, 7(January), 2356–2381. https://www.researchgate.net/publication/348150568
- Alaboudi, A. R., Almashhadany, D. A., & Jarrah Bashar Suleiman. (2021). Effect of Cooking and Freezing on Levels of
 pesticides residues in Local Fresh Fish. *Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj- Napoca. Animal Science and Biotechnologies*, 78(2), 28. https://doi.org/10.15835/buasvmcn-asb:2021.0007
- Alvarez, F., Arena, M., Auteri, D., Binaglia, M., Federica Castoldi, A., Chiusolo, A., Colagiorgi, A., Colas, M., Crivellente,
 F., de Lentdecker, C., Egsmose, M., Fait, G., Ferilli, F., Gouliarmou, V., Nogareda, L. H., Ippolito, A., Istace, F.,
 Jarrah, S., Kardassi, D., ... Villamar Bouza, L. (2022). Peer review of the pesticide risk assessment of the active
 substance oxamyl. *EFSA Journal*, 20(5). https://doi.org/10.2903/j.efsa.2022.7296
- Aly, O. M., & El-Dib, M. A. (1971). Studies on the persistence of some carbamate insecticides in the Aquatic Environment I. Hydrolysis of sevin, baygon, pyrolan, and dimetilan in waters. *Water Research Pergamon Press*, 5, 1191–1205.
- Amenyogbe, E., Huang, J. S., Chen, G., & Wang, Z. (2021). An overview of the pesticides' impacts on fishes and humans.
 International Journal of Aquatic Biology, 9(1), 55–65. https://doi.org/10.22034/ijab.v9i1.972
- 402 Araújo, C. V. M., Shinn, C., Müller, R., Moreira-Santos, M., Shinn, C., Espíndola, E. L., & Ribeiro, R. (2015). *The*403 *Ecotoxicity of Pyrimethanil for Aquatic Biota Skin microbiome of Pelophylax perezi frogs View project The*404 *Ecotoxicity of Pyrimethanil for Aquatic Biota*. https://doi.org/10.13140/RG.2.1.1568.0485
- Barzman, M., Bàrberi, P., Birch, A. N. E., Boonekamp, P., Dachbrodt-Saaydeh, S., Graf, B., Hommel, B., Jensen, J. E.,
 Kiss, J., Kudsk, P., Lamichhane, J. R., Messéan, A., Moonen, A. C., Ratnadass, A., Ricci, P., Sarah, J. L., & Sattin,
 M. (2015). Eight principles of integrated pest management. *Agronomy for Sustainable Development*, *35*(4), 1199–
 1215. https://doi.org/10.1007/s13593-015-0327-9
- Bernabò, I., Guardia, A., MacIrella, R., Tripepi, S., & Brunelli, E. (2017). Chronic exposures to fungicide pyrimethanil:
 Multi-organ effects on Italian tree frog (Hyla intermedia). *Scientific Reports*, 7(1). https://doi.org/10.1038/s41598017-07367-6
- Boran, H., Altinok, I., & Capkin, E. (2010). Histopathological changes induced by maneb and carbaryl on some tissues of
 rainbow trout, Oncorhynchus mykiss. *Tissue and Cell*, 42(3), 158–164. https://doi.org/10.1016/j.tice.2010.03.004
- 414 Braunbeck, T..., Hinton, D. E., & Streit, B. (1998). Bioaccumulation of contaminants in fish. In T. Braunbeck, D. E. Hinton,
 415 & B. Streit (Eds.), *Fish Ecotoxicology* (86., Vol. 86). Birkhäuser, Basel. https://doi.org/10.1007/978-3-0348-8853416 0 12
- 417 Çilingir Yeltekin, A., Oğuz, A. R., Kankaya, E., Özok, N., & Güneş, İ. (2020). Hematological and biochemical response
 418 in the blood of alburnus tarichi (Actinopterygii: Cypriniformes: Cyprinidae) exposed to tebuconazole. *Acta*419 *Ichthyologica et Piscatoria*, 50(4), 373–379. https://doi.org/10.3750/AIEP/02931
- Andrée, D. C., Marie Anne Eurie, F., Niels, D. T., Isabel, G. A., Arne, D., Wout, V. E., Lenin, R. F., Jasmine, D. R.,
 Liesbeth, J., Pieter, S., Luis, D. G., & Peter L.M., G. (2021). From field to plate: Agricultural pesticide presence in

- the guayas estuary (Ecuador) and commercial mangrove crabs. *Environmental Pollution*, 289(August).
 https://doi.org/10.1016/j.envpol.2021.117955
- 424 Deribe, E., Rosseland, B. O., Borgstrøm, R., Salbu, B., Gebremariam, Z., Dadebo, E., Skipperud, L., & Eklo, O. M. (2013).
 425 Biomagnification of DDT and its metabolites in four fish species of a tropical lake. *Ecotoxicology and Environmental* 426 Safety, 95, 10–18. https://doi.org/10.1016/j.ecoenv.2013.03.020
- 427 Di, S., Liu, R., Tian, Z., Cheng, C., Chen, L., Zhang, W., Zhou, Z., & Diao, J. (2017). Assessment of tissue-specific
 428 accumulation, elimination and toxic effects of dichlorodiphenyltrichloroethanes (DDTs) in carp through aquatic food
 429 web. *Scientific Reports*, 7(1), 1–15. https://doi.org/10.1038/s41598-017-02612-4
- 430 Dosnon-Olette, R., Couderchet, M., & Eullaffroy, P. (2009). Phytoremediation of fungicides by aquatic macrophytes:
 431 Toxicity and removal rate. *Ecotoxicology and Environmental Safety*, 72(8), 2096–2101.
 432 https://doi.org/10.1016/j.ecoenv.2009.08.010
- Dosnon-Olette, R., Trotel-Aziz, P., Couderchet, M., & Eullaffroy, P. (2010). Fungicides and herbicide removal in
 Scenedesmus cell suspensions. *Chemosphere*, *79*(2), 117–123. https://doi.org/10.1016/j.chemosphere.2010.02.005
- 435 EFSA. (2011). Review of the existing maximum residue levels (MRLs) for pyrimethanil according to Article 12 of
 436 Regulation (EC) No 396/2005. EFSA Journal, 9(11). https://doi.org/10.2903/j.efsa.2011.2454
- 437 FAO. (2021). FAOSTAT ANALYTICAL BRIEF 16: Pesticides use Global, regional and country trends.
 438 https://www.fao.org/3/cb3411en/cb3411en.pdf
- Forsgren, K. L., Riar, N., & Schlenk, D. (2013). The effects of the pyrethroid insecticide, bifenthrin, on steroid hormone
 levels and gonadal development of steelhead (Oncorhynchus mykiss) under hypersaline conditions. *General and Comparative Endocrinology*, *186*, 101–107. https://doi.org/10.1016/j.ygcen.2013.02.047
- Frankart, C., Eullaffroy, P., & Vernet, G. (2003). Comparative effects of four herbicides on non-photochemical
 fluorescence quenching in Lemna minor. *Environmental and Experimental Botany*, 49, 159–168.
 www.elsevier.com/locate/envexpbot
- Gebremedhin, S., Bruneel, S., Getahun, A., Anteneh, W., & Goethals, P. (2019). The Endemic Species Flock of
 Labeobarbus spp. in L. Tana (Ethiopia) Threatened by Extinction: Implications for Conservation Management. *Water*, 11(12), 2560. https://doi.org/10.3390/w11122560
- 448 Gultekin, F., Ozturk, M., & Akdogan, M. (2000). The effect of organophosphate insecticide chlorpyrifos-ethyl on lipid 449 peroxidation and antioxidant enzymes (in vitro). Archives of Toxicology, 74(9), 533-538. 450 https://doi.org/10.1007/s002040000167
- Hamid, A., Yaqub, G., Ahmed, S. R., & Aziz, N. (2017). Assessment of human health risk associated with the presence of
 pesticides in chicken eggs. *Food Science and Technology (Brazil)*, *37*(3), 378–382. https://doi.org/10.1590/1678453 457x.11616

Hernández, A. F., Parrón, T., Tsatsakis, A. M., Requena, M., Alarcón, R., & López-Guarnido, O. (2013). Toxic effects of
pesticide mixtures at a molecular level: Their relevance to human health. *Toxicology*, 307, 136–145.
https://doi.org/10.1016/j.tox.2012.06.009

- Holden, A. v. (1972). The Effects of Pesticides on Life in Fresh Waters Author (s): Proceedings of the Royal Society of
 London . Series B , Biological Sciences , Mar . 21 , 1972 , Vol . 180 , No . 1061 , A Discussion on Freshwater and
 Estaurine Studie. *Biological Sciences*, 180(1061), 383–394.
- Houbraken, M., Spranghers, T., de Clercq, P., Cooreman-Algoed, M., Couchement, T., de Clercq, G., Verbeke, S., &
 Spanoghe, P. (2016). Pesticide contamination of Tenebrio molitor (Coleoptera: Tenebrionidae) for human consumption. *Food Chemistry*, 201, 264–269. https://doi.org/10.1016/j.foodchem.2016.01.097
- Islam, M. A., Nurul Amin, S. M., Brown, C. L., Juraimi, A. S., Uddin, M. K., & Arshad, A. (2022). Determination of the
 Most Efficient Household Technique for the Reduction of Pesticide Residues from Raw Fish Muscles. *Foods*, 11(9).
 https://doi.org/10.3390/foods11091254
- Jiang, J., Zhang, C., Wang, L., Wang, X., He, H., Wu, S., & Zhao, X. (2022). Insights into the combined effects of
 environmental concentration of difenoconazole and tebuconazole on zebrafish early life stage. *Science of The Total Environment*, 830, 154687. https://doi.org/10.1016/j.scitotenv.2022.154687
- Kim, L., Lee, D., Cho, H. K., & Choi, S. D. (2019). Review of the QuEChERS method for the analysis of organic pollutants:
 Persistent organic pollutants, polycyclic aromatic hydrocarbons, and pharmaceuticals. *Trends in Environmental Analytical Chemistry*, 22(e00063). https://doi.org/10.1016/j.teac.2019.e00063
- 472 Kortenkamp, A. (2007). Ten years of mixing cocktails: A review of combination effects of endocrine-disrupting chemicals.
 473 In *Environmental Health Perspectives* (Vol. 115, Issue SUPPL1, pp. 98–105). https://doi.org/10.1289/ehp.9357
- 474 Kumari, K. (2020). Pesticides toxicity in fishes : A review. Journal of Entomology and Zology Studies, 8(5), 1640–1642.
- Kushwaha, M., Verma, S., & Chatterjee, S. (2016). Profenofos, an Acetylcholinesterase-Inhibiting Organophosphorus
 Pesticide: A Short Review of Its Usage, Toxicity, and Biodegradation. *Journal of Environmental Quality*, 45(5),
 1478–1489. https://doi.org/10.2134/jeq2016.03.0100
- 478 Lengai, G. M. W., Muthomi, J. W., & Mbega, E. R. (2020). Phytochemical activity and role of botanical pesticides in pest
- 479 management for sustainable agricultural crop production. In *Scientific African* (Vol. 7, Issue e00239). Elsevier B.V.
 480 https://doi.org/10.1016/j.sciaf.2019.e00239
- 481 Lewis, K. A., Tzilivakis, J., Warner, D. J., & Green, A. (2016). An international database for pesticide risk assessments
 482 and management. *Human and Ecological Risk Assessment*, 22(4), 1050–1064.
 483 https://doi.org/10.1080/10807039.2015.1133242
- 484 Lozowicka, B. (2015). Health risk for children and adults consuming apples with pesticide residue. *Science of the Total* 485 *Environment*, 502, 184–198. https://doi.org/10.1016/j.scitotenv.2014.09.026
- 486 Macirella, R., Curcio, V., Ahmed, A. I. M., Pellegrino, D., & Brunelli, E. (2022). Effect of short-term exposure to low
 487 concentration of tebuconazole: morphological, histometric and functional modifications in Danio rerio liver.
 488 *European Zoological Journal*, 89(1), 324–338. https://doi.org/10.1080/24750263.2022.2043469
- 489 MacKay, D., & Fraser, A. (2000). Bioaccumulation of persistent organic chemicals: Mechanisms and models.
 490 *Environmental Pollution*, 110(3), 375–391. https://doi.org/10.1016/S0269-7491(00)00162-7

- 491 Mahmood, A., & Malik, R. N. (2014). Human health risk assessment of heavy metals via consumption of contaminated
 492 vegetables collected from different irrigation sources in Lahore, Pakistan. *Arabian Journal of Chemistry*, 7(1), 91–
 493 99. https://doi.org/10.1016/j.arabjc.2013.07.002
- 494 Manuelmolina-Ruiz, J., Cieslik, E., Cieslik, I., & Walkowska, I. (2014). Determination of pesticide residues in fish tissues
 495 by modified QuEChERS method and dual-d-SPE clean-up coupled to gas chromatography–mass spectrometry.
 496 *Environmental Science and Pollution Research*, 22(1), 369–378. https://doi.org/10.1007/s11356-014-3361-2
- Marchand, M. J., Pieterse, G. M., & Barnhoorn, I. E. J. (2010). Sperm motility and testicular histology as reproductive
 indicators of fish health of two feral fish species from a currently DDT sprayed area, South Africa. *Journal of Applied Ichthyology*, 26(5), 707–714. https://doi.org/10.1111/j.1439-0426.2010.01558.x
- Matos, P., Fontaínhas-Fernandes, A., Peixoto, F., Carrola, J., & Rocha, E. (2007). Biochemical and histological hepatic
 changes of Nile tilapia Oreochromis niloticus exposed to carbaryl. *Pesticide Biochemistry and Physiology*, 89(1),
 73–80. https://doi.org/10.1016/j.pestbp.2007.03.002
- Mengistie, B. T., Mol, A. P. J., & Oosterveer, P. (2016). Private Environmental Governance in the Ethiopian Pesticide
 Supply Chain: Importation, Distribution and Use. NJAS Wageningen Journal of Life Sciences, 76, 65–73.
 https://doi.org/10.1016/j.njas.2015.11.005
- 506 Mhadhbi, L., & Beiras, R. (2012). Acute toxicity of seven selected pesticides (alachlor, atrazine, dieldrin, diuron,
 507 pirimiphos-methyl, chlorpyrifos, diazinon) to the marine fish (turbot, Psetta maxima). *Water, Air, and Soil Pollution*,
 508 223(9), 5917–5930. https://doi.org/10.1007/s11270-012-1328-9
- 509 Müller, R., Shinn, C., Waldvogel, A. M., Oehlmann, J., Ribeiro, R., & Moreira-Santos, M. (2019). Long-term effects of 510 the fungicide pyrimethanil on aquatic primary producers in macrophyte-dominated outdoor mesocosms in two 511 665, 982-994. European ecoregions. of the Total Science Environment, 512 https://doi.org/10.1016/j.scitotenv.2019.02.050
- 513 Olette, R., Couderchet, M., Biagianti, S., & Eullaffroy, P. (2008). Toxicity and removal of pesticides by selected aquatic
 514 plants. *Chemosphere*, 70(8), 1414–1421. https://doi.org/10.1016/j.chemosphere.2007.09.016
- Patnaik, L., & Patra, ; (2006). Haemoatopoietic Alterations induced by Carbaryl in Clarias batrachus (LINN). J. Appl. Sci. *Environ. Mgt.*, 10(3), 5–7. www.bioline.org.br/ja
- Polat, A., Polat, S., Simsek, A., Kurt, T. T., & Ozyurt, G. (2018). Pesticide residues in muscles of some marine fish species
 and seaweeds of Iskenderun Bay (Northeastern Mediterranean), Turkey. *Environmental Science and Pollution Research*, 25(4), 3756–3764. https://doi.org/10.1007/s11356-017-0756-x
- Rasool, S., Rasool, T., & Gani, K. M. (2022). A review of interactions of pesticides within various interfaces of intrinsic
 and organic residue amended soil environment. In *Chemical Engineering Journal Advances* (Vol. 11). Elsevier B.V.
 https://doi.org/10.1016/j.ceja.2022.100301
- Ribera, D., Narbonne, J. F., Arnaud, C., & Saint-Denis, M. (2001). Biochemical responses of the earthworm Eisenia fetida
 andrei exposed to contaminated arti®cial soil, effects of carbaryl. *Soil Biology & Biochemistry*, *33*, 1123–1130.
 www.elsevier.com/locate/soilbio

- 526 Rosseland, B. O., Massabuau, J.-C., Grimalt, J., Hofer, R., Lackner, R., Raddum, G., Rognerud, S., & Vives, I. (2001).
- 527 Fish Ecotoxicology, The EMERGE Fish Sampling Manual for Live Fish. The EMERGE project (European mountain
 528 lake Ecosystem: regionalisation, diaGnostic and socio-economics valuation). 1–7.
- 529 Sabzevari, S., & Hofman, J. (2022). A worldwide review of currently used pesticides' monitoring in agricultural soils. In 530 Science of the Total Environment (Vol. 812. Issue 152344). Elsevier B.V. 531 https://doi.org/10.1016/j.scitotenv.2021.152344
- Sarkar, S., Dias Bernardes Gil, J., Keeley, J., Möhring, N., & Jansen, K. (2021). *The use of pesticides in developing countries and their impact on health and the right to food*□: *study.* (the European union).
 https://doi.org/10.2861/28995
- Araújo, C. V. M., Shinn, C., Müller, R., Moreira-Santos, M., Shinn, C., Espíndola, E. L., & Ribeiro, R. (2012). Aquatic
 ecotoxicity of the fungicide pyrimethanil: Effect profile under optimal and thermal stress conditions. *Environmental Pollution*, *168*, 161–169. https://doi.org/10.1016/j.envpol.2012.04.020
- Sharma, A., Kumar, V., Shahzad, B., Tanveer, M., Sidhu, G. P. S., Handa, N., Kohli, S. K., Yadav, P., Bali, A. S., Parihar,
 R. D., Dar, O. I., Singh, K., Jasrotia, S., Bakshi, P., Ramakrishnan, M., Kumar, S., Bhardwaj, R., & Thukral, A. K.
 (2019). Worldwide pesticide usage and its impacts on ecosystem. In *SN Applied Sciences* (Vol. 1, Issue 11). Springer
 Nature. https://doi.org/10.1007/s42452-019-1485-1
- 542 Sibbing, F. A., & Nagelkerke, L. A. J. (2001). Resource partitioning by Lake Tana barbs predicted from fish morphometrics Fish 543 Biology and and prey characteristics. Reviews in Fisheries, 10(4), 393-437. 544 https://doi.org/10.1023/A:1012270422092
- Sifakis, S., Androutsopoulos, V. P., Tsatsakis, A. M., & Spandidos, D. A. (2017). Human exposure to endocrine disrupting
 chemicals: effects on the male and female reproductive systems. *Environmental Toxicology and Pharmacology*, *51*,
 56–70. https://doi.org/10.1016/j.etap.2017.02.024
- 548 Smith, L. W. (1982). Wildlife toxicity studies with Oxamyl. Environmental Toxicology and Chemistry, I, 157–165.
- Srivastava, P., Singh, A., & Pandey, A. K. (2016). Pesticides toxicity in fishes: Biochemical, Physiological and Genotoxic
 aspects. *Biochem. Cell. Arch.*, 16(2), 199–218. www.connectjournals.com/bca
- 551 Teklu, B. M., Adriaanse, P. I., ter Horst, M. M. S., Deneer, J. W., & van den Brink, P. J. (2015). Surface water risk 552 assessment of pesticides in Ethiopia. Science the Total Environment, 508, 566-574. of 553 https://doi.org/10.1016/j.scitotenv.2014.11.049
- Teng, M., Zhu, W., Wang, D., Qi, S., Wang, Y., Yan, J., Dong, K., Zheng, M., & Wang, C. (2018). Metabolomics and
 transcriptomics reveal the toxicity of difenoconazole to the early life stages of zebrafish (Danio rerio). *Aquatic Toxicology*, *194*, 112–120. https://doi.org/10.1016/j.aquatox.2017.11.009
- 557 Tesfahun, A., & Alebachew, S. (2023). Food and feeding habits of the Nile tilapia Oreochromis niloticus (Linnaeus, 1758) 558 from Ribb reservoir, Lake Tana sub-basin, Ethiopia. Cogent Food Agriculture, æ 9(1). 559 https://doi.org/10.1080/23311932.2023.2212457
- Tournas, V. H. (2005). Spoilage of vegetable crops by bacteria and fungi and related health hazards. *Critical Reviews in Microbiology*, *31*(1), 33–44. https://doi.org/10.1080/10408410590886024

- US EPA. (2003). Revised EFED Risk Assessment of Carbaryl in Support of the Reregistration Eligibility Decision RED.
 https://www3.epa.gov/pesticides/endanger/litstatus/effects/carb-riskass.pdf
- 564USEPA.(2007).PesticidefactsheetforFlazasulfuron.565https://www3.epa.gov/pesticides/chemsearch/regactions/registration/fsPC-11910117-May-07.pdf
- 566 Vignet, C., Cappello, T., Fu, Q., Lajoie, K., de Marco, G., Clérandeau, C., Mottaz, H., Maisano, M., Hollender, J., Schirmer, 567 K., & Cachot, J. (2019). Imidacloprid induces adverse effects on fish early life stages that are more severe in Japanese 568 medaka (Oryzias latipes) than in zebrafish (Danio rerio). Chemosphere. 225. 470-478. 569 https://doi.org/10.1016/j.chemosphere.2019.03.002
- 570 Vonesh, J. R., & Kraus, J. M. (2009). Pesticide alters habitat selection and aquatic community composition. *Oecologia*, 571 *160*(2), 379–385. https://doi.org/10.1007/s00442-009-1301-5
- 572 WHO and FAO. (2019). Global situation of pesticide management in agriculture and public health.
 573 http://www.fao.org/3/ca7032en/ca7032en.pdf
- Witczak, A. (2009). Effect of heat treatment on organochlorine pesticide residues in selected fish species. *Pol. J. Food Nutr. Sci, 59*(3), 231–235. http://journal.pan.olsztyn.pl
- Worku, S. A., Dubale, Y. M., Woldemariam, E. D., & Eklo, O. M. (2022). Levels of organochlorine pesticides in five
 species of fish from Lake Ziway, Ethiopia. *Scientific African*, e01252. https://doi.org/10.1016/j.sciaf.2022.e01252
- 578 Yohannes, Y. B., Ikenaka, Y., Nakayama, S. M. M., Mizukawa, H., & Ishizuka, M. (2017). DDTs and other organochlorine
 579 pesticides in tissues of four bird species from the Rift Valley region, Ethiopia. *Science of the Total Environment*,
 580 *574*, 1389–1395. https://doi.org/10.1016/j.scitotenv.2016.08.056
- 581 Yohannes, Y. B., Ikenaka, Y., Saengtienchai, A., Watanabe, K. P., Nakayama, S. M. M., & Ishizuka, M. (2014). 582 Concentrations and human health risk assessment of organochlorine pesticides in edible fish species from a Rift 583 Ziway, Ethiopia. Vallev lake-Lake Ecotoxicology and Environmental Safety, 106. 95-101. https://doi.org/10.1016/j.ecoenv.2014.04.014 584 R