









Evidence-based guideline: unexplained infertility[†]

The Guideline Group on Unexplained Infertility, D. Romualdi ^{1,*}, B. Ata ^{2,3}, S. Bhattacharya⁴, E. Bosch ⁵, M. Costello^{6,7}, K. Gersak⁸, R. Homburg⁹, M. Mincheva ¹⁰, R.J. Norman ^{7,11}, T. Piltonen ¹², S. Dos Santos-Ribeiro ¹³, D. Scicluna¹⁴, S. Somers¹⁵, S.K. Sunkara¹⁶, H.R. Verhoeve¹⁷, and N. Le Clef ¹⁸

¹Department of Woman and Child Health and Public Health, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Rome, Italy

²Department of Obstetrics and Gynaecology, Koc University, Istanbul, Turkey

³ART Fertility Clinics, Dubai, United Arab Emirates

⁴School of Medicine, Medical Sciences & Nutrition, University of Aberdeen, Aberdeen, UK

⁵IVI-RMA Valencia, Valencia, Spain

⁶University of New South Wales, Sydney, Australia

⁷NHMRC Centre of Research Excellence Women's Health in Reproductive Life (WHiRL), Monash University, Melbourne, Australia

⁸Department of Obstetrics and Gynaecology, Faculty of Medicine, University Medical Centre Ljubljana, University of Ljubljana, Ljubljana, Slovenia

⁹Liverpool Womens' Hospital, Hewitt Fertility Centre, Liverpool, UK

¹⁰Centre for Tumour Microenvironment, Barts Cancer Institute, Queen Mary University of London, London, UK

¹¹The Robinson Research Institute The University of Adelaide, Adelaide, Australia

¹²Department of Obstetrics and Gynaecology, Reproductive Endocrinology and IVF Unit, PEDEGO Research Unit, Medical Research Centre, Oulu University Hospital, University of Oulu, Oulu, Finland

¹³IVI-RMA Lisboa, Lisbon, Portugal


¹⁴Gudja, Malta

¹⁵Department of Reproductive Medicine, Ghent University Hospital, Ghent, Belgium

¹⁶King's College London, London, UK

¹⁷Department of Gynaecology, OLVG, Amsterdam, The Netherlands

¹⁸European Society of Human Reproduction and Embryology, Grimbergen, Belgium

*Correspondence address. ESHRE Central Office, BXL7—Building 1, Nijverheidslaan 3, B-1853 Strombeek-Bever, Belgium. E-mail: guidelines@eshre.eu  <https://orcid.org/0000-0002-5236-2586>

[†]ESHRE Pages content is not externally peer reviewed. The manuscript has been approved by the Executive Committee of ESHRE.

ABSTRACT

STUDY QUESTION: What is the recommended management for couples presenting with unexplained infertility (UI), based on the best available evidence in the literature?

SUMMARY ANSWER: The evidence-based guideline on UI makes 52 recommendations on the definition, diagnosis, and treatment of UI.

WHAT IS KNOWN ALREADY: UI is diagnosed in the absence of any abnormalities of the female and male reproductive systems after 'standard' investigations. However, a consensual standardization of the diagnostic work-up is still lacking. The management of UI is traditionally empirical. The efficacy, safety, costs, and risks of treatment options have not been subjected to robust evaluation.

STUDY DESIGN, SIZE, DURATION: The guideline was developed according to the structured methodology for ESHRE guidelines. Following formulation of key questions by a group of experts, literature searches, and assessments were undertaken. Papers written in English and published up to 24 October 2022 were evaluated.

PARTICIPANTS/MATERIALS, SETTING, METHODS: Based on the available evidence, recommendations were formulated and discussed until consensus was reached within the guideline development group (GDG). Following stakeholder review of an initial draft, the final version was approved by the GDG and the ESHRE Executive Committee.

MAIN RESULTS AND THE ROLE OF CHANCE: This guideline aims to help clinicians provide the best care for couples with UI. As UI is a diagnosis of exclusion, the guideline outlined the basic diagnostic procedures that couples should/could undergo during an infertility work-up, and explored the need for additional tests. The first-line treatment for couples with UI was deemed to be IUI in combination with ovarian stimulation. The place of additional and alternative options for treatment of UI was also evaluated. The GDG made 52 recommendations on diagnosis and treatment for couples with UI. The GDG formulated 40 evidence-based recommendations—of which 29 were formulated as strong recommendations and 11 as weak—10 good practice points and two research only recommendations. Of the evidence-based recommendations, none were supported by high-quality evidence, one by moderate-quality evidence, nine by low-quality evidence, and 31 by very low-quality evidence. To support future research in UI, a list of research recommendations was provided.

Received: June 19, 2023. Editorial decision: July 5, 2023.

© The Author(s) 2023. Published by Oxford University Press on behalf of European Society of Human Reproduction and Embryology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

LIMITATIONS, REASONS FOR CAUTION: Most additional diagnostic tests and interventions in couples with UI have not been subjected to robust evaluation. For a large proportion of these tests and treatments, evidence was very limited and of very low quality. More evidence is required, and the results of future studies may result in the current recommendations being revised.

WIDER IMPLICATIONS OF THE FINDINGS: The guideline provides clinicians with clear advice on best practice in the care of couples with UI, based on the best evidence currently available. In addition, a list of research recommendations is provided to stimulate further studies in the field. The full guideline and a patient leaflet are available in www.eshre.eu/guideline/UI.

STUDY FUNDING/COMPETING INTEREST(S): The guideline was developed by ESHRE, who funded the guideline meetings, literature searches, and dissemination of the guideline in collaboration with the Monash University led Australian NHMRC Centre of Research Excellence in Women's Health in Reproductive Life (CREWHIRL). The guideline group members did not receive any financial incentives; all work was provided voluntarily. D.R. reports honoraria from IBSA and Novo Nordisk. B.A. reports speakers' fees from Merck, Gedeon Richter, Organon and Intas Pharma; is part of the advisory board for Organon Turkey and president of the Turkish Society of Reproductive Medicine. S.B. reports speakers' fees from Merck, Organon, Ferring, the Obstetric and Gynaecological Society of Singapore and the Taiwanese Society for Reproductive Medicine; editor and contributing author, Reproductive Medicine for the MRCOG, Cambridge University Press; is part of the METAFOR and CAPE trials data monitoring committee. E.B. reports research grants from Roche diagnostics, Gedeon Richter and IBSA; speaker's fees from Merck, Ferring, MSD, Roche Diagnostics, Gedeon Richter, IBSA; E.B. is also a part of an Advisory Board of Ferring Pharmaceuticals, MSD, Roche Diagnostics, IBSA, Merck, Abbott and Gedeon Richter. M.M. reports consulting fees from Mojo Fertility Ltd. R.J.N. reports research grant from Australian National Health and Medical Research Council (NHMRC); consulting fees from Flinders Fertility Adelaide, VinMec Hospital Hanoi Vietnam; speaker's fees from Merck Australia, Cadilla Pharma India, Ferring Australia; chair clinical advisory committee Westmead Fertility and research institute MyDuc Hospital Vietnam. T.P. is a part of the Research Council of Finland and reports research grants from Roche Diagnostics, Novo Nordics and Sigrid Juselius foundation; consulting fees from Roche Diagnostics and organon; speaker's fees from Gedeon Richter, Roche, Exeltis, Organon, Ferring and Korento patient organization; is a part of NFOG, AE-PCOS society and several Finnish associations. S.S.R. reports research grants from Roche Diagnostics, Organon, Theramex; consulting fees from Ferring Pharmaceuticals, MSD and Organon; speaker's fees from Ferring Pharmaceuticals, MSD/Organon, Besins, Theramex, Gedeon Richter; travel support from Gedeon Richter; S.S.R. is part of the Data Safety Monitoring Board of TTRANSPORT and deputy of the ESHRE Special Interest Group on Safety and Quality in ART; stock or stock options from IVI Lisboa, Clínica de Reprodução assistida Lda; equipment/medical writing/gifts from Roche Diagnostics and Ferring Pharmaceuticals. S.K.S. reports speakers' fees from Merck, Ferring, MSD, Pharmasure. HRV reports consulting and travel fees from Ferring Pharmaceuticals. The other authors have nothing to disclose.

DISCLAIMER: This guideline represents the views of ESHRE, which were achieved after careful consideration of the scientific evidence available at the time of preparation. In the absence of scientific evidence on certain aspects, a consensus between the relevant ESHRE stakeholders has been obtained.

Adherence to these clinical practice guidelines does not guarantee a successful or specific outcome, nor does it establish a standard of care. Clinical practice guidelines do not replace the need for application of clinical judgment to each individual presentation, nor variations based on locality and facility type.

ESHRE makes no warranty, express or implied, regarding the clinical practice guidelines and specifically excludes any warranties of merchantability and fitness for a particular use or purpose. (Full disclaimer available at www.eshre.eu/guidelines.)

Keywords: unexplained infertility / guideline / evidence-based / medically assisted reproduction / IUI / IVF / pregnancy / ESHRE / ART

Introduction

Approximately 30% of infertile couples are considered to experience 'unexplained infertility' (UI) (2019, 2020). This controversial diagnosis is made when no abnormalities of the female and male reproductive systems are identified. UI is inevitably a diagnosis by exclusion, following otherwise 'standard' investigations. However, a consensual standardization of the diagnostic work-up is still lacking. The International Committee for Monitoring Assisted Reproductive Technologies (ICMART) defined UI as 'infertility in couples with apparently normal ovarian function, fallopian tubes, uterus, cervix and pelvis and with adequate coital frequency; and apparently normal testicular function, genito-urinary anatomy and a normal ejaculate. The potential for this diagnosis is dependent upon the methodologies used and/or those methodologies available' (Zegers-Hochschild et al., 2017).

The proportion of couples with UI is related to the extent of diagnostic examination performed to uncover putative causes for unsuccessful attempts at pregnancy (ESHRE Capri Workshop Group, 2004). Furthermore, the criteria for labelling specific features as 'normal' are heterogeneous. Finally, apart from the clearly recognized causes of infertility, several undetectable defects in the reproductive process might prevent conception.

In the absence of an identified cause, the management of UI is traditionally empirical. The efficacy, safety, costs, and risks of treatment options have not been subjected to robust evaluation.

Materials and methods

The guideline was developed according to a well-documented methodology that is universal to ESHRE guidelines (Vermeulen et al., 2019). The guideline development group (GDG) was composed of members of the ESHRE Special Interest Group (SIG) Reproductive Endocrinology, SIG Andrology, SIG Safety and Quality in ART, SIG Nurses and Midwives, and a patient representative from Fertility Europe. This guideline was developed in collaboration with Monash University NHMRC Centre for Research Excellence in Women's Reproductive Health.

In short, 21 key questions were formulated by the GDG, of which four were answered as narrative questions, and 17 as PICO (Patient, Intervention, Comparison, Outcome) questions. For each PICO question, databases (PUBMED/MEDLINE and Cochrane library) were searched from inception to 24 October 2022, for publications written in English. From the literature searches, studies were selected on the basis of their relevance

to the PICO questions, assessed for quality, and summarized in evidence tables (Supplementary File S1). At GDG meetings, the evidence and draft recommendations were presented by the assigned GDG member and discussed until consensus was reached within the group. Each recommendation was labelled as strong or conditional and a grade was assigned based on the strength of the supporting evidence (High ⊕⊕⊕⊕, Moderate ⊕⊕⊕○, Low ⊕⊕○○, Very low ⊕○○○). Good practice points (GPPs) based on clinical expertise were added, where relevant, to clarify the recommendations or to provide further practical advice. Two 'research only' recommendations were also made for tests which should only be applied within the context of research.

Strong recommendations should be applied to most patients, while weak recommendations require discussion and shared decision-making.

For some of the narrative questions, a similar literature search was conducted. Collected data were summarized in a narrative summary and conclusions were formulated.

The draft guideline along with an invitation to participate in the stakeholder review were published on the ESHRE website between 12 December 2022 and 30 January 2023. All comments were processed by the GDG, either by adapting the content of the guideline and/or by responding to the reviewer. The review process was summarized in the review report, which is published on the ESHRE website (www.eshre.eu/Guidelines). The list of experts who contributed to the stakeholder review is included in Supplementary File S2. Overall, 31% of the 260 comments resulted in an amendment to the guideline text. A flowchart on diagnosis and management of UI is also available on the ESHRE website.

This guideline will be considered for update 4 years after publication, with an intermediate assessment of the need for updating 2 years after publication.

Results

Key questions and recommendations

The current document summarizes all the key questions and the recommendations from the guideline on 'Unexplained Infertility'. Further background information and the supporting evidence for each recommendation can be found in the full version of the guideline available at www.eshre.eu/guideline/UI.

Definition

The GDG defines UI as follows: infertility in couples with apparently normal ovarian function, fallopian tubes, uterus, cervix and pelvis, age ≤40 years and with adequate coital frequency; and apparently normal testicular function, genito-urinary anatomy, and a normal ejaculate.

As per the ICMART definition of infertility, couples should have at least 12 months of regular, unprotected sexual intercourse before investigations are initiated.

The GDG recommends routinely taking a medical, reproductive and sexual history from both the male and female partner.

The GDG considers a regular menstrual cycle to be 24–38 days, up to 8 days in duration, and shortest to longest cycle variation of <7–9 days (Munro et al., 2018).

The GDG recommends at least one basic semen examination, according to World Health Organization (WHO) criteria,

performed by a laboratory which subscribes to an external quality control programme. If the result from first basic semen analysis is below the lower fifth percentile reference limit as per WHO criteria (6th edition), a second analysis should be performed after a 3-month interval.

Diagnosis

Confirmation of ovulation

In women with regular menstrual cycles, tests for confirmation of ovulation are not routinely recommended.	GPP
In women with regular menstrual cycles, if confirmation of ovulation is warranted, tests such as urinary LH measurements, ultrasound monitoring or mid-luteal progesterone measurement can be used (Gregoriou et al., 1990; Bischof et al., 1991; Martinez et al., 1991; Guermandi et al., 2001)	Conditional ⊕○○○

Oocyte/corpus luteum quality

In women with regular menstrual cycles, it is suggested not to routinely measure midluteal serum progesterone levels (Hull et al., 1982).	Conditional ⊕○○○
In women investigated for infertility, endometrial biopsy for histological examination is not recommended in the absence of other indications (Coutifaris et al., 2004; Edi-Osagie et al., 2004).	Strong ⊕⊕○○

Ovarian reserve

In women with regular menstrual cycles, ovarian reserve testing is not required to identify the aetiology of infertility or to predict the probability of spontaneous conception over 6–12 months (Scott et al., 1993; Rosen et al., 2011; Hagen et al., 2012; Casadei et al., 2013; Murto et al., 2013; Hvidman et al., 2016; Depmann et al., 2017; Greenwood et al., 2017; Steiner et al., 2017; Yücel et al., 2018; Nguyen et al., 2022).	Strong ⊕⊕○○
--	----------------

Tubal factor

Hysterosalpingo-contrast-sonography (HyCoSy) and hysterosalpingography (HSG) are valid tests for tubal patency compared to laparoscopy and chromopertubation (Broeze et al., 2011; Wang and Qian, 2016; Alcázar et al., 2020).	Strong ⊕⊕⊕○
HSG and HyCoSy are comparable in diagnostic capacity, thus selection of the technique depends on the preference of the clinician and the patient.	GPP
Chlamydia antibody testing for tubal patency could be considered a non-invasive test to differentiate between patients at low and at high risk for tubal occlusion (Mol et al., 1997).	Conditional ⊕○○○
In patients at high risk for tubal abnormality, visual demonstration of tubal patency is necessary.	GPP

Uterine factor

Ultrasound, preferably 3D, is recommended to exclude uterine anomalies in women with unexplained infertility (Jurkovic et al., 1995; Caliskan et al., 2010; Ludwin et al., 2013).	Strong ⊕○○○
MRI is not recommended as a first-line test to confirm a normal uterine structure and anatomy in women with unexplained infertility.	Strong ⊕○○○
If ultrasound assessment of the uterine cavity is normal, no further evaluation is needed (Fatemi et al., 2010; Almog et al., 2011; Bakas et al., 2014; Makled et al., 2014; Yang et al., 2019).	Strong ⊕○○○

Laparoscopy

Routine diagnostic laparoscopy is not recommended for the diagnosis of unexplained infertility (Tanahatog et al., 2003, 2005; Lavy et al., 2004).	Strong ⊕○○○
---	----------------

Cervical/vaginal investigations

The post-coital test is not recommended in couples with unexplained infertility (Oei et al., 1995).	Strong ⊕⊕○○
Vaginal microbiota testing could be considered in couples with unexplained infertility only in a research setting.	Research only

Male genito-urinary anatomy

Testicular imaging is not recommended when semen analysis according to WHO criteria is normal (Lotti et al., 2021).	Strong ⊕○○○
---	----------------

Male additional tests

Testing for anti-sperm antibodies in the semen is not recommended when semen analysis according to WHO criteria is normal (Ayvaliotis et al., 1985; Lähteenmäki, 1993; Rajah et al., 1993; Pagidas et al., 1994; Lähteenmäki et al., 1995; Vazquez-Levin et al., 1997; Bozhedomov et al., 2015; Barbonetti et al., 2020).	Strong ⊕○○○
Testing for sperm DNA fragmentation is not recommended when semen analysis according to WHO criteria is normal (O'Neill et al., 2018; Borges et al., 2019; Repalle et al., 2022).	Strong ⊕○○○
Sperm chromatin condensation test is not recommended when semen analysis according to WHO criteria is normal.	Strong ⊕○○○
Sperm aneuploidy screening is not recommended when semen analysis according to WHO criteria is normal.	Strong ⊕○○○
Serum hormonal testing is not required when semen analysis according to WHO criteria is normal.	Strong ⊕○○○
Human papillomavirus (HPV) testing of semen is not recommended when semen analysis according to WHO criteria is normal.	Strong ⊕○○○
Microbiology testing of semen is not recommended when semen analysis according to WHO criteria is normal.	Strong ⊕○○○

Systemic additional tests

Testing for anti-sperm antibodies in serum of either males or females with unexplained infertility is not recommended (Menge et al., 1982; Mardesic et al., 2000; Monem and Moalla, 2003; Yasin et al., 2016).	Strong ⊕○○○
Testing for coeliac disease in women with unexplained infertility can be considered (Tersigni et al., 2014).	Conditional ⊕⊕○○
Testing for thyroid antibody and other autoimmune conditions (apart from coeliac disease) in women with unexplained infertility is not recommended (Poppe et al., 2002; Abalovich et al., 2007; Kilic et al., 2008).	Strong ⊕○○○
TSH measurement is considered good practice in pre-conception care.	GPP
No additional thyroid evaluation in women is recommended if TSH is within the normal range (Duran et al., 2013; Unuane et al., 2013; Orouji Jokar et al., 2018; Rehman et al., 2020).	Strong ⊕○○○
Testing for thrombophilia in women with unexplained infertility is not recommended (Behjati et al., 2006; Bellver et al., 2008; Coulam and Jeyendran, 2009; Casadei et al., 2010; Fatini et al., 2012; Steinvil et al., 2012; Kydonopoulou et al., 2017; Milenkovic et al., 2020).	Strong ⊕○○○
Measurement of oxidative stress in semen of males with unexplained infertility should only be considered in the context of research.	Research only
Measurement of oxidative stress in women with unexplained infertility is not recommended (Veena et al., 2008; Pekel et al., 2015; Lazzarino et al., 2021; Şentürk et al., 2021).	Strong ⊕⊕○○
Genetic or genomic tests are currently not recommended in couples with unexplained infertility (Trková et al., 2000; Papanikolaou et al., 2005; Witkin et al., 2010; Sahmani et al., 2011; Vani et al., 2012; Suganya et al., 2015; Salas-Huetos et al., 2016; Rull et al., 2018; Ertosun et al., 2022).	Strong ⊕○○○
Testing for vitamin D deficiency in women is not recommended for diagnosis of unexplained infertility (Rudick et al., 2012; Lopes et al., 2017; Butts et al., 2019; Güngör et al., 2022; Ko et al., 2022).	Strong ⊕○○○
Prolactin testing in women is not recommended (Subramanian et al., 1997; Veena et al., 2008; Orouji Jokar et al., 2018; Qu et al., 2020).	Strong ⊕○○○
BMI measurement in women is considered good practice in pre-conception care.	GPP

Treatment**Expectant management**

IUI with ovarian stimulation is recommended as a first-line treatment for couples with unexplained infertility (Fisch et al., 1989; Bhattacharya et al., 2008; Pandian et al., 2015; Ayeleke et al., 2020).	Strong ⊕○○○
The GDG advises to base the decision to start active treatment on prognosis in couples with unexplained infertility.	GPP

Active treatment

IUI with ovarian stimulation is recommended as a first-line treatment for couples with unexplained infertility (Agarwal and Mittal, 2004; Bhattacharya et al., 2008; Ayeleke et al., 2020; Harira 2018; Ibrahim et al., 2012).	Strong ⊕○○○
To avoid multiple pregnancies and ovarian hyperstimulation syndrome (OHSS), care is needed by using gonadotrophin treatment only in a low-dose regimen with adequate monitoring.	GPP
IVF is probably not recommended over IUI with ovarian stimulation in couples with unexplained infertility (Pandian et al., 2015; Nandi et al., 2022).	Conditional ⊕○○○
It is expected that the decision to use IVF is individualized by patient characteristics such as age, duration of infertility, previous treatment, and previous pregnancy.	GPP
ICSI is not recommended over conventional IVF in couples with unexplained infertility (Bhattacharya et al., 2001; Foong et al., 2006; Dang et al., 2021).	Strong ⊕○○○

Mechanical–surgical procedures

Hysteroscopy for the detection and possible correction of intrauterine abnormalities not seen at routine imaging is not recommended (Casini et al., 2006; Seyam et al., 2015).	Strong ⊕⊕○○
HSG (i.e. tubal flushing) with an oil-soluble contrast medium is preferable over a water-soluble contrast medium. Risks and benefits of tubal flushing with oil-soluble contrast medium should be discussed with all couples with unexplained infertility (Wang et al., 2020).	Conditional ⊕⊕○○
Endometrial scratching should not be offered for unexplained infertility (Parsanezhad et al., 2013; Senocak et al., 2017; Ghuman et al., 2020; Jafarabadi et al., 2020; Yildiz et al., 2021; Wong et al., 2022).	Strong ⊕⊕○○

If incidentally minimal to mild endometriosis is found at laparoscopy, this is not further considered unexplained infertility by the GDG.

Alternative therapeutic approaches

Adjunct oral antioxidant therapy to women undergoing fertility treatment is probably not recommended (Showell et al., 2020).	Conditional ⊕○○○
Adjunct oral antioxidant therapy to males undergoing fertility treatment is probably not recommended.	Conditional ⊕○○○
Acupuncture in women is probably not recommended (Guven et al., 2020).	Conditional ⊕⊕○○
Inositol supplementation in women is probably not recommended (Montanino Oliva et al., 2020).	Conditional ⊕○○○
Psychological support, including psychotherapy, is recommended for patients when needed.	GPP
A healthy diet and regular exercise, supported by behavioural therapy when necessary, are recommended.	GPP

Quality of life

Healthcare professionals should be aware that	Conditional ⊕○○○
<ul style="list-style-type: none"> There is probably no difference in quality of life (QoL) between women with unexplained infertility versus women in couples with known causes of infertility, except when the cause of infertility is PCOS, where the QoL is lower. QoL is probably higher in men from a couple with unexplained infertility compared to men from a couple with known causes of infertility except when the cause of infertility is men with a partner with PCOS, then the men from a couple with unexplained infertility have a lower QoL (Kowalcek et al., 2001; Santoro et al., 2016; Warchol-Biedermann, 2021). 	

Discussion

The current paper presents the 52 recommendations on management of UI from the evidence-based guideline on 'Unexplained Infertility'. This guideline covers all aspects of the definition, diagnosis and treatment of couples with UI. The guideline was written by a multidisciplinary group of experts in reproductive endocrinology, reproductive surgery, and andrology, along with a nurse and a patient representative and developed in collaboration with the Monash University led NHMRC Centre of Research Excellence in Women's Health in Reproductive Life (CREWHIRL).

Notwithstanding the importance and relevance of the topic, research data on many key aspects are scarce. As a basis for the current guideline, a formal literature review was conducted. Most studies on diagnosis were old, with often incomplete reporting of methodology. Review of additional tests for establishing the diagnosis of UI was plagued by heterogeneity of the study population and lack of standardization of assays. The literature on the diagnosis of a putative male cause for unexplained infertility was complicated by the interchangeable use of the terms unexplained and idiopathic male infertility.

The recommendation against laparoscopy as a routine procedure in the diagnostic infertility work-up generated considerable debate during the stakeholder review. Therefore, this topic, which was extensively analysed during the previous meetings, was further reconsidered. The GDG agreed that HSG and HyCoSy/HyFoSy do not detect mild endometriosis, adhesions, or subtle tubal lesions. However, there are insufficient good quality data to suggest that clinically relevant diagnoses will be missed by omitting a laparoscopy in patients at low risk for tubal pathology. As evidence is lacking to justify routine laparoscopy for every patient with otherwise UI given possible surgical and anaesthesiological risks, the recommendation was retained. Nevertheless, clinicians are advised to counsel women at high risk for tubal pathology (a history of pelvic inflammatory disease, previous ectopic pregnancy) or endometriosis about the benefits and risks of laparoscopy.

The GDG received several comments on the relatively minor role of investigations of the male partner in the standard diagnostic work-up for UI. The literature on most of the possible additional tests proposed for the male partner in the last decades was reviewed in the guideline development process. Insufficient evidence was found to suggest the diagnostic benefit of these investigations in men with normal semen parameters according to WHO criteria (6th edition). The tests under scrutiny were characterized by limited capacity to discriminate between

couples who would benefit from a specific medically assisted reproduction (MAR) technique; inconsistent and heterogeneous cut-offs and unvalidated thresholds; lack of reliable predictive value in terms of reproductive outcomes; and lack of proven value in informing clinical decision making. Hence, recommendations in favour of their routine use in the initial evaluation of couples with UI were not adequately supported, taking into account their possible economic and psychological burden.

Nevertheless, the GDG acknowledges that the quality of data is generally very low and that the male factor is neglected in the scientific literature. It was decided to amend the former version of the guideline by underlining that more research is needed in this area. Re-focusing research efforts on addressing gaps in the understanding of male infertility, such as identifying new aetiological causes, clinical diagnostics, and MAR treatment options, will enable the development of more personalized therapeutic options to manage couple's infertility and improve reproductive outcomes. Furthermore, a statement was added on the importance to investigate the general and reproductive history of the male, with particular attention to sexual dysfunction. Should any abnormality emerge, physical examination, and appropriate investigations would be warranted. However, in these circumstances, the diagnosis of UI would no longer be applicable, and further specifications would fall outside the scope of the present guideline.

Very few high-quality randomized controlled trials were available to the GDG to make sound recommendations with regard to treatment of couples with unexplained infertility. The GDG also received criticism for not including a section on prognosis-based treatment in the guideline. This feedback was held in high regard and a new section was included in the guideline. In brief, it was reported that prognostic models, as well as patients' preference, can help the decision-making on a treatment plan in couples with UI. Overall, in the case of expectant management, the most important prognostic factors are age, duration of infertility, previous treatment, and previous pregnancies. It is, however, important to note that none of the currently available prediction models is fully evolved. Promising models are currently under development, however, they need to be implemented and validated before legitimately entering into common use.

Research gaps were detected in several areas, and the top three topics are documented in a list of recommendations for further research ([Supplementary File S3](#)).

Despite the limitations of guidelines in general, and the limitations in the evidence supporting the current guideline, the GDG is confident that this document will help best practice in the management of couples with UI.

Supplementary data

[Supplementary data](#) are available at *Human Reproduction* online.

Data availability

The data underlying this article are available in the article and in its [supplementary material](#).

Acknowledgements

The GDG would like to acknowledge the help of many clinicians and professional organizations who refereed the content of the guideline and submitted helpful comments to the draft version. Special thanks to the steering committee of the ESHRE SIG

Andrology for the feedback on the formulations of the key questions and the final draft of the guideline.

Authors' roles

D.R. chaired the GDG and hence fulfilled a leading role in collecting the evidence, writing the manuscript and dealing with reviewer comments. N.L.C., as methodological expert, performed all literature searches for the guideline, provided methodological support, and coordinated the guideline development. All other authors, listed in alphabetical order, as GDG members, contributed equally to the manuscript, by drafting key questions, synthesizing evidence, writing the different parts of the guideline, and discussing recommendations until consensus within the group was reached.

Funding

The study has no external funding; all costs for meetings were covered by ESHRE.

Conflict of interest

The guideline was developed by ESHRE, who funded the guideline meetings, literature searches, and dissemination of the guideline in collaboration with the Monash University led Australian NHMRC Centre of Research Excellence in Women's Health in Reproductive Life (CREWHIRL). The guideline group members did not receive any financial incentives; all work was provided voluntarily. D.R. reports honoraria from IBSA and Novo Nordisk. B.A. reports speakers' fees from Merck, Gedeon Richter, Organon and Intas Pharma; is part of the advisory board for Organon Turkey and president of the Turkish Society of Reproductive Medicine. S.B. reports speakers' fees from Merck, Organon, Ferring, the Obstetric and Gynaecological Society of Singapore and the Taiwanese Society for Reproductive Medicine; editor and contributing author, Reproductive Medicine for the MRCOG, Cambridge University Press; is a part of the METAFOR and CAPE trials data monitoring committee. E.B. reports research grants from Roche diagnostics, Gedeon Richter and IBSA; speaker's fees from Merck, Ferring, MSD, Roche Diagnostics, Gedeon Richter, IBSA; E.B. is also a part of an Advisory Board of Ferring Pharmaceuticals, MSD, Roche Diagnostics, IBSA, Merck, Abbott and Gedeon Richter. M.M. reports consulting fees from Mojo Fertility Ltd. R.J.N. reports research grant from Australian National Health and Medical Research Council (NHMRC); consulting fees from Flinders Fertility Adelaide, VinMec Hospital Hanoi Vietnam; speaker's fees from Merck Australia, Cadilla Pharma India, Ferring Australia; chair clinical advisory committee Westmead Fertility and research institute MyDuc Hospital Vietnam. T.P. is part of the Research Council of Finland and reports research grants from Roche Diagnostics, Novo Nordics and Sigrid Juselius foundation; consulting fees from Roche Diagnostics and organon; speaker's fees from Gedeon Richter, Roche, Exeltis, Organon, Ferring and Korento patient organization; is a part of NFOG, AE-PCOS society, and several Finnish associations. S.S.R. reports research grants from Roche Diagnostics, Organon, Theramex; consulting fees from Ferring Pharmaceuticals, MSD and Organon; speaker's fees from Ferring Pharmaceuticals, MSD/Organon, Besins, Theramex, Gedeon Richter; travel support from Gedeon Richter; S.S.R. is a part of the Data Safety Monitoring Board of TTRANSPORT and deputy of the ESHRE Special Interest Group on Safety and Quality in ART; stock or stock options from IVI Lisboa,

Clínica de Reprodução assistida Lda; equipment/medical writing/ gifts from Roche Diagnostics and Ferring Pharmaceuticals. S.K.S. reports speakers' fees from Merck, Ferring, MSD, Pharmasure. H.R.V. reports consulting and travel fees from Ferring Pharmaceuticals. The other authors have nothing to disclose.

References

- Infertility Workup for the Women's Health Specialist: ACOG Committee Opinion, Number 781. *Obstet Gynecol* 2019;**133**: e377–e384.
- Evidence-based treatments for couples with unexplained infertility: a guideline. *Fertil Steril* 2020;**113**:305–322.
- Abalovich M, Mitelberg L, Allami C, Gutierrez S, Alcaraz G, Otero P, Levalle O. Subclinical hypothyroidism and thyroid autoimmunity in women with infertility. *Gynecol Endocrinol* 2007;**23**:279–283.
- Agarwal S, Mittal S. A randomised prospective trial of intrauterine insemination versus timed intercourse in superovulated cycles with clomiphene. *Indian J Med Res* 2004;**120**:519–522.
- Alcázar JL, Martínez A, Duarte M, Welly A, Marín A, Calle A, Garrido R, Pascual MA, Guerriero S. Two-dimensional hysterosalpingo-contrast-sonography compared to three/four-dimensional hysterosalpingo-contrast-sonography for the assessment of tubal occlusion in women with infertility/subfertility: a systematic review with meta-analysis. *Hum Fertil (Camb)* 2020;**25**:43–55.
- Almog B, Shalom-Paz E, Shehata F, Ata B, Levin D, Holzer H, Tan SL. Saline instillation sonohysterography test after normal baseline transvaginal sonography results in infertility patients. Is it justified? *Gynecol Endocrinol* 2011;**27**:286–289.
- Ayeleke RO, Asseler JD, Cohlen BJ, Veltman-Verhulst SM. Intra-uterine insemination for unexplained subfertility. *Cochrane Database Syst Rev* 2020;**3**:Cd001838.
- Ayvaliotis B, Bronson R, Rosenfeld D, Cooper G. Conception rates in couples where autoimmunity to sperm is detected. *Fertil Steril* 1985;**43**:739–742.
- Bakas P, Hassiakos D, Grigoriadis C, Vlahos N, Liapis A, Gregoriou O. Role of hysteroscopy prior to assisted reproduction techniques. *J Minim Invasive Gynecol* 2014;**21**:233–237.
- Barbonetti A, Castellini C, D'Andrea S, Minaldi E, Totaro M, Francavilla S, Francavilla F. Relationship between natural and intrauterine insemination-assisted live births and the degree of sperm autoimmunisation. *Hum Reprod* 2020;**35**:1288–1295.
- Behjati R, Modarressi MH, Jeddi-Tehrani M, Dokoohaki P, Ghasemi J, Zarnani AH, Aarabi M, Memariani T, Ghaffari M, Akhondi MA. Thrombophilic mutations in Iranian patients with infertility and recurrent spontaneous abortion. *Ann Hematol* 2006;**85**:268–271.
- Bellver J, Soares SR, Alvarez C, Muñoz E, Ramírez A, Rubio C, Serra V, Remohí J, Pellicer A. The role of thrombophilia and thyroid autoimmunity in unexplained infertility, implantation failure and recurrent spontaneous abortion. *Hum Reprod* 2008;**23**:278–284.
- Bhattacharya S, Hamilton MP, Shaaban M, Khalaf Y, Seddler M, Ghobara T, Braude P, Kennedy R, Rutherford A, Hartshorne G et al. Conventional in-vitro fertilisation versus intracytoplasmic sperm injection for the treatment of non-male-factor infertility: a randomised controlled trial. *Lancet* 2001;**357**:2075–2079.
- Bhattacharya S, Harrild K, Mollison J, Wordsworth S, Tay C, Harrold A, McQueen D, Lyall H, Johnston L, Burrage J et al. Clomifene citrate or unstimulated intrauterine insemination compared with expectant management for unexplained infertility: pragmatic randomised controlled trial. *BMJ* 2008;**337**:a716.
- Bischof P, Bianchi PG, Campana A. Comparison of a rapid, quantitative and automated assay for urinary luteinizing hormone (LH), with an LH detection test, for the prediction of ovulation. *Hum Reprod* 1991;**6**:515–518.
- Borges E Jr, Zanetti BF, Setti AS, Braga D, Provenza RR, Iaconelli A Jr. Sperm DNA fragmentation is correlated with poor embryo development, lower implantation rate, and higher miscarriage rate in reproductive cycles of non-male factor infertility. *Fertil Steril* 2019;**112**:483–490.
- Bozhedomov VA, Nikolaeva MA, Ushakova IV, Lipatova NA, Bozhedomova GE, Sukhikh GT. Functional deficit of sperm and fertility impairment in men with antisperm antibodies. *J Reprod Immunol* 2015;**112**:95–101.
- Broeze KA, Opmeer BC, Van Geloven N, Coppus SF, Collins JA, Den Hartog JE, Van der Linden PJ, Marianowski P, Ng EH, Van der Steeg JW et al. Are patient characteristics associated with the accuracy of hysterosalpingography in diagnosing tubal pathology? An individual patient data meta-analysis. *Hum Reprod Update* 2011;**17**:293–300.
- Butts SF, Seifer DB, Koelper N, Senapati S, Sammel MD, Hoofnagle AN, Kelly A, Krawetz SA, Santoro N, Zhang H et al.; Eunice Kennedy Shriver National Institute of Child Health and Human Development Reproductive Medicine Network. Vitamin D deficiency is associated with poor ovarian stimulation outcome in PCOS but not unexplained infertility. *J Clin Endocrinol Metab* 2019;**104**:369–378.
- Caliskan E, Ozkan S, Cakiroglu Y, Sarisoy HT, Corakci A, Ozeren S. Diagnostic accuracy of real-time 3D sonography in the diagnosis of congenital Mullerian anomalies in high-risk patients with respect to the phase of the menstrual cycle. *J Clin Ultrasound* 2010;**38**:123–127.
- Casadei L, Manicuci C, Puca F, Madrigale A, Emidi E, Piccione E. Can anti-Müllerian hormone be predictive of spontaneous onset of pregnancy in women with unexplained infertility? *J Obstet Gynaecol* 2013;**33**:857–861.
- Casadei L, Puca F, Privitera L, Zamaro V, Emidi E. Inherited thrombophilia in infertile women: implication in unexplained infertility. *Fertil Steril* 2010;**94**:755–757.
- Casini ML, Rossi F, Agostini R, Unfer V. Effects of the position of fibroids on fertility. *Gynecol Endocrinol* 2006;**22**:106–109.
- Coulam CB, Jeyendran RS. Thrombophilic gene polymorphisms are risk factors for unexplained infertility. *Fertil Steril* 2009;**91**: 1516–1517.
- Coutifaris C, Myers ER, Guzick DS, Diamond MP, Carson SA, Legro RS, McGovern PG, Schlaff WD, Carr BR, Steinkampf MP et al.; NICHD National Cooperative Reproductive Medicine Network. Histological dating of timed endometrial biopsy tissue is not related to fertility status. *Fertil Steril* 2004;**112**:e116–e124.
- Dang VQ, Vuong LN, Luu TM, Pham TD, Ho TM, Ha AN, Truong BT, Phan AK, Nguyen DP, Pham TN et al. Intracytoplasmic sperm injection versus conventional in-vitro fertilisation in couples with infertility in whom the male partner has normal total sperm count and motility: an open-label, randomised controlled trial. *Lancet* 2021;**397**:1554–1563.
- Depmann M, Broer SL, Eijkemans MJC, van Rooij IAJ, Scheffer GJ, Heimensem J, Mol BW, Broekmans FJM. Anti-Müllerian hormone does not predict time to pregnancy: results of a prospective cohort study. *Gynecol Endocrinol* 2017;**33**:644–648.
- Duran B, Özlü T, Koç O, Eşitken C, Topçuoğlu A. Relationship of thyroid hormone levels and thyroid autoantibodies with early pregnancy loss and infertility. *J Obstet Gynaecol* 2013;**33**:862–864.
- Edi-Osagie EC, Seif MW, Aplin JD, Jones CJ, Wilson G, Lieberman BA. Characterizing the endometrium in unexplained and tubal factor infertility: a multiparametric investigation. *Fertil Steril* 2004;**82**: 1379–1389.

- Ertosun MG, Araci DG, Peker A, Uzuner SY, Toyulu A, Ozekinci M, Usta MF, Clark OA. Investigation of the relationship between reproductive disorders and chromosomal abnormalities in a large-scale, single-center 10-year retrospective study. *J Gynecol Obstet Hum Reprod* 2022;**51**:102467.
- ESHRE Capri Workshop Group. Diagnosis and management of the infertile couple: missing information. *Hum Reprod Update* 2004;**10**: 295–307.
- Fatemi HM, Kasius JC, Timmermans A, van Disseldorp J, Fauser BC, Devroey P, Broekmans FJ. Prevalence of unsuspected uterine cavity abnormalities diagnosed by office hysteroscopy prior to in vitro fertilization. *Hum Reprod* 2010;**25**:1959–1965.
- Fatini C, Conti L, Turillazzi V, Sticchi E, Romagnuolo I, Milanini MN, Cozzi C, Abbate R, Noci I. Unexplained infertility: association with inherited thrombophilia. *Thromb Res* 2012;**129**:e185–e188.
- Fisch P, Casper RF, Brown SE, Wrixon W, Collins JA, Reid RL, Simpson C. Unexplained infertility: evaluation of treatment with clomiphene citrate and human chorionic gonadotropin. *Fertil Steril* 1989;**51**:828–833.
- Foong SC, Fleetham JA, O'Keane JA, Scott SG, Tough SC, Greene CA. A prospective randomized trial of conventional in vitro fertilization versus intracytoplasmic sperm injection in unexplained infertility. *J Assist Reprod Genet* 2006;**23**:137–140.
- Ghuman NK, Raikar S, Singh P, Gothwal M, Yadav G. Improving reproductive outcomes of intrauterine insemination: Does endometrial scratch injury help? A randomised controlled trial. *Eur J Obstet Gynecol Reprod Biol* 2020;**253**:225–231.
- Greenwood EA, Cedars MI, Santoro N, Eisenberg E, Kao CN, Haisenleder DJ, Diamond MP, Huddleston HG; National Institutes of Health/Eunice Kennedy Shriver National Institute of Child Health and Human Development Cooperative Reproductive Medicine Network. Antimüllerian hormone levels and antral follicle counts are not reduced compared with community controls in patients with rigorously defined unexplained infertility. *Fertil Steril* 2017;**108**:1070–1077.
- Gregoriou O, Kassanos D, Vitoratos N, Papadias C, Zourlas PA. Clinical efficacy of LH-color: a new home ovulation test. *Int J Gynaecol Obstet* 1990;**32**:141–143.
- Guermandi E, Vegetti W, Bianchi MM, Uglietti A, Ragni G, Crosignani P. Reliability of ovulation tests in infertile women. *Obstet Gynecol* 2001;**97**:92–96.
- Güngör K, Güngör ND, Başar MM, Cengiz F, Erşahin SS, Çil K. Relationship between serum vitamin D levels semen parameters and sperm DNA damage in men with unexplained infertility. *Eur Rev Med Pharmacol Sci* 2022;**26**:499–505.
- Guyen PG, Cayir Y, Borekci B. Effectiveness of acupuncture on pregnancy success rates for women undergoing in vitro fertilization: a randomized controlled trial. *Taiwan J Obstet Gynecol* 2020;**59**: 282–286.
- Hagen CP, Vestergaard S, Juul A, Skakkebaek NE, Andersson AM, Main KM, Hjellund NH, Ernst E, Bonde JP, Anderson RA et al. Low concentration of circulating antimüllerian hormone is not predictive of reduced fecundability in young healthy women: a prospective cohort study. *Fertil Steril* 2012;**98**:1602–1608.e2.
- Harira M. Use of letrozole versus clomiphene-estradiol for treating infertile women with unexplained infertility not responding well to clomiphene alone, comparative study. *Middle East Fertil Soc J* 2018;**23**:384–387.
- Hull MG, Savage PE, Bromham DR, Ismail AA, Morris AF. The value of a single serum progesterone measurement in the midluteal phase as a criterion of a potentially fertile cycle ("ovulation") derived from treated and untreated conception cycles. *Fertil Steril* 1982;**37**:355–360.
- Hvidman HW, Bentzen JG, Thuesen LL, Lauritsen MP, Forman JL, Loft A, Pinborg A, Nyboe Andersen A. Infertile women below the age of 40 have similar anti-Müllerian hormone levels and antral follicle count compared with women of the same age with no history of infertility. *Hum Reprod* 2016;**31**:1034–1045.
- Ibrahim MI, Moustafa RA, Abdel-Azeem AA. Letrozole versus clomiphene citrate for superovulation in Egyptian women with unexplained infertility: a randomized controlled trial. *Arch Gynecol Obstet* 2012;**286**:1581–1587.
- Jafarabadi MN, Bagheri M, Ebrahimi Z, Shariat M, Haghollahi F. Endometrial scratching effect on pregnancy rate in intrauterine insemination cycles: a randomized controlled trial. *Int J Women's Health Reprod Sci* 2020;**8**:85–89.
- Jurkovic D, Geipel A, Gruboeck K, Jauniaux E, Natucci M, Campbell S. Three-dimensional ultrasound for the assessment of uterine anatomy and detection of congenital anomalies: a comparison with hysterosalpingography and two-dimensional sonography. *Ultrasound Obstet Gynecol* 1995;**5**:233–237.
- Kilic S, Tasdemir N, Yilmaz N, Yuksel B, Gul A, Batioglu S. The effect of anti-thyroid antibodies on endometrial volume, embryo grade and IVF outcome. *Gynecol Endocrinol* 2008;**24**:649–655.
- Ko JKY, Shi J, Li RHW, Yeung WSB, Ng EHY. 100 YEARS OF VITAMIN D: Effect of serum vitamin D level before ovarian stimulation on the cumulative live birth rate of women undergoing in vitro fertilization: a retrospective analysis. *Endocr Connect* 2022;**11**:e210444.
- Kowalczyk I, Wihstutz N, Buhrow G, Diedrich K. Subjective well-being in infertile couples. *J Psychosom Obstet Gynaecol* 2001;**22**:143–148.
- Kydonopoulou K, Delkos D, Rousso D, Ilonidis G, Mandala E. Association of plasminogen activator inhibitor-type 1 (PAI-1)-675 4G/5G polymorphism with unexplained female infertility. *Hippokratia* 2017;**21**:180–185.
- Lähteenmäki A. In-vitro fertilization in the presence of antisperm antibodies detected by the mixed antiglobulin reaction (MAR) and the tray agglutination test (TAT). *Hum Reprod* 1993;**8**:84–88.
- Lähteenmäki A, Reima I, Hovatta O. Treatment of severe male immunological infertility by intracytoplasmic sperm injection. *Hum Reprod* 1995;**10**:2824–2828.
- Lavy Y, Lev-Sagie A, Holtzer H, Revel A, Hurwitz A. Should laparoscopy be a mandatory component of the infertility evaluation in infertile women with normal hysterosalpingogram or suspected unilateral distal tubal pathology? *Eur J Obstet Gynecol Reprod Biol* 2004;**114**:64–68.
- Lazzarino G, Pallisco R, Bilotta G, Listorti I, Mangione R, Saab MW, Caruso G, Amorini AM, Brundo MV, Lazzarino G et al. Altered follicular fluid metabolic pattern correlates with female infertility and outcome measures of in vitro fertilization. *Int J Mol Sci* 2021;**22**:8735.
- Lopes VM, Lopes JR, Brasileiro JP, Oliveira I, Lacerda RP, Andrade MR, Tierno NI, Souza RC, Motta LA. Highly prevalence of vitamin D deficiency among Brazilian women of reproductive age. *Arch Endocrinol Metab* 2017;**61**:21–27.
- Lotti F, Frizza F, Balercia G, Barbonetti A, Behre HM, Calogero AE, Cremers JF, Francavilla F, Isidori AM, Kliesch S et al. The European Academy of Andrology (EAA) ultrasound study on healthy, fertile men: Scrotal ultrasound reference ranges and associations with clinical, seminal, and biochemical characteristics. *Andrology* 2021;**9**:559–576.
- Ludwin A, Pityński K, Ludwin I, Banas T, Knafel A. Two- and three-dimensional ultrasonography and sonohysterography versus hysteroscopy with laparoscopy in the differential diagnosis of septate, bicornuate, and arcuate uteri. *J Minim Invasive Gynecol* 2013;**20**:90–99.

- Makled AK, Farghali MM, Shenouda DS. Role of hysteroscopy and endometrial biopsy in women with unexplained infertility. *Arch Gynecol Obstet* 2014;**289**:187–192.
- Mardesic T, Ulcova-Gallova Z, Huttelova R, Muller P, Voboril J, Mikova M, Hulvert J. The influence of different types of antibodies on in vitro fertilization results. *Am J Reprod Immunol* 2000;**43**:1–5.
- Martinez AR, Bernardus RE, Kucharska D, Schoemaker J. Urinary luteinizing hormone testing and prediction of ovulation in spontaneous, clomiphene citrate and human menopausal gonadotropin-stimulated cycles. A clinical evaluation. *Acta Endocrinol (Copenh)* 1991;**124**:357–363.
- Menge AC, Medley NE, Mangione CM, Dietrich JW. The incidence and influence of antisperm antibodies in infertile human couples on sperm-cervical mucus interactions and subsequent fertility. *Fertil Steril* 1982;**38**:439–446.
- Milenkovic J, Milojkovic M, Mitic D, Stoimenov TJ, Smelcerovic Z, Stojanovic D, Vujic S, Bojanic N. Interaction of thrombophilic SNPs in patients with unexplained infertility-multifactor dimensionality reduction (MDR) model analysis. *J Assist Reprod Genet* 2020;**37**:1449–1458.
- Mol BW, Dijkman B, Wertheim P, Lijmer J, van der Veen F, Bossuyt PM. The accuracy of serum chlamydial antibodies in the diagnosis of tubal pathology: a meta-analysis. *Fertil Steril* 1997;**67**:1031–1037.
- Monem FM, Moalla HA. Antisperm antibodies and unexplained infertility in Syria. An Unsolved Problem? *Saudi Med J* 2003;**24**:912–913.
- Montanino Oliva M, Buonomo G, Carra MC, Lippa A, Lisi F. Myo-inositol impact on sperm motility in vagina and evaluation of its effects on foetal development. *Eur Rev Med Pharmacol Sci* 2020;**24**:2704–2709.
- Munro MG, Critchley HOD, Fraser IS; FIGO Menstrual Disorders Committee. The two FIGO systems for normal and abnormal uterine bleeding symptoms and classification of causes of abnormal uterine bleeding in the reproductive years: 2018 revisions. *Int J Gynaecol Obstet* 2018;**143**:393–408.
- Murto T, Bjuresten K, Landgren BM, Stavreus-Evers A. Predictive value of hormonal parameters for live birth in women with unexplained infertility and male infertility. *Reprod Biol Endocrinol* 2013;**11**:61.
- Nandi A, Raja G, White D, Tarek ET. Intrauterine insemination + controlled ovarian hyperstimulation versus in vitro fertilisation in unexplained infertility: a systematic review and meta-analysis. *Arch Gynecol Obstet* 2022;**305**:805–824.
- Nguyen DK, O'Leary S, Gadalla MA, Roberts B, Alvino H, Tremellen KP, Mol BW. The predictive value of anti-Müllerian hormone for natural conception leading to live birth in subfertile couples. *Reprod Biomed Online* 2022;**44**:557–564.
- O'Neill CL, Parrella A, Keating D, Cheung S, Rosenwaks Z, Palermo GD. A treatment algorithm for couples with unexplained infertility based on sperm chromatin assessment. *J Assist Reprod Genet* 2018;**35**:1911–1917.
- Oei SG, Helmerhorst FM, Keirse MJ. When is the post-coital test normal? A critical appraisal. *Hum Reprod* 1995;**10**:1711–1714.
- Orouji Jokar T, Fourman LT, Lee H, Mentzinger K, Fazeli PK. Higher TSH levels within the normal range are associated with unexplained infertility. *J Clin Endocrinol Metab* 2018;**103**:632–639.
- Pagidas K, Hemmings R, Falcone T, Miron P. The effect of antisperm autoantibodies in male or female partners undergoing in vitro fertilization-embryo transfer. *Fertil Steril* 1994;**62**:363–369.
- Pandian Z, Gibreel A, Bhattacharya S. In vitro fertilisation for unexplained subfertility. *Cochrane Database Syst Rev* 2015;**2015**:Cd003357.
- Papanikolaou EG, Vernaev V, Kolibianakis E, Assche EV, Bonduelle M, Liebaers I, Van Steirteghem A, Devroey P. Is chromosome analysis mandatory in the initial investigation of normovulatory women seeking infertility treatment? *Hum Reprod* 2005;**20**:2899–2903.
- Parsanezhad ME, Dadras N, Maharlouei N, Neghaban L, Keramati P, Amini M. Pregnancy rate after endometrial injury in couples with unexplained infertility: a randomized clinical trial. *Iran J Reprod Med* 2013;**11**:869–874.
- Pekel A, Gönenç A, Turhan N, Kafalı H. Changes of sFas and sFasL, oxidative stress markers in serum and follicular fluid of patients undergoing IVF. *J Assist Reprod Genet* 2015;**32**:233–241.
- Poppe K, Glinioer D, Van Steirteghem A, Tournaye H, Devroey P, Schiettecatte J, Velkeniers B. Thyroid dysfunction and autoimmunity in infertile women. *Thyroid* 2002;**12**:997–1001.
- Qu T, Yan M, Shen WJ, Li L, Zhu P, Li Z, Huang J, Han T, Hu W, Zhou R et al. Predictive serum markers for unexplained infertility in child-bearing aged women. *Am J Reprod Immunol* 2020;**83**:e13194.
- Rajah SV, Parslow JM, Howell RJ, Hendry WF. The effects on in-vitro fertilization of autoantibodies to spermatozoa in subfertile men. *Hum Reprod* 1993;**8**:1079–1082.
- Rehman R, Rajpar HI, Ashraf M, Iqbal NT, Lalani S, Alam F. Role of oxidative stress and altered thyroid hormones in unexplained infertility. *J Pak Med Assoc* 2020;**70**:1345–1349.
- Repalle D, Saritha KV, Bhandari S. Sperm DNA fragmentation negatively influences the cumulative live birth rate in the intracytoplasmic sperm injection cycles of couples with unexplained infertility. *Clin Exp Reprod Med* 2022;**49**:185–195.
- Rosen MP, Johnstone E, Addauan-Andersen C, Cedars MI. A lower antral follicle count is associated with infertility. *Fertil Steril* 2011;**95**:1950–1954.e1.
- Rudick B, Ingles S, Chung K, Stanczyk F, Paulson R, Bendikson K. Characterizing the influence of vitamin D levels on IVF outcomes. *Hum Reprod* 2012;**27**:3321–3327.
- Rull K, Grigороva M, Ehrenberg A, Vaas P, Sekavin A, Nömmemees D, Adler M, Hanson E, Juhanson P, Laan M. FSHB -211 G>T is a major genetic modulator of reproductive physiology and health in childbearing age women. *Hum Reprod* 2018;**33**:954–966.
- Sahmani M, Sakhinia E, Farzadi L, Najafipour R, Darabi M, Mehdizadeh A, Shahnazi V, Shaaker M, Noori M. Two common polymorphisms in the peroxisome proliferator-activated receptor γ gene may improve fertilization in IVF. *Reprod Biomed Online* 2011;**23**:355–360.
- Salas-Huetos A, Blanco J, Vidal F, Grossmann M, Pons MC, Garrido N, Anton E. Spermatozoa from normozoospermic fertile and infertile individuals convey a distinct miRNA cargo. *Andrology* 2016;**4**:1028–1036.
- Santoro N, Eisenberg E, Trussell JC, Craig LB, Gracia C, Huang H, Alvero R, Casson P, Christman G, Coutifaris C et al.; Reproductive Medicine Network Investigators. Fertility-related quality of life from two RCT cohorts with infertility: unexplained infertility and polycystic ovary syndrome. *Hum Reprod* 2016;**31**:2268–2279.
- Scott RT, Leonardi MR, Hofmann GE, Illions EH, Neal GS, Navot D. A prospective evaluation of clomiphene citrate challenge test screening of the general infertility population. *Obstet Gynecol* 1993;**82**:539–544.
- Senocak GC, Yapca OE, Borekci B. Comparison of pregnancy rates between patients with and without local endometrial scratching before intrauterine insemination. *J Gynecol Obstet Hum Reprod* 2017;**46**:687–690.
- Şentürk R, Tola EN, Bozkurt M, Doğuç DK. The role of oxidant status on the etiopathogenesis of unexplained infertility and intracytoplasmic sperm injection - embryo transfer success: a case-control study. *J Obstet Gynaecol* 2021;**42**:1312–1318.

- Seyam EM, Hassan MM, Mohamed Sayed Gad MT, Mahmoud HS, Ibrahim MG. Pregnancy outcome after office microhysteroscopy in women with unexplained infertility. *Int J Fertil Steril* 2015;**9**: 168–175.
- Showell MG, Mackenzie-Proctor R, Jordan V, Hart RJ. Antioxidants for female subfertility. *Cochrane Database Syst Rev* 2020;**8**: Cd007807.
- Steiner AZ, Pritchard D, Stanczyk FZ, Kesner JS, Meadows JW, Herring AH, Baird DD. Association between biomarkers of ovarian reserve and infertility among older women of reproductive age. *JAMA* 2017;**318**:1367–1376.
- Steinvil A, Raz R, Berliner S, Steinberg DM, Zeltser D, Levran D, Shimron O, Sella T, Chodick G, Shalev V et al. Association of common thrombophilias and antiphospholipid antibodies with success rate of in vitro fertilisation. *Thromb Haemost* 2012;**108**: 1192–1197.
- Subramanian MG, Kowalczyk CL, Leach RE, Lawson DM, Blacker CM, Ginsburg KA, Randolph JF Jr, Diamond MP, Moghissi KS. Midcycle increase of prolactin seen in normal women is absent in subjects with unexplained infertility. *Fertil Steril* 1997;**67**:644–647.
- Suganya J, Kujur SB, Selvaraj K, Suruli MS, Haripriya G, Samuel CR. Chromosomal abnormalities in infertile men from Southern India. *J Clin Diagn Res* 2015;**9**:Gc05–10.
- Tanahatoo S, Hompes PG, Lambalk CB. Accuracy of diagnostic laparoscopy in the infertility work-up before intrauterine insemination. *Fertil Steril* 2003;**79**:361–366.
- Tanahatoo SJ, Lambalk CB, Hompes PG. The role of laparoscopy in intrauterine insemination: a prospective randomized reallocation study. *Hum Reprod* 2005;**20**:3225–3230.
- Tersigni C, Castellani R, de Waure C, Fattorossi A, De Spirito M, Gasbarrini A, Scambia G, Di Simone N. Celiac disease and reproductive disorders: meta-analysis of epidemiologic associations and potential pathogenic mechanisms. *Hum Reprod Update* 2014;**20**:582–593.
- Trková M, Kapras J, Bobková K, Stanková J, Mejstnarová B. Increased micronuclei frequencies in couples with reproductive failure. *Reprod Toxicol* 2000;**14**:331–335.
- Unuane D, Velkeniers B, Anckaert E, Schiettecatte J, Tournaye H, Haentjens P, Poppe K. Thyroglobulin autoantibodies: is there any added value in the detection of thyroid autoimmunity in women consulting for fertility treatment? *Thyroid* 2013;**23**:1022–1028.
- Vani GT, Mukesh N, Rama Devi P, Usha Rani P, Reddy PP. Methylenetetrahydrofolate reductase C677T polymorphism is not associated with male infertility in a South Indian population. *Andrologia* 2012;**44**(Suppl 1):252–259.
- Vazquez-Levin MH, Notrica JA, Polak de Fried E. Male immunologic infertility: sperm performance on in vitro fertilization. *Fertil Steril* 1997;**68**:675–681.
- Veena BS, Upadhyaya S, Adiga SK, Pratap KN. Evaluation of oxidative stress, antioxidants and prolactin in infertile women. *Indian J Clin Biochem* 2008;**23**:186–190.
- Vermeulen N, Le Clef N, Veleza Z, D'Angelo A, Tilleman K. European recommendations for good practice in addition to an evidence-based guidelines programme: rationale and method of development. *BMJ Evid Based Med* 2019;**24**:30–34.
- Wang R, Watson A, Johnson N, Cheung K, Fitzgerald C, Mol BWJ, Mohiyiddeen L. Tubal flushing for subfertility. *Cochrane Database Syst Rev* 2020;**10**:Cd003718.
- Wang Y, Qian L. Three- or four-dimensional hysterosalpingo contrast sonography for diagnosing tubal patency in infertile females: a systematic review with meta-analysis. *Br J Radiol* 2016;**89**:20151013.
- Warchol-Biedermann K. The etiology of infertility affects fertility quality of life of males undergoing fertility workup and treatment. *Am J Mens Health* 2021;**15**:1557988320982167.
- Witkin SS, Bierhals K, Linhares I, Normand N, Dieterle S, Neuer A. Genetic polymorphism in an inflammasome component, cervical mycoplasma detection and female infertility in women undergoing in vitro fertilization. *J Reprod Immunol* 2010;**84**:171–175.
- Wong TY, Lensen S, Wilkinson J, Glanville EJ, Acharya S, Clarke F, Das S, Dawson J, Hammond B, Jayaprakasan K et al. Effect of endometrial scratching on unassisted conception for unexplained infertility: a randomized controlled trial. *Fertil Steril* 2022;**117**: 612–619.
- Yang JH, Chen MJ, Yang PK. Factors increasing the detection rate of intrauterine lesions on hysteroscopy in infertile women with sonographically normal uterine cavities. *J Formos Med Assoc* 2019;**118**:488–493.
- Yasin AL, Yasin AL, Basha WS. The epidemiology of anti-sperm antibodies among couples with unexplained infertility in North West Bank, Palestine. *J Clin Diagn Res* 2016;**10**:QC01–QC03.
- Yildiz G, Kurt D, Mat E, Yildiz P. The effect of local endometrial injury on the success of intrauterine insemination. *J Exp Clin Med (Turkey)* 2021;**38**:521–524.
- Yücel B, Kelekci S, Demirel E. Decline in ovarian reserve may be an undiagnosed reason for unexplained infertility: a cohort study. *Arch Med Sci* 2018;**14**:527–531.
- Zegers-Hochschild F, Adamson GD, Dyer S, Racowsky C, de Mouzon J, Sokol R, Rienzi L, Sunde A, Schmidt L, Cooke ID et al. The International Glossary on Infertility and Fertility Care, 2017. *Hum Reprod* 2017;**32**:1786–1801.