# Explaining Decision-Making between Exploration and Repetition: Key Factors for Physical Activity Recommendations

Ine Coppens imec - WAVES - Ghent University Ghent, Belgium Ine.Coppens@UGent.be Toon De Pessemier imec - WAVES - Ghent University Ghent, Belgium Toon.DePessemier@UGent.be Luc Martens imec - WAVES - Ghent University Ghent, Belgium Luc1.Martens@UGent.be

# Abstract

A challenge in promoting physical activity with recommender systems lies in balancing repeat recommendations to create habits, with exploration to prevent boredom. This study aims to identify the key variables influencing users' decision-making between the two. Through the analysis of data from an eight-week Micro-Randomized Trial conducted via a mobile health app, using random forest variable importance measures and SHAP analyses, we identified factors affecting these decisions. Our findings reveal that participants were more likely to explore new activities during the first two weeks of the intervention, in the afternoons and evenings, on Sundays, and when activities involved specific locations or workouts. These findings provide valuable insights into the transition from exploration to repetition, contributing to more effective recommender systems for physical activity promotion.

# **CCS** Concepts

Information systems → Personalization; Recommender systems; • Computing methodologies → Feature selection;
 Human-centered computing → Empirical studies in ubiquitous and mobile computing; User studies; • Applied computing → Psychology.

# Keywords

recommender systems; physical activity; decision-making; variable importance analysis; random forest; repetition; exploration

#### **ACM Reference Format:**

Ine Coppens, Toon De Pessemier, and Luc Martens. 2024. Explaining Decision-Making between Exploration and Repetition: Key Factors for Physical Activity Recommendations. In Proceedings of the 6th International Workshop on Health Recommender Systems co-located with 18th ACM Conference on Recommender Systems (HealthRecSys'24), Bari, Italy, October 18, 2024. 7 pages.

#### 1 Introduction

Insufficient physical activity (PA) is associated with adverse health outcomes [28]. To support motivation and long-term health benefits, individuals with low activity levels may benefit from engaging in enjoyable activities. Recommender Systems (RSs) can automatically personalize such enjoyable PAs by utilizing a wide range of information about users and their preferences [24, p 9].

As these physically inactive people might not have established healthy habits yet, the RS should repeat some activities to create habits, because repetition makes the behavior more automatic [15], increasing the chances for long-term engagement and positive health outcomes. However, the RS should also provide sufficient opportunity to explore new and varied activities to prevent boredom [2, 14, 32].

In addition to their recommended minimum of 150 minutes of moderate-to-vigorous PA (MVPA) per week, the World Health Organization (WHO) also recommends a mix of aerobic and musclestrengthening activities [28]. As such, both exploration for variety and repetition for habit formation are important for healthy behavior. The challenge lies in determining when an RS should prioritize one approach over the other.

This study focuses on identifying the key variables that impact users' decision-making process to choose either an exploration or repetition PA item, arguing that future RSs can integrate these variables in their own decision-making processes as well. The research question is:

*RQ*: Which variables affect users' decision to choose an exploration or repetition *PA* recommendation, and in which conditions are they more likely to choose one over the other?

By analyzing data from an eight-week Micro-Randomized Trial (MRT) conducted with a mobile health (mHealth) app, we aim to uncover the most important factors, determined with Random Forests' (RFs) variable importances [27] and SHapley Additive exPlanations (SHAP) [20] analyses. RFs are a machine learning technique used for classification or regression to predict an outcome variable based on several input variables by aggregating the results from multiple decision trees. RFs are widely used for data exploration and understanding using variable importance measures [27].

This paper merges two study waves: one that ran from October 2023 until January 2024 (started in autumn), and one from March until June 2024 (spring). Our previous research was solely conducted on the data from the study starting in autumn [7]. To increase participation and dataset, we replicated the study in spring and combined both study's data. In a previous paper [5], we already conducted a first analysis on the subjective perceptions and preferences measured with star rating of the repetition vs. exploration PA recommendations. The current paper dives deeper in the decisionmaking of the users using a set of factors that might influence their decision, contributing to the design of more effective RS algorithms in the PA domain.

The rest of the paper is organized as follows. Section 2 covers previous work on RFs in the domain of PA. The methods are discussed in Sect. 3. Next, the results and their discussion are elaborated in Sect. 4, followed by the conclusion in Sect. 5.

HealthRecSys'24, Bari, Italy, October 18, 2024

<sup>© 2024</sup> Copyright for the individual papers remains with the authors. Use permitted under Creative Commons License Attribution 4.0 International (CC BY 4.0). This volume is published and copyrighted by its editors

HealthRecSys'24, Bari, Italy, October 18, 2024

# 2 Related work

Previous work with RFs in the domain of PA mainly focused on the classification and recognition of the activity using sensors, such as accelerometers [13, 17, 21]. In those RFs, the input data consists of the data collected from the sensors and the predicted outcome variable is the detected type of PA. In our study, the input data is a combination of manual information (e.g., the self-reported motivation of the user at that time) and automatic information about the context (e.g., the weather).

Other studies used PA as an input variable in the RF to predict a certain outcome variable. In [9], body mass index and depressive symptoms are predicted with sedentary time and MVPA. Another study investigated severity of menopausal symptoms with an RF regression and found that PA level was in the top four of variable importances [29]. In [11], COVID-19 death rates were estimated with 29 socioeconomic and health-related factors with an RF.

PA amount was also used as outcome variable in several studies. Meeting the guidelines for sufficient aerobic PA in a target group of autistic adults was assessed in [12]. Another study found that greenery in the streets impacts the duration of light intensity PA the highest for older adults [31]. In [8], an RF algorithm performed best to predict the probability of achieving a daily steps threshold. In our study, the predicted outcome class is whether the user decides to engage in an exploration or repetition item, thus resulting in a binary classification.

Previous work has also investigated the exploration and repetition decision in next basket recommendations [18] and sequential recommendations [19]. While these are mainly focused on e-commerce and grocery shopping, our study is situated in the health domain and PA promotion with mHealth interventions.

#### 3 Methods

An Android app was created to display the personalized PA recommendations as shown in Fig. 1. Our PA dataset is based on activities from the compendium of physical activities [1], for which we distinguish between *workout* and *location* PA types. In our dataset, we also integrated *general* PA tips from the Belgian website for health (www.gezondleven.be), which contains small-effort activities people can integrate in all four PA situations of their daily life (*free time, during work, household task, active transport*). The resulting dataset contains 237 PA items, connected to the corresponding PA type and situation. For the content-based RS algorithm, all PA items were manually extended with 24 binary attributes to describe their content. For a full implementation description of content-based RS algorithm, we recommend reading our previous paper [6]. The exact same RS algorithm was used in this study, with an adjustment of the dataset and output list that now shows six recommendations.

A list of six items was chosen because six combinations can be made in our MRT study design: 2 from the outcome variable (repetition vs. exploration) x 3 from the PA type (general vs. location vs. workout). A repetition item refers to a personalized recommendation for an activity that the user has previously submitted during the study, while an exploration item is a personalized recommendation for an activity that the user has not yet submitted. By randomly positioning each of the six combination in the list for every participant at every delivery time, the micro-randomization of this study Ine Coppens, Toon De Pessemier, and Luc Martens



Figure 1: The main screen of the app displays six recommendations in a random order, corresponding to the six combinations of exploration versus repetition and the PA type.

is achieved [30] and position bias effects are prevented. As a result, the user can freely choose between a repetition or exploration item of all three PA types at any time.

We recruited healthy adults (between 18 and 65 years old), who have less than 150 min of MVPA per week, via the Sona Platform and Facebook groups for paid studies of Ghent University. For both study waves (starting in autumn and spring), participants were asked to use the app for at least eight consecutive weeks, after which they receive 30 EUR. To prevent false submissions about their PA behavior, participants were informed that they do not receive more money for submitting more PAs, but only qualify for the incentive if they actively use the app for eight weeks and complete all questionnaires. At the start and after eight weeks, they answer the pre-test and post-test questionnaire, respectively, both containing the European Health Interview Survey - Physical Activity Questionnaire (EHIS-PAQ) to measure weekly PA [10] and a question about their age group (18 - 44 or 45 - 65 years). The study received ethical approval from the Ethical Committee (www.ugent.be/pp/en/research/ec) on August 22, 2023 (reference number: 2023-061A).

An RF classifier is tuned with Randomized Search and trained using scikit-learn (scikit-learn.org/) version 1.5.1. Table 1 provides an overview of all 15 input variables of the RF model. The predicted output of the RF is whether the user engaged in a repetition or exploration item (binary classification). SHAP provides an explanation for this output by assigning feature importance values to each variable for a specific prediction [20]. For the variables with the highest feature importance, we apply the dependence plot function from SHAP package version 0.46.0 to show the relation between the input variable and the corresponding SHAP value for the RF's prediction [20]. For an additional statistical analyses on the EHIS-PAQ answers, SPSS Statistics v. 28 is used for the analyses with a Linear Mixed Model (LMM) using the MIXED procedure [26].

#### 4 Results and Discussion

Of the 62 participants who started the study (100% in the age group of 18 - 44 years), 34 continued for at least eight weeks. Throughout the study, the participants submitted 457 recommended items, of which the amounts are categorized in Table 2.

Firstly, the hyperparameters of the model were tuned with Randomized Search with 5-fold cross validation and a max depth of 6 to prevent overfitting. The resulting optimal hyperparameters are: n estimators = 493, min samples split = 10, min samples leaf = 10, max features = sqrt, max depth = 6, bootstrap = True, resulting in a train accuracy of .764 and test accuracy of .663. The corresponding feature importances are shown in Fig. 2, in which the time in study showed the highest relevance.

Figure 3 shows the SHAP dependence plots that visualize the relation between the six input variables with the highest feature importances and their corresponding SHAP value. We added a color legend to show possible interaction effects with days in study, as this variable ranked highest on the variable importances.

#### 4.1 The input variables

4.1.1 Day in study. The first dependence plot shows that days in the beginning of the study are associated with a higher prediction for an exploration item (above the y=0 line). The threshold at which the model switches from predicting exploration to repetition lies around day 15 in the study.

The switch from exploration to repetition after two weeks was also found in our previous study, which showed a significant interaction effect on the star rating feedback after this two-week mark for general PAs [5]. In this previous study [5], we referred to the lower effort and complexity of general PAs as the reason for the quicker shift to repetition, based on [15].

4.1.2 Hour of day. The second dependence plot reveals that exploration predictions are associated with afternoon and evening hours, but not with mornings or nights, with this effect becoming more pronounced later in the study. The highest predictions for explorations are between 4 and 10 pm, especially for later days in the study. An explanation for this pattern could be that our participants (who all belong in the age group of 18 - 44 years) prefer fixed routines in the morning before they go to work or school, and are more likely to explore new activities when they finished their daytime commitments.

This pattern could also be attributed to people's circadian rhythm and the distinction between morning chronotypes who rise and peak early in the day, and evening chronotypes who experience arousal in the afternoon or evening [22]. However, as we did not collect the chronotype of our participants, nor an exact age or occupation, we cannot derive a clear explanation from this. 4.1.3 Outdoor temperature. The third plot suggests a larger prediction for exploration when the outdoor temperature is around 10 degrees Celsius, with two threshold values around 5 and 15 degrees. According to [4], the first and last warm days of the year may motivate people for more PA, which could explain why they are more open to exploration in this temperature range, unlike higher temperatures above 20 degrees.

4.1.4 PA type and situation. High variable importance was also found for PA type and situation, of which the dependence plots are shown at the bottom of Fig. 3. The higher association with repetition in free-time, household, and general contexts can be due to established habits in these contexts. For example, exploration could be less preferred in household tasks because people already have their own routine for these.

Exploration was mostly associated with activities for transportation, during work, at a location, and as a workout. The results of our previous analysis [5] showed that exploration for general PAs was only preferred in the first two weeks of the study, after which higher star ratings were given to repetitions. This can explain why general PAs are mostly associated with repetition prediction in the SHAP dependence plot. Similarly, the star rating was consistently higher for exploration of location and workout PAs in the LMM [5], which corresponds to the conclusions of this SHAP analysis.

As 94% of the active transport and 75% of during work submits were an exploration, as displayed in Table 2, this explains the higher association with exploration. Nonetheless, submits in the situations of active transport and during work are limited, thwarting reliable analyses for these situations.

4.1.5 Day of week. The fifth plot shows that exploration predictions are more associated with Sundays, likely because more people have time off then, allowing more time for exploration. As Saturdays are not associated with more exploration, this can explain the lower variable importance of the "weekend/week day" variable.

4.1.6 Season. The season in which the participant started the study (autumn vs. spring) ranked seventh in the variable importances. We did not create a dependence plot for this variable, but conducted an LMM analysis with this variable to investigate the effect on the amount of weekly MVPA.

To investigate the increase in weekly PA, we compare the total MVPA measured by the EHIS-PAQ ( $Q4^*Q5 + Q7$ ) in the pre-test and post-test questionnaires [10]. An LMM was fitted with the time (pre-test vs. post-test), the season in which the participant started (autumn vs. spring), and their interaction as fixed effects [23, 26]. To account for possible variations between users, the user ID was considered as a random effect in a random coefficient model with a random slope [26].

We found a significant interaction effect, of which the mean fixed predicted values are shown in Fig. 4, illustrating that the group of participants who started the study in spring had a higher increase in weekly MVPA, on average (F(1,46.094)=4.079, p=.049). While a general increase in MVPA in both study groups was expected [16, 25], the stronger increase in the spring group can be explained because PA is higher in warmer months [4].

Table 1: We	put 15 input variables in the <b>R</b>	F, which can be categorize	d in eight categories and	d originate from various sources.

category	type	input variables for the RF	source
Time	continuous	day in study	device's clock
		hour of day	
	categorical	start season: spring / autumn	
		day of week	
		weekend / week day	
Weather	continuous	outdoor temperature	GPS + weather API
	categorical	clear sky	(OpenWeatherMap)
		clouds	
		rain	
Situation	categorical	free time / work / household / transport	dataset
PA type	categorical	general (e.g., walk during breaks) / location (e.g., minigolf) / workout (e.g., pilates)	dataset
Company	categorical	alone / with a buddy	self-reported
Motivation	continuous	score on 4	self-reported
Location	categorical	indoors / outdoors	self-reported
Step count	continuous	amount of steps already detected that day	accelerometer

#### input variable for the RF



Figure 2: The feature importances show that day in study, hour of day, outdoor temperature, PA situation, day of week, and PA type score the highest on the feature importances in the RF's exploration/repetition prediction.

# 4.2 Limitations and future work

As only 34 participants finished the full eight-week study, a large amount of data of the later weeks are missing. Additionally, the study was not conducted year-round or across different climate zones, limiting reliable conclusions about the impact of outdoor temperature. Given the limited submissions for transportation and work PAs, we suggest future research to include longer studies with a larger participant pool. Explaining Decision-Making between Exploration and Repetition



SHAP value for exploration submit prediction

Figure 3: The dependence plots from SHAP show a switch in predicting an exploration item (above the y=0 line) to a repetition item (under the y=0 line).

Table	2: The	amount	of s	submits	per	situation	and	per	PA
type,	includii	ng the pe	ercei	ntage of	the	situation.			

situation	PA type	amount	% of situation
transport	exploration	17	94%
	repetition	1	6%
during work	exploration	45	75%
	repetition	15	25%
household	exploration	73	55%
	repetition	59	45%
free time	exploration	153	62%
	repetition	94	38%





Figure 4: This interaction plot shows the significant interaction effect of the starting season on the increase in weekly MVPA from pre-test to post-test.

Our results show that the variables company, location, and motivation had lower importance. However, this company variable could be extended with the presence of a human trainer, as their supervision and guided workout plans, often containing repetition of activities, increase engagement in trainings delivered via mobile apps [3].

Nonetheless, our results indicate that time-related factors primarily drive the decision to explore new activities. We suggest integrating these time-related variables in future RSs for PA promotion. Although we identified specific conditions for exploration, we propose to tailor these conditions to the user. For example, an RS could learn at what moments a user prefers to explore a new activity. Additional information could be integrated in the system, such as the user's chronotype to take into account moments at which the user is most active [22].

#### 5 Conclusion

Two eight-week user studies with a total of 62 physically inactive participants (<150 minutes MVPA/week) were conducted to investigate factors influencing people's decision to either repeat or explore a PA recommendation. In the MRT, repetition and exploration PAs were provided to the user in an mHealth app at random positions, allowing participants to freely choose between the options at each delivery time.

The RF and SHAP approach identified key factors and conditions influencing the likelihood of exploring a new activity: in the first two weeks of the mHealth intervention, in afternoons and evenings, on Sundays, and for activities at a location or as a workout. Other factors scored lower on the variable importances of the decision between an exploration or repetition item, such as the season, company, motivation, and location. However, we did find a significant higher increase in weekly MVPA for the group that started the study in spring, suggesting that the season might not largely influence the exploration/repetition decision, but does affect the amount of PA.

By defining moments and contexts at which people are more open to explore new activities, this study contributes to future PA recommenders to balance between repeating favorite activities and introducing new ones.

#### Acknowledgments

This research received funding from the Flemish Government under the "Onderzoeksprogramma Artificiële Intelligentie (AI) Vlaanderen" programme.

#### References

- [1] Barbara E. Ainsworth, William L. Haskell, Stephen D. Herrmann, Nathanael Meckes, David R. Bassett, Catrine Tudor-Locke, Jennifer L. Greer, Jesse Vezina, Melicia C. Whitt-Glover, and Arthur S. Leon. 2011. 2011 Compendium of Physical Activities: A Second Update of Codes and MET Values. *Medicine & Science in Sports & Exercise* 43, 8 (Aug. 2011), 1575–1581. https://doi.org/10.1249/MSS. 0b013e31821ece12
- [2] Ashton Anderson, Ravi Kumar, Andrew Tomkins, and Sergei Vassilvitskii. 2014. The dynamics of repeat consumption. In *Proceedings of the 23rd international conference on World wide web*. ACM, Seoul Korea, 419–430. https://doi.org/10. 1145/2566486.2568018
- [3] Ludovico Boratto, Salvatore Carta, Fabrizio Mulas, and Paolo Pilloni. 2017. An e-coaching ecosystem: design and effectiveness analysis of the engagement of remote coaching on athletes. *Personal and Ubiquitous Computing* 21, 4 (Aug. 2017), 689–704. https://doi.org/10.1007/s00779-017-1026-0
- [4] Mathieu Bélanger, Katherine Gray-Donald, Jennifer O'loughlin, Gilles Paradis, and James Hanley. 2009. Influence of Weather Conditions and Season on Physical Activity in Adolescents. *Annals of Epidemiology* 19, 3 (2009), 180–186. https://doi.org/10.1016/j.annepidem.2008.12.008
  [5] Inc Coppens, Toon De Pessemier, and Luc Martens. 2024. Balancing Habit
- [5] Ine Coppens, Toon De Pessemier, and Luc Martens. 2024. Balancing Habit Repetition and New Activity Exploration: A Longitudinal Micro-Randomized

Trial in Physical Activity Recommendations. In Proceedings of the 18th ACM Conference on Recommender Systems (Bari, Italy). Association for Computing Machinery, New York, NY, USA, 5 pages. https://doi.org/10.1145/3640457.3691715

- [6] Ine Coppens, Toon De Pessemier, and Luc Martens. 2024. Exploring the added effect of three recommender system techniques in mobile health interventions for physical activity: a longitudinal randomized controlled trial. User Modeling and User-Adapted Interaction (July 2024). https://doi.org/10.1007/s11257-024-09407-z
- [7] Ine Coppens, Toon De Pessemier, and Luc Martens. 2024. Repeating my Workouts or Exploring new Activities? A Longitudinal Micro-Randomized User Study for Physical Activity Recommender Systems. In Adjunct Proceedings of the 32nd ACM Conference on User Modeling, Adaptation and Personalization (Cagliari, Italy) (UMAP Adjunct '24). Association for Computing Machinery, New York, NY, USA, 176–182. https://doi.org/10.1145/3631700.3664867
- [8] Talko B. Dijkhuis, Frank J. Blaauw, Miriam W. Van Ittersum, Hugo Velthuijsen, and Marco Aiello. 2018. Personalized Physical Activity Coaching: A Machine Learning Approach. Sensors 18, 2 (Feb. 2018), 623. https://doi.org/10.3390/s18020623
- [9] Glen E. Duncan, Ally R. Avery, Siny Tsang, Nathaniel F. Watson, Bethany D. Williams, and Eric Turkheimer. 2022. The pillars of health: influence of multiple lifestyle behaviors on body mass index and depressive symptoms in adult twins. *BMC Public Health* 22, 1 (Aug. 2022), 1487. https://doi.org/10.1186/s12889-022-13901-7
- [10] Jonas D. Finger, Jean Tafforeau, Lydia Gisle, Leila Oja, Thomas Ziese, Juergen Thelen, Gert B. M. Mensink, and Cornelia Lange. 2015. Development of the European Health Interview Survey - Physical Activity Questionnaire (EHIS-PAQ) to monitor physical activity in the European Union. Archives of Public Health 73, 1 (Dec. 2015), 59. https://doi.org/10.1186/s13690-015-0110-z
- [11] George Grekousis, Zhixin Feng, Joannis Marakakis, Yi Lu, and Ruoyu Wang. 2022. Ranking the importance of demographic, socioeconomic, and underlying health factors on US COVID-19 deaths: A geographical random forest approach. *Health & Place* 74 (March 2022), 102744. https://doi.org/10.1016/j.healthplace.2022.102744
- [12] Sean Healy, Benjamin Brewer, Elissa Hoopes, Alexis Paller, Shannon Mayberry, Joseph Maguire, Julie Daly, Paige Laxton, and Freda Patterson. 2022. Identifying the most proximal multi-level factors associated with meeting each of the 24-h movement behavior recommendations in a sample of autistic adults. *Disability and Health Journal* 15, 4 (Oct. 2022), 101367. https://doi.org/10.1016/j.dhjo.2022. 101367
- [13] Zhang Junjie, Cai Shenghao, Xu Jie, and Yuan Hua. 2022. Random forest-based physical activities recognition by usingwearable sensors. *Industria Textila* 73, 01 (March 2022), 27–33. https://doi.org/10.35530/IT.073.01.20215
- [14] Komal Kapoor, Karthik Subbian, Jaideep Srivastava, and Paul Schrater. 2015. Just in Time Recommendations: Modeling the Dynamics of Boredom in Activity Streams. In Proceedings of the Eighth ACM International Conference on Web Search and Data Mining. ACM, Shanghai China, 233–242. https://doi.org/10.1145/ 2684822.2685306
- [15] Phillippa Lally, Cornelia H. M. Van Jaarsveld, Henry W. W. Potts, and Jane Wardle. 2010. How are habits formed: Modelling habit formation in the real world. *European Journal of Social Psychology* 40, 6 (Oct. 2010), 998–1009. https: //doi.org/10.1002/ejsp.674
- [16] Liliana Laranjo, Ding Ding, Bruno Heleno, Baki Kocaballi, Juan C Quiroz, Huong Ly Tong, Bahia Chahwan, Ana Luisa Neves, Elia Gabarron, Kim Phuong Dao, David Rodrigues, Gisela Costa Neves, Maria L Antunes, Enrico Coiera, and David W Bates. 2021. Do smartphone applications and activity trackers increase physical activity in adults? Systematic review, meta-analysis and metaregression. *British Journal of Sports Medicine* 55, 8 (2021), 422–432. https://doi.org/10.1136/ bjsports-2020-102892 arXiv:https://bjsm.bmj.com/content/55/8/422.full.pdf
- [17] Kangjae Lee and Mei-Po Kwan. 2018. Physical activity classification in free-living conditions using smartphone accelerometer data and exploration of predicted results. *Computers, Environment and Urban Systems* 67 (Jan. 2018), 124–131. https://doi.org/10.1016/j.compenvurbsys.2017.09.012
- [18] Ming Li, Sami Jullien, Mozhdeh Ariannezhad, and Maarten de Rijke. 2023. A Next Basket Recommendation Reality Check. ACM Trans. Inf. Syst. 41, 4, Article 116 (apr 2023), 29 pages. https://doi.org/10.1145/3587153
- [19] Ming Li, Ali Vardasbi, Andrew Yates, and Maarten De Rijke. 2023. Repetition and Exploration in Sequential Recommendation. In Proceedings of the 46th International ACM SIGIR Conference on Research and Development in Information Retrieval. ACM, Taipei Taiwan, 2532–2541. https://doi.org/10.1145/3539618.3591914
- [20] Scott M. Lundberg and Su-In Lee. 2017. A unified approach to interpreting model predictions. In Proceedings of the 31st International Conference on Neural Information Processing Systems (Long Beach, California, USA) (NIPS'17). Curran Associates Inc., Red Hook, NY, USA, 4768–4777. https://doi.org/10.5555/3295222. 3295230
- [21] Mamoun T. Mardini, Chen Bai, Amal A. Wanigatunga, Santiago Saldana, Ramon Casanova, and Todd M. Manini. 2021. Age Differences in Estimating Physical Activity by Wrist Accelerometry Using Machine Learning. *Sensors* 21, 10 (May 2021), 3352. https://doi.org/10.3390/s21103352
- [22] Cynthia P. May, Lynn Hasher, and Karl Healey. 2023. For Whom (and When) the Time Bell Tolls: Chronotypes and the Synchrony Effect. Perspectives on

Explaining Decision-Making between Exploration and Repetition

HealthRecSys'24, Bari, Italy, October 18, 2024

Psychological Science 18, 6 (Nov. 2023), 1520–1536. https://doi.org/10.1177/ 17456916231178553

- [23] Geert Molenberghs and Geert Verbeke. 2009. A Model for Longitudinal Data. In Linear Mixed Models for Longitudinal Data. Springer New York, New York, NY, 19–29. https://doi.org/10.1007/978-1-4419-0300-6\_3 Series Title: Springer Series in Statistics.
- [24] Francesco Ricci, Lior Rokach, and Bracha Shapira (Eds.). 2022. Recommender Systems Handbook. Springer US, New York, NY. https://doi.org/10.1007/978-1-0716-2197-4
- [25] Amelia Romeo, Sarah Edney, Ronald Plotnikoff, Rachel Curtis, Jillian Ryan, Ilea Sanders, Alyson Crozier, and Carol Maher. 2019. Can Smartphone Apps Increase Physical Activity? Systematic Review and Meta-Analysis. *Journal of Medical Internet Research* 21, 3 (March 2019), e12053. https://doi.org/10.2196/12053
- [26] SPSS Inc. 2005. Linear Mixed-Effects Modeling in SPSS: An Introduction to the MIXED Procedure. Technical Report. SPSS Inc. https://imaging.mrc-cbu.cam.ac. uk/statswiki/FAQ/multilevel?action=AttachFile&do=get&target=spssegs.pdf
- [27] A. Verikas, A. Gelzinis, and M. Bacauskiene. 2011. Mining data with random forests: A survey and results of new tests. *Pattern Recognition* 44, 2 (Feb. 2011), 330-349. https://doi.org/10.1016/j.patcog.2010.08.011

- [28] World Health Organization. 2020. WHO guidelines on physical activity and sedentary behaviour. http://www.ncbi.nlm.nih.gov/books/NBK566045/ OCLC: 1237095892.
- [29] SongWen Wu, Yi Shi, Qiao Zhao, and Ke Men. 2023. The relationship between physical activity and the severity of menopausal symptoms: a cross-sectional study. *BMC Women's Health* 23, 1 (April 2023), 212. https://doi.org/10.1186/ s12905-023-02347-7
- [30] Jing Xu, Xiaoxi Yan, Caroline Figueroa, Joseph Jay Williams, and Bibhas Chakraborty. 2023. A flexible micro-randomized trial design and sample size considerations. *Statistical Methods in Medical Research* 32, 9 (Sept. 2023), 1766–1783. https://doi.org/10.1177/09622802231188513
- [31] Peng Zang, Hualong Qiu, Fei Xian, Linchuan Yang, Yanan Qiu, and Hongxu Guo. 2022. Nonlinear Effects of the Built Environment on Light Physical Activity among Older Adults: The Case of Lanzhou, China. International Journal of Environmental Research and Public Health 19, 14 (July 2022), 8848. https://doi. org/10.3390/ijerph1914848
- [32] Yuan Zhang. 2022. Variety-Seeking Behavior in Consumption: A Literature Review and Future Research Directions. *Frontiers in Psychology* 13 (June 2022), 874444. https://doi.org/10.3389/fpsyg.2022.874444



# The 6th International Workshop on Health Recommender Systems

Organizers: Hanna Hauptman, Christoph Trattner, Helma Torkamaan



October 18, 2024

Bari, Italy

 $\bullet \bullet \bullet \bullet \bullet \bullet$