


ESSAY

Conservation-focused mapping of avian migratory routes using a pan-European automated telemetry network

Lucy Mitchell¹  | Vera Brust²  | Thiemo Karwinkel³  | Susanne Åkesson⁴  |
 Dmitry Kishkinev⁵  | Gabriel Norevik⁴  | Tibor Szep^{6,7}  | Anders Hedenström⁴  |
 Sander Lagerveld⁸  | Barbara Helm⁹  | Heiko Schmaljohann³ 

¹Terrestrial Ecology Group, Department of Biology, Faculty of Sciences, Ghent University, Gent, Belgium

²Naturschutzbund Deutschland (NABU) e.V., Berlin, Germany

³Institute of Biology and Environmental Sciences, School of Mathematics and Science, Carl von Ossietzky Universität Oldenburg, Oldenburg, Germany

⁴Department of Biology, Lund University, Lund, Sweden

⁵School of Life Sciences, Keele University, Newcastle-under-Lyme, UK

⁶Institute of Environmental Sciences, University of Nyíregyháza, Nyíregyháza, Hungary

⁷MME/BirdLife Hungary, Nyíregyháza, Hungary

⁸Wageningen Marine Research, Wageningen University and Research, Den Helder, The Netherlands

⁹Bird Migration Unit, Swiss Ornithological Institute, Sempach, Switzerland

Correspondence

Lucy Mitchell, Terrestrial Ecology Group, Department of Biology, Faculty of Sciences, Ghent University, K. L. Ledeganckstraat 35, 9000 Gent, Belgium. Email: lucy.mitchell@ugent.be

Article impact statement: Demographic knowledge gaps in the conservation of migratory species can be filled using Motus automated VHF telemetry.

Funding information

Deutsche Forschungsgemeinschaft (DFG), Grant/Award Numbers: 395940726, SCHM 2647/3-1, SCHM 2647/4-1, SCHM 2647/7-1; Federal Agency for Nature Conservation (BfN), Grant/Award Numbers: 351582210A, 351986140A, 352315100B; Yorkshire Water Biodiversity Fund, Grant/Award Number: YW 270521; Scottish Ornithological Club; British Ecological Society, Grant/Award Number: SR20/1532

Abstract

Accelerated biodiversity loss has destabilized functional links within and between ecosystems. Species that cross different ecosystems during migration between breeding and nonbreeding sites are particularly sensitive to global change because they are exposed to various, often ecosystem-specific, threats. Because these threats have lethal and nonlethal effects on populations, many migratory species are declining, making this group especially vulnerable to global change. To mitigate their decline, research at a continental and flyway scale is required to adequately monitor changes in the migratory and demographic processes of populations during all parts of the annual cycle. The Motus Wildlife Tracking System (Motus) could provide a solution to data gaps that exist for small, migratory species. Motus is an automated telemetry system for animal tracking that uses a single very-high-frequency radio signal to track tagged individuals. Motus can provide information on movements made by individuals of small migrant species, thereby aiding the understanding of aspects of their migration that could affect demographic parameters. Conservation-focused research opportunities related to Motus include identification of critical stopover sites that support and connect multiple species and insight into migratory decisions in small migrant birds related to environmental stressors, such as artificial light at night. Examples of stopover studies from the existing network that demonstrate its utility include identification of a high-conservation-value stopover area for the blackpoll warbler (*Setophaga striata*) in the eastern United States. Geographical gaps in the network across the Mediterranean region and across eastern Europe need to be filled to track continent-wide movements. Motus can provide individual-level migration information for a variety of small-bodied taxa, and a drive to expand the network will improve its ability to direct conservation plans for such species.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Conservation Biology* published by Wiley Periodicals LLC on behalf of Society for Conservation Biology.

KEYWORDS

automated radio tracking, demographic parameters, flyway conservation, management plans, migration

INTRODUCTION

Biodiversity loss driven by land-use change and exploitation of natural resources and affected further by climatic disruption is a defining feature of the Anthropocene (Sala et al., 2000). A decline in habitat availability and disruption to ecosystem structure, reducing critical services, such as nutrient cycling, carbon storage, and flood control, have led to declines in a wide range of taxa globally (Jaureguiberry et al., 2022). The impacts of anthropogenic development not only manifest through physical changes (i.e., habitat loss) but also through increases in zoonotic and vector-borne diseases (Jaureguiberry et al., 2022) and pest outbreaks (Ayres & Lombardero, 2018). These impacts affect species' distributions, abundances, and fitness and consequently their ability to complete their life cycle successfully (Bellard et al., 2012).

Of particular concern are migratory species, which serve as ecological indicators and providers of vital contributions to ecosystem functioning, including biomass production, pollination, and pest control (Bauer & Hoye, 2014; Satterfield et al., 2020). Migratory species experience a variety of environmental conditions on their seasonal, sometimes intercontinental, journeys (Horton et al., 2020; Howard et al., 2020; Turbek et al., 2018; Zurell et al., 2018). Rapid changes in land use and configuration occurring within their annual cycle may mean that their requirements for reproduction and survival are compromised (Birnie-Gauvin et al., 2020; Marcacci et al., 2022; Rigal et al., 2023). There are also additional threats, such as hunting (Jiguet et al., 2019), modification of physical barriers due to the addition of anthropogenic structures (Gauld et al., 2022), and increasingly unpredictable climatic patterns decoupling the phenology of ecologically linked species (Clarke et al., 2022; Iler et al., 2021).

These challenges directly conflict with the multifactorial optimization of migration, which is often based on inherited, integrated migration strategies (Åkesson & Helm, 2020; Fattorini et al., 2023; Schmaljohann et al., 2022). Although migrant species differ in their migratory timing, distance, speed, and route, their journeys all involve repeated, alternating migratory endurance flights and stopover periods for resting, recovering, and refueling (Åkesson & Hedenström, 2007; Alerstam et al., 2003; Schmaljohann et al., 2022). Understanding the factors affecting population trends of these species (i.e., the changes in vital rates that drive population growth or decline) is essential (Morrison et al., 2016) because many migratory species cannot respond to changes at a sufficiently rapid pace, which can lead to widespread population declines (Both et al., 2006; Frick et al., 2020; Vickery et al., 2023; Wilcover & Wikelski, 2008).

The Convention on the Conservation of Migratory Species highlights the need for a multispecies, flyway-level perspective in terms of research into population declines (Chowdury et al., 2023; Frick et al., 2020; Marcacci et al., 2022; UNEP, 2020; Vickery et al., 2023). However, gathering data from a sufficiently high number of individuals from different populations at this

scale is extremely challenging (McKinnon & Love, 2018; Morrison et al., 2016) and relies on international collaboration (Nadal et al., 2020; Serratos et al., 2024; Vickery et al., 2023). Particularly for small and light migratory passerines, waders, and highly aerial species such as swifts, their size and behavior make studying their movements difficult (Fiedler, 2009; Wikelski et al., 2007).

CURRENT METHODS AND THEIR LIMITATIONS

Studying when and where differences in population processes occur in migratory birds is notoriously difficult (Border et al., 2017; Doerr & Doerr, 2005). However, quantifying variation in survival, mortality, emigration, and immigration (summarized as dispersal) is crucially important to formulating effective conservation measures for populations and species that are at risk of decline (DeMars et al., 2023; Gómez et al., 2021).

There is little detailed spatial and temporal information on small birds that migrate. Broudscale migration patterns across Europe, including concentrations of avian and insect migrants passing through marine and mountainous regions, have been identified using radar (Bruderer & Jenni, 1990; Bruderer & Liechti, 1999; Hirschhofer et al., 2024; Nilsson et al., 2019; Weisshaupt et al., 2021). Yet, radar data largely do not allow researchers to tease out species-specific and individual-level variation in large-scale movements (Schmaljohann et al., 2008; Zugg et al., 2008). Such information would facilitate linking individual migratory behaviors to demography, physiology, and ecology.

Several million individuals have been marked with metal or color rings across Europe (Du Feu et al., 2016; Spina et al., 2022), contributing to fundamental knowledge of bird movements. Yet, recapture, recovery, and resighting probabilities are relatively low (across 32 European-level ringing schemes, recovery rate for all species combined ranges from 0.6% to 7.6% [Baillie, 1995]). This is particularly the case on wintering grounds but is highly variable among species and locations (Thorup et al., 2014). For example, the willow warbler (*Phylloscopus trochilus*) is ringed in huge numbers on its breeding ground in northern Europe, but only a very small number of recoveries occur in Africa. For each recovery of this species in sub-Saharan Africa, 16,000 individuals need to be marked on the breeding grounds in Finland (Hedenström & Pettersson, 1987).

The disadvantages of these methods can be mitigated by tracking individual migratory prebreeding and postbreeding dispersal (Mukhin et al., 2005; Züst et al., 2023) and nomadic, nonbreeding movements (McKinnon et al., 2019; Snell et al., 2018). However, individual tracking of small migrants requires tracking devices weighing a maximum of 3%–5% of an individual's body weight (Barron et al., 2010), which excludes most tracking technology on the market (Figure 1) (Bridge et al., 2011; McKinnon & Love, 2018), including new low-power, wide-area




FIGURE 1 Number of publications per year resulting from Motus data (Source: motus.org).

devices, such as SigFox and LoRaWAN (Wild et al., 2023). Radio transmitters, however, have already reached minimum weights of currently 0.13 g (Lotek NanoPin), lighter than the smallest light-level geolocators (Lotek: 0.3 g) and significantly smaller than GPS units that require significant energy to relay information to a satellite and fix a position. Some radio-tracking systems (e.g., tRackIT System and ATLAS project) have narrow spatial coverage and a limit to how many individuals (ca. 200) can be monitored concurrently (Beardsworth et al., 2022; Gottwald et al., 2019).

The Motus Wildlife Tracking System (Motus) exploits a network of very-high-frequency (VHF) receiving stations equipped with directional Yagi antennae on the same frequency. These receivers continuously receive and record uniquely coded signals from tagged individuals, and there is no need for recapture (Imlay et al., 2020; Taylor et al., 2017). We focused on how Motus can fill conservation and demographic-specific knowledge gaps through tracking of migratory birds. We sought to spark further collaborative use of Motus to create a denser network in Europe that resembles the situation in North America.

MOTUS AND ITS BENEFITS

Motus was devised through a partnership between Acadia University and Birds Canada researchers (Taylor et al., 2011, 2017),

and its spread across the Americas is a great success story of collaborative research (<https://motus.org>). Globally, to date (December 2024) there are 899 tagging projects, which combined have tagged 51,089 animals of 403 species. The entire Motus network consists of 2066 receivers, and the largest single project array consists of 109 receivers in Ontario, Canada.

Publications resulting from Motus data total 218, which combined were cited, according to Zotero, >500 times. The number of publications based on Motus data has more than doubled since 2015 (Figure 2), and the lead and coauthors of these publications are rarely limited to academics. The application of Motus has been recognized by multiple stakeholders, such as the US Fish and Wildlife Service, National Parks Service, Canadian Wildlife Service, and BirdLife Europe (Machado et al., 2024).

Investment in the network across North America continues to grow. In March 2024, a grant of CSD3.1 million was awarded to a consortium of 5 Canadian universities and Birds Canada to continue installing Motus receiving stations across the country and to further community-based science. This investment, combined with a specific mention in the Convention on Migratory Species of automated radio tracking deployed at a flyway scale (COP13, Resolution 12.26), demonstrates the current and potential future value of Motus to conservation.

The initiation of Motus in Europe started in 2017. Although the network has grown more slowly than in the Americas, there is now a dense network of passive receiving stations (Figure 3) along the coasts of Germany, the Netherlands, and the United




FIGURE 2 Capability and context of tags enabled for Motus (gray dotted lines, variation on both axes taking into account programming influence on battery life and differences among and between device types; orange Wi-Fi symbols, transmission capability independent of bird's return to a specific location). Tag types are positioned approximately in relation to their mean battery lifetime and size.

Kingdom and to a slightly lesser extent in Sweden, Denmark, Belgium, and France. There are a number of stations in other countries as well as offshore (Figure 4).

The network of stations in Europe is patchy, particularly in eastern Europe, and there is a lack of a universally permitted tracking frequency, so it does not yet allow continuous tracking across the continent. In many European countries, the frequency of 150.1 MHz is authorized temporarily or permanently for wild animal telemetry tracking. Multifrequency detection by Motus receivers is possible, but it incurs additional expense for extra equipment. For example, adding antennas to receiving stations for monitoring the license-free frequency of 434 MHz would cost approximately €80–300 per station, for an additional 1–4 antennas, plus the cost of extra cables. For the tags, researchers can select from among a number of options and device parameters, including burst interval (usually from 1 s to 1 min), battery or solar power, attachment type, and antenna type, to meet their specific scientific requirements (Figure 1).

Motus has many promising features, including its extended temporal and spatial data gathering capacity, relative to standard radio tracking. In addition to autonomous, near-real-time recording of the receivers and sub-0.5-g tags, the spatial scale of detections is in the order of several kilometers, rather than orders of magnitude higher, as with geolocators (Taylor et al., 2017), although new multisensor tags have improved substantially in positional accuracy (Nussbaumer et al., 2023). Receiving stations can, in theory, be placed anywhere (Figure 3b) and have a detection range of >10 km; therefore, data capture is

less limited by researcher effort, in contrast to commonly used methods, such as bird ringing (Flack et al., 2022; Griffin et al., 2020). Fixed positioning of the receiving stations (ideally at sites of importance to the species of interest to maximize detection probability) and an unrestricted recording period allow standardized data collection and thus reduce observer bias (Griffin et al., 2020).

Despite the advantages, there are drawbacks of radio-tracking studies in general. Most studies do not detect all tagged individuals. The reasons for this are many and not mutually exclusive but include loss of the tag, predation, emigration, tag failure, topography, and weather conditions. Crewe et al. (2020) reported detection rates of 50%–70%, whereas a dense coverage of receivers on the small island of Helgoland, Germany, resulted repeatedly in detection rates of 95%–100% (Karwinkel et al., 2022, 2024). There are also uncommon occurrences of high levels of false positive, or uncoded detections, which can appear if large numbers of individuals are released at once close to a receiver. Mitigations, such as staggered switching on of the tags to encourage differential pulse emission, can be put in place, and the numbers required to cause this confusion mean this is unlikely to happen in a natural scenario.

Motus is already producing important insights into the movements of migrating and wintering European birds, including a better understanding of the migratory and premigratory movements of sea-crossing thrushes (Brust et al., 2019) and differences between long- and short-distance migrants in stopover time and flight direction (Packmor et al., 2020; Ruppel et al.,




FIGURE 3 (a) European robin (*Erithacus rubecula*) with attached radio transmitter and leg-loop harness (shown above the bird) and (b) a Motus receiving station (6-m tall) with 4 six-element Yagi antennas pointing in 4 directions. The station is solar powered and has a buffer battery (in aluminum box on ground). Electronics are in the yellow box on the pole. Detailed information about tagging animals and building stations is on the Motus Webpage (motus.org/resources/) and available from regional Motus coordinators (motus.org/groups/regional-coordination-groups/). Photos by T.K.

2023). Examples from Europe and North America show that Motus can gather long-term, annual cycle data, for a relatively low cost, on groups and in periods (e.g., juvenile fledging) that are often missing from population studies (Martell et al., 2023; Satterfield et al., 2020).

HOW MOTUS CAN HELP TO ADDRESS KNOWLEDGE GAPS IN SMALL MIGRATORY BIRD DEMOGRAPHICS

Survival and mortality

Despite the biological significance of survival and mortality to population size and its dynamics (Sandercock et al., 2020), little is known about either rate in migratory passerines. In migratory species, variation in survival among populations can be linked to alternative routes and their different pressures (Hewson et al., 2016). The latter may increase population-specific immediate and delayed fitness costs (Dhanjal-Adams et al., 2017), which might be particularly prevalent in areas that support high numbers of comigrants (multiple species moving through major

sites and corridors simultaneously [Cohen & Satterfield, 2020]). The convergence of otherwise spatially segregated populations at single locations may also increase the probability of disease transmission, which can have delayed fitness costs (Cohen & Satterfield, 2020).

To obtain information on route- or area-specific mortality rates, focusing receiving station placement in closely packed fence or curtain formation (Figure 4) would provide check-points for tagged migrants along their migratory routes. If there are sufficient stations intersecting migratory routes (and adequate numbers of individuals are tagged), then obstacles that slow down migration can be identified and estimates of mortality rates for such areas can be made (Buechley et al., 2021; Klaassen et al., 2014). Survival has been estimated successfully based on Motus data for Kirtland's warbler (*Setophaga kirtlandii*) (Cooper et al., 2024). This species, with its limited population size and discrete wintering range, lends itself to Motus tagging, and a robust design Cormack–Jolly–Seber model allowed the calculation of apparent survival rates with a high level of certainty, knowing that a high proportion of marked individuals had been detected.

Gonzalez et al. (2021) used Motus data to identify habitat-specific overwinter survival rates in Swainson's thrush (*Catharus ustulatus*), which can be used to inform habitat protection and management on the wintering grounds. Motus data have also been used by Brunner et al. (2022) to identify high migratory connectivity among populations of the elusive Swainson's warbler (*Limnothlypis swainsonii*), which has implications for population-specific changes and can direct future monitoring work. These cross-continental studies demonstrate the power of Motus to collect data at multiple scales along the length of a flyway. Extensive testing of detection capability of an antenna array in a fixed area is essential to maximizing coverage and the ability to produce survival estimates. It is better yet if survival is estimated across a limited area (Cooper et al., 2024) and a restricted temporal period to increase the robustness of the estimates (Evans et al., 2020).

Identification of stopover sites

Motus can be used in regional arrays that expand outward from a known stopover site, allowing identification of exploratory and regional movements by birds that may be assessing the wider area, often at night (Brown & Taylor, 2015; Schmaljohann & Eikenaar, 2017; Taylor et al., 2011). In Europe, this could build on current ringing efforts at hotspots of bird occurrence (e.g., Bay of Biscay, Strait of Gibraltar) at spatial scales not feasible with ringing alone. Pinpointing specific sites for targeted conservation efforts is important; limited, localized stopover site use can increase vulnerability in certain migrating species (Bayly et al., 2013; Gómez et al., 2014; Hagelin et al., 2021).

Information on arrival and departure times from Motus data on multiple individuals of different species can help elucidate the functions of stopover sites (Linscott & Senner, 2021; Moore, 2018; Schmaljohann et al., 2022). Identification of these functions could be very valuable in the context of future global




FIGURE 4 Current Motus receiving station network (purple dots) across Europe and hypothetical future stations (yellow dots) to show the potential Motus data have to answer demographic and conservation-focused questions about bird migration (blue arrows, flyways and movements of interest).

climate change, when the current conditions of stopover sites may degrade or even disappear (Bayly et al., 2018). Smetzer and King (2018) used data from a regional Motus network to identify a major stopover area for blackpoll warblers (*Setophaga striata*) and red-eyed vireos (*Vireo olivaceus*) in the Gulf of Maine of the United States. The prolonged stopovers recorded for both species suggest that the region may be a major refueling area for birds preparing for very long-distance migratory flights, thus demonstrating the area's high conservation value.

Stopover sites on either side of ecological barriers could be equipped with Motus stations at high densities (e.g., 5–10 km between stations, but variation in detection distance due to topography and the behavior of the species must be taken into account) to provide insights into how birds respond to such barriers (e.g., Sjöberg et al., 2015; Zenzal et al., 2021). This might include local- to regional-scale movements before crossing a barrier, intrinsic and extrinsic conditions required for a successful crossing, stopover duration, departure directions, and potential differences between populations and seasons. Holberton et al. (2019) and Herbert et al. (2022) used Motus data to demonstrate site-based variation in stopover duration, which

was related, at least in part, to bird condition and morphology. This indicates some level of migratory connectivity and thus that loss or degradation of one or more stopover sites could have population-level effects.

Dispersal, immigration, and emigration

Natal and breeding dispersal are critical but understudied fundamental biological processes, partly because the survival rates of nestlings and juveniles are generally so low that high personnel and financial investments are required to track a few individuals. Dispersal consists of the initial process of emigration from a breeding site and the subsequent immigration to another (Matthysen & Clobert, 2012).

Species with discrete breeding sites restricted by habitat may become increasingly inbred, and this inbreeding could be exacerbated by habitat loss, climate change, and a lack of immigration (Day et al., 2024). Such changes may consequently lead to a species' rapid decline if survival is also low (Schaub et al., 2012, 2013). Understanding how these populations are

connected through immigration and emigration (e.g., as in le Roux & Nocera [2021] who used Motus data on chimney swifts [*Chaetura pelagica*]) is important for deciding what conservation measures might be useful to avoid loss of genetic diversity (Driscoll et al., 2014). Researchers can estimate emigration and immigration rates of a species of interest through comprehensive tagging campaigns (ethical considerations of such projects notwithstanding [Soulsbury et al., 2020]), where Motus stations can cover initial breeding sites and potential areas where birds might disperse.

Regional-scale movements of juvenile blackpoll warblers, Kirtland's warbler, and barn swallow (*Hirundo rustica*) have been demonstrated prior to migration based on data from the Motus network (Brown & Taylor, 2015, 2017; Cooper & Marra, 2020; Evans, 2018). Data are particularly needed from juveniles to assess when, how, and why they decide on breeding site settlement. Studies by Doerr and Doerr (2005) and Mukhin et al. (2018) suggest that tracking the dispersal of breeders and fledging juveniles to new habitats in the region is feasible using this system.

Questions remain about the function of exploratory dispersal movements, which may be preparatory information-gathering trips ("homing target" or "habitat optimization" hypotheses [Mitchell et al., 2015]), and premigratory flights (Züst et al., 2023). These flights may relate to range expansion and individual or species responses to climate change (Driscoll et al., 2014; Dufour et al., 2021, 2022). Tracking individuals during the dispersal phase could improve understanding of the role of (long-distance) dispersal in the evolution of new migration routes and wintering grounds, perhaps as part of the wider phenomenon of vagrancy (Dufour et al., 2021, 2022; Lees & Gilroy, 2009).

Motus' ability to expand spatially and temporally beyond the capabilities of manual VHF tracking, thus reducing bias and monitoring hidden movements (Züst et al., 2023), can then increase the power of studies on juvenile fledging movement (Cox & Kesler, 2012) and medium- to long-distance postbreeding dispersal (Evans et al., 2018; Hayes et al., 2024). Results from such studies can inform conservation decisions and improve understanding of how far and in what direction juveniles disperse. Tracking of many different young individuals can also highlight how individual phenotypes and differences in body condition might lead to differential postfledging survival (Motus data used in a study of fledging barn swallows [Evans et al., 2020]) and how this might be affected by surrounding habitat quality (wood thrush [*Hylocichla mustelina*] [Hayes et al., 2024]). These practical elements are invaluable to formulating effective conservation measures and facilitating population stability (Endriss et al., 2019; Niebuhr et al., 2015; Travis & Dytham, 2013).

Understanding migratory decisions

In addition to using Motus data to describe migration, it can facilitate an experimental approach (i.e., extending laboratory-based studies in natural scenarios) (Goymann et al., 2010; Schmaljohann & Klinner, 2020). For instance, by radio tag-

ging multiple lean and fat individuals of a species on a single day, researchers can minimize the effect of weather variation on the birds' departure decision and separate out phenotypic, condition-related variation (e.g., Karwinkel et al., 2022, 2024). Subjecting numerous individuals to the same external conditions and tracking them at the same time may allow estimation of conditions when most individuals migrate (Delingat et al., 2008; Schmaljohann & Klinner, 2020), for example, during favorable winds (Lagerveld et al., 2024).

Parameters derived from flights of individuals tracked with Motus, such as departure and landing decisions, speed, and routes (Figure 5a; Brunner et al., 2022; Brust et al., 2019; Linhart et al., 2023; Packmor et al., 2020; Ruppel et al., 2023), can allow comparisons in behavior among individuals of different populations and those that orient across and around barriers (Figure 5b) (Brust & Hüppop, 2022; Schmaljohann & Naef-Daenzer, 2011; Woodworth et al., 2015). An improved understanding of migration behavior, its limitations, and flexibility among different species can help improve predictions of how species might adapt to changes around them and thus improve efforts toward their conservation (Sutherland, 1998).

Obtaining individual responses to environmental stressors

Motus data can also be used to identify issues of conservation concern and detect responses of migrating birds to specific forms of anthropogenic or environmental disruption. Anthropogenic structures, such as offshore wind turbines, can attract migrating birds, potentially causing increased mortality through collision (Perrow, 2019) or evoking avoidance behavior that can lead to increased or delayed fitness costs due to longer routes and higher energy expenditure (Schwemmer et al., 2023). Such impacts are still largely unquantified for migratory populations of birds (Marques et al., 2021). One possibility is to use Motus in combination with acoustic monitoring (as in Lagerveld et al. [2023]), whereby it is possible to localize the interaction of tracked individuals with near- and offshore infrastructure while also contextualizing these known individuals among con- and allospecifics detected by the acoustic recorders (Loring et al., 2019; Willmott et al., 2023).

Other anthropogenic disruptors are agrochemicals, such as neonicotinoids, which can impair the progress of migration in different taxa (Cabrera-Cruz et al., 2020). Eng et al. (2019) used Motus tracking data to show responses to neonicotinoid ingestion by white-crowned sparrows (*Zonotrichia leucophrys*). They showed that migrating birds on stopover were severely impaired in their ability to refuel, despite significantly increasing the length of stopover.

Artificial light at night (ALAN) can attract night-migrating birds to bright, often urban, areas (Horton et al., 2023; McLaren et al., 2018; Smith et al., 2021). These areas may act as ecological traps (i.e., inadequate stopover sites that might present higher risk of mortality [Van Doren et al., 2021]). Similarly, anthropogenic electromagnetic radiation (electrosmog) may disrupt the magnetic compass of night-migrating songbirds (Engels




FIGURE 5 (a) Currently operational Motus receiving stations (green dots) along the North Sea coast and examples of movement tracks of birds and bats and (b) examples of potential station placement (yellow dots) and data collection at Gibraltar, Iberian Peninsula, where many thousands of migratory species cross an important migratory barrier, the Mediterranean Sea (blue arrows, expected flight paths that could be detected by Motus).

et al., 2014). Engels et al. observed these responses in a laboratory environment with caged birds; whether electrosmog is also a hazard for freely moving birds in the wild remains to be tested. Here, researchers could apply Motus tracking, where directional and departure time data can be collected by local and regional arrays of receivers positioned in and around areas of high urban density.

Combining Motus tracking with physical samples

Simultaneously collecting samples (e.g., feathers, saliva, blood, or feces) that reveal something about the physiological state of

the animals, together with movement behavior, can improve understanding of how the physiology of an individual influences its migratory decisions. The high temporal resolution of Motus tracking data allows one to identify more closely links between physiological indicators, especially those changing rapidly (e.g., hormones), and species' movement (e.g., Eikenaar et al., 2020). This could, for example, include site quality by correlating stopover duration and habitat use, as recorded by Motus, with body condition and immune function (Brust et al., 2022; Hegemann et al., 2018; Schmaljohann & Naef-Daenzer, 2011). This would allow researchers to determine whether the sites provide the necessary functions for stopover. If not, targeted conservation measures could be taken to restore the missing functions.

Genetic analyses in conjunction with recorded migratory behavior (direction and routes, which are accessible with the high spatiotemporal accuracy of Motus) could indicate population-specific differences and possibly regions in the genetic structure that are important for the genetic coding of migratory behavior (Bossu et al., 2022; Ruegg et al., 2014; Sharma et al., 2023). Blood and fecal samples could be used to monitor the prevalence of pathogens that can be linked to body condition, population origin, and migration decisions (ideally seasonal migration success) (Morales et al., 2022; Neima et al., 2020). In the long term, standardized studies of migratory behavior combined with sampling of tagged individuals could allow predictions of responses to global climate and habitat changes (Anderson et al., 2019; Saura et al., 2014).

Logistics of developing Motus for flyway-level research

Achieving greater geographical (i.e., near-continental) coverage of the Motus network stations is underway. However, this requires a strategic placement plan (Lefevre & Smith, 2020) based around the key questions we have discussed and the special physical features of European landscapes (Figures 4 & 5). The network will require significant capital investment and a collaborative spirit among researchers, conservationists, and volunteers because this task is too big for single groups.

Single groups can realize regional-scale networks through discrete projects, which is a necessary way of completing a continent-wide network (Griffin et al., 2020; Taylor et al., 2017). Ideally, such projects fill in geographical gaps based on species ecology and migratory behavior already garnered from other technologies (e.g., geolocators [Bayly et al., 2018] or radar [Robinson, 2023]). As well as capital, the development of the network will require significant time and focus to maintain equipment and retrieve data, particularly in remote areas. Such receivers are less likely to be monitored remotely because of signal and power restrictions; therefore, greater logistical efforts are required to obtain the stored data and undertake maintenance.

Cost per receiver can be realized for as little as €3000–5000 (~4 directional antennas, Sensorgnome receiver), but it may approach €10000 depending on requirements for installation and precise configuration of antennas. Each tag, whether from CTT or Lotek, is approximately €200, although this approaches €300 for the very smallest models. Although cheaper than large, satellite-enabled tags, this does not approach the low cost of metal or color rings that allow researchers to capture and mark many thousands of birds. Cost reduction is hampered by limited market competition and a lack of open-source development, which contrasts with the collaborative nature of Motus entirely and must be addressed to allow tagging on a much larger scale.

Finally, the amount of data collected from Motus is enormous and is likely to continue to grow alongside other biologging data (López-López, 2016), so appropriate statistical tools will need to continue to be developed. Complex Bayesian modeling frame-

works to appropriately analyze Motus data have been developed and tested in limited circumstances (e.g., modeling movement offshore [Baldwin et al., 2018; Cranmer et al., 2017] and estimating flight heights [Lagerveld et al., 2024]). Extending the applicability of these methods and developing integrated frameworks with multiple data types would allow researchers to make better use of Motus data and make further inferences about migratory parameters that can inform conservation (Gregory et al., 2023).

These challenges can only be solved over the long term with a coordinated, international, collaborative effort to develop joint funding applications and to work together for the benefit of the wider Motus community. This community must contain academics, policy makers, government officials, conservationists, and amateur biologists and ecologists who can develop well-defined, focused study objectives. The involvement of a diverse number of stakeholders is required, not just to share the cost burden and coordination responsibilities but also to ensure fair data sharing and the direct integration of such data into policy and conservation actions (Gregory et al., 2023; Guilherme et al., 2023; UNEP, 2020).

FINAL OUTLOOK

In this time of rapid ecosystem disruption, it is vital to work collaboratively to conserve migratory species. Work needs to be done at multiple scales to answer questions about how species are confronting environmental changes. Motus can provide multiscale data on the movements of bird without the need for recapture, even on species as small as Nathusius's pipistrelles (*Pipistrellus nathusii*) (Bach et al., 2022; Briggs et al., 2023; Lagerveld et al., 2024), yellow-browed warbler *Phylloscopus inornatus*, and large insects, such as the monarch butterfly (*Danaus plexippus*) (Knight et al., 2019; Wilcox et al., 2021). Motus' features and capabilities make it an attractive and exciting prospect for exploring as-yet-unanswered ecological, evolutionary, and behavioral questions.












There is a significant amount of logistical and planning work to be done to develop and increase the size of the network so that it can reach its full potential in terms of basic and applied science. Such efforts should also focus on expanding the collaboration between parties and realizing conservation strategies that will benefit birds, nature as a whole, and ultimately, by supporting the One Health approach, humans.

ACKNOWLEDGMENTS

Funding to install Motus stations in Germany was provided by the Deutsche Forschungsgemeinschaft (DFG) within the Sonderforschungsbereich (SFB) 1372 Magnetoreception and Navigation in Vertebrates (project number 395940726) and other projects (SCHM 2647/3-1, SCHM 2647/4-1, SCHM 2647/7-1) and by the Federal Agency for Nature Conservation (BfN) with funds from the Federal Ministry for the Environment, Nature Conservation and Nuclear Safety (BMU), Germany (grants 351582210A and 351986140A to O. Hüppop and grant 352315100B to H.S.). In the United Kingdom,

funding for station installation and tags was from the Yorkshire Water Biodiversity Fund (YW 270521 Connected Ecology), the Scottish Ornithological Club, and the British Ecological Society (SR20/1532), as well as some very generous personal donations. We explicitly thank all team members behind the realized Motus stations in all countries in this collaboration (O. Hüppop, H.-H. Blikslager, T. Klinner, F. Matthews, T. Mertens, M. de Neidels, F. Packmor, E. Parsons, S. Parsons, A. Peter, G. Rüppel, Z. Züst) and the entire team behind the Motus Wildlife Tracking System for their invaluable technical support. We also thank J. Waldenstrom and N. Isaakson for comments before submission and W. Cresswell, N. Sapir, and one other anonymous reviewer for their comments on the manuscript.

ORCID

Lucy Mitchell  <https://orcid.org/0000-0003-2162-013X>
 Vera Brust  <https://orcid.org/0000-0003-4128-4495>
 Thimo Karwinkel  <https://orcid.org/0000-0001-6433-3013>
 Susanne Åkesson  <https://orcid.org/0000-0001-9039-2180>
 Dmitry Kishkinev  <https://orcid.org/0000-0002-2619-1197>
 Gabriel Norevik  <https://orcid.org/0000-0002-4112-0346>
 Tibor Szép  <https://orcid.org/0000-0002-4569-7425>
 Anders Hedenström  <https://orcid.org/0000-0002-1757-0945>
 Sander Lagerveld  <https://orcid.org/0000-0003-1291-4021>
 Barbara Helm  <https://orcid.org/0000-0002-6648-1463>
 Heiko Schmaljohann  <https://orcid.org/0000-0002-0886-4319>

REFERENCES

- Åkesson, S., & Hedenström, A. (2007). How migrants get there: Migratory performance and orientation. *Bioscience*, 57(2), 123–133.
- Åkesson, S., & Helm, B. (2020). Endogenous programs and flexibility in bird migration. *Frontiers in Ecology and Evolution*, 8, Article 78.
- Alerstam, T., Hedenström, A., & Åkesson, S. (2003). Long-distance migration: Evolution and determinants. *Oikos*, 103(2), 247–260.
- Anderson, A. M., Duijns, S., Smith, P. A., Friis, C., & Nol, E. (2019). Migration distance and body condition influence shorebird migration strategies and stopover decisions during southbound migration. *Frontiers in Ecology and Evolution*, 7, Article 251.
- Ayres, M. P., & Lombardero, M. J. (2018). Forest pests and their management in the Anthropocene. *Canadian Journal of Forest Research*, 48(3), 292–301.
- Bach, P., Voigt, C. C., Göttsche, M., Bach, L., Brust, V., Hill, R., Hüppop, O., Lagerveld, S., Schmaljohann, H., & Seebens-Hoyer, A. (2022). Offshore and coastline migration of radio-tagged *Nathusius' pipistrelles*. *Conservation Science and Practice*, 4(10), Article e12783.
- Baillie, S. R. (1995). Uses of ringing data for the conservation and management of bird populations: A ringing scheme perspective. *Journal of Applied Statistics*, 22(5-6), 967–988.
- Baldwin, J. W., Leap, K., Finn, J. T., & Smetzer, J. R. (2018). Bayesian state-space models reveal unobserved off-shore nocturnal migration from Motus data. *Ecological Modelling*, 386, 38–46.
- Barron, D. G., Brawn, J. D., & Weatherhead, P. J. (2010). Meta-analysis of transmitter effects on avian behaviour and ecology. *Methods in Ecology and Evolution*, 1, 180–187.
- Bauer, S., & Hoyer, B. J. (2014). Migratory animals couple biodiversity and ecosystem functioning worldwide. *Science*, 344(6179), Article 1242552.
- Bayly, N. J., Gómez, C., & Hobson, K. A. (2013). Energy reserves stored by migrating Gray-cheeked Thrushes *Catharus minimus* at a spring stopover site in northern Colombia are sufficient for a long-distance flight to North America. *Ibis*, 155, 271–283.
- Bayly, N. J., Rosenberg, K. V., Easton, W. E., Gomez, C., Carlisle, J. A. Y., Ewert, D. N., Drake, A., & Goodrich, L. (2018). Major stopover regions and migratory bottlenecks for Nearctic-Neotropical landbirds within the Neotropics: A review. *Bird Conservation International*, 28(1), 1–26.
- Beardsworth, C. E., Gobbens, E., van Maarseveen, F., Denissen, B., Dekinga, A., Nathan, R., ... Bijleveld, A. I. (2022). Validating ATLAS: A regional-scale high-throughput tracking system. *Methods in Ecology and Evolution*, 13(9), 1990–2004.
- Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., & Courchamp, F. (2012). Impacts of climate change on the future of biodiversity. *Ecology Letters*, 15(4), 365–377.
- Birnie-Gauvin, K., Lennox, R. J., Guglielmo, C. G., Teffer, A. K., Crossin, G. T., Norris, D. R., Aarestrup, K., & Cooke, S. J. (2020). The value of experimental approaches in migration biology. *Physiological and Biochemical Zoology*, 93(3), 210–226.
- Border, J. A., Henderson, I. G., Ash, D., & Hartley, I. R. (2017). Characterising demographic contributions to observed population change in a declining migrant bird. *Journal of Avian Biology*, 48, 1139–1149.
- Bossu, C. M., Heath, J. A., Kaltenecker, G. S., Helm, B., & Ruegg, K. C. (2022). Clock-linked genes underlie seasonal migratory timing in a diurnal raptor. *Proceedings of the Royal Society B: Biological Sciences*, 289(1974), Article 20212507.
- Both, C., Bouwuis, S., Lessells, C. M., & Visser, M. E. (2006). Climate change and population declines in a long-distance migratory bird. *Nature*, 441(7089), 81–83.
- Bridge, E., Thorup, K., Bowlin, M., Chilson, P., Diehl, R., Fléron, R., Hartl, P., Kays, R., Kelly, J., Robinson, W., & Wikelski, M. (2011). Technology on the move: Recent and forthcoming innovations for tracking migratory birds. *Bioscience*, 61(9), 689–698.
- Briggs, P., Briggs, B., & Parsons, E. A. (2023). First Motus tag results from Bedford Lakes study of *Nathusius' pipistrelle*. *British Islands Bats*, 4, 46–60.
- Brown, J. M., & Taylor, P. D. (2015). Adult and hatch-year blackpoll warblers exhibit radically different regional-scale movements during post-fledging dispersal. *Biology Letters*, 11(12), Article 20150593.
- Brown, J. M., & Taylor, P. D. (2017). Migratory blackpoll warblers (*Setophaga striata*) make regional-scale movements that are not oriented toward their migratory goal during fall. *Movement Ecology*, 5, Article 15.
- Bruderer, B., & Jenni, L. (1990). Migration across the Alps. In E. Gwinner (Ed.), *Bird migration: Physiology and ecophysiology* (pp. 61–77). Springer Verlag.
- Bruderer, B., & Liechti, F. (1999). Bird migration across the Mediterranean. In N. Adams & R. Slotow (Eds.), *Proceedings of the International Ornithological Congress, Durban* (pp. 1983–1999). BirdLife South Africa.
- Brunner, A. R., Dossman, B. C., Jirinec, V., Percy, K. L., Tonra, C. M., Johnson, E. I., & Marra, P. P. (2022). Migratory behavior and connectivity revealed in a secretive Neotropical migratory songbird, the Swainson's Warbler. *Journal of Field Ornithology*, 93(3), Article 5.
- Brust, V., Eikenaar, C., Packmor, F., Schmaljohann, H., Hüppop, O., & Czirják, G. Á. (2022). Do departure and flight route decisions correlate with immune parameters in migratory songbirds? *Functional Ecology*, 36, 3007–3021.
- Brust, V., & Hüppop, O. (2022). Underestimated scale of songbird offshore migration across the south-eastern North Sea during autumn. *Journal of Ornithology*, 163, 51–60.
- Brust, V., Michalik, B., & Hüppop, O. (2019). To cross or not to cross—Thrushes at the German North Sea coast adapt flight and routing to wind conditions in autumn. *Movement Ecology*, 7, Article 32.
- Buechley, E. R., Oppel, S., Efrat, R., Phipps, W. L., Carbonell Alanis, I., Álvarez, E., ... Marra, P. P. (2021). Differential survival throughout the full annual cycle of a migratory bird presents a life-history trade-off. *Journal of Animal Ecology*, 90(5), 1228–1238.
- Cabrera-Cruz, S. A., Cohen, E. B., Smolinsky, J. A., & Buler, J. J. (2020). Artificial light at night is related to broad-scale stopover distributions of nocturnally migrating landbirds along the Yucatan Peninsula, Mexico. *Remote Sensing*, 12, Article 395.
- Chowdhury, S., Jennions, M. D., Zalucki, M. P., Maron, M., Watson, J. E., & Fuller, R. A. (2023). Protected areas and the future of insect conservation. *Trends in Ecology & Evolution*, 38(1), 85–95.
- Clarke, B., Otto, F., Stuart-Smith, R., & Harrington, L. (2022). Extreme weather impacts of climate change: An attribution perspective. *Environmental Research: Climate*, 1(1), Article 012001.

- Cohen, E. B., & Satterfield, D. A. (2020). 'Chancing on a spectacle': co-occurring animal migrations and interspecific interactions. *Ecography*, 43(11), 1657–1671.
- Cooper, N. W., Haradon, H. A., Hartman, J. P., Hayden, B. R., Kersh, L., & Riley, A. (2024). Preliminary evaluation of habitat use, pairing rate, and reproductive success in an experimental breeding habitat planted for Kirtland's Warblers (*Setophaga kirtlandii*). *Journal of Field Ornithology*, 95(3), Article 3.
- Cooper, N. W., & Marra, P. P. (2020). Hidden long-distance movements by a migratory bird. *Current Biology*, 30, 4056.e3–4062.e3.
- Cox, A. S., & Kesler, D. C. (2012). Reevaluating the cost of natal dispersal: Post-fledging survival of red-bellied woodpeckers. *The Condor*, 114(2), 341–347.
- Cranmer, A., Smetzer, J. R., Welch, L., & Baker, E. (2017). A Markov model for planning and permitting offshore wind energy: A case study of radio-tracked terns in the Gulf of Maine, USA. *Journal of Environmental Management*, 193, 400–409.
- Crewe, T. L., Kendal, D., & Campbell, H. A. (2020). Motivations and fears driving participation in collaborative research infrastructure for animal tracking. *PLoS ONE*, 15, Article e0241964.
- Day, G., Fox, G., Hipperson, H., Maher, K., Tucker, R., Horsburgh, G., Waters, D., Durrant, K. L., Arnold, K., Burke, T., & Slate, J. (2024). Revealing the demographic history of the European nightjar (*Caprimulgus europaeus*). *Ecology and Evolution*, 14(10), Article e70460.
- Delingat, J., Bairlein, F., & Hedenström, A. (2008). Obligatory barrier crossing and adaptive fuel management in migratory birds: The case of the Atlantic crossing in Northern Wheatears (*Oenanthe oenanthe*). *Behavioral Ecology and Sociobiology*, 62, 1069–1078.
- DeMars, C. A., Johnson, C. J., Dickie, M., Habib, T. J., Cody, M., Saxena, A., Boutin, S., & Serrouya, R. (2023). Incorporating mechanism into conservation actions in an age of multiple and emerging threats: The case of boreal caribou. *Ecosphere*, 14(7), Article e4627.
- Dhanjal-Adams, K. L., Klaassen, M., Nicol, S., Possingham, H. P., Chadès, I., & Fuller, R. A. (2017). Setting conservation priorities for migratory networks under uncertainty. *Conservation Biology*, 31, 646–656.
- Doerr, E. D., & Doerr, V. A. J. (2005). Dispersal range analysis: Quantifying individual variation in dispersal behaviour. *Oecologia*, 142, 1–10.
- Driscoll, D. A., Banks, S. C., Barton, P. S., Ikin, K., Lentini, P., Lindenmayer, D. B., Smith, A. L., Berry, L. E., Burns, E. L., Edworthy, A., & Evans, M. J. (2014). The trajectory of dispersal research in conservation biology. Systematic review. *PLoS ONE*, 9(4), Article e95053.
- Du Feu, C. R., Clark, J. A., Schaub, M., Fiedler, W., & Baillie, S. R. (2016). The EURING Data Bank—A critical tool for continental-scale studies of marked birds. *Ringed & Migration*, 31(1), 1–18.
- Dufour, P., Åkesson, S., Hellström, M., Hewson, C., Lagerveld, S., Mitchell, L., Chernetsov, N., Schmaljohann, H., & Crochet, P. A. (2022). The Yellow-browed Warbler (*Phylloscopus inornatus*) as a model to understand vagrancy and its potential for the evolution of new migration routes. *Movement Ecology*, 10(1), Article 59.
- Dufour, P., de Franceschi, C., Doniol-Valcroze, P., Jiguet, F., Guéguen, M., Renaud, J., Lavergne, S., & Crochet, P. A. (2021). A new westward migration route in an Asian passerine bird. *Current Biology*, 31(24), 5590–5596.
- Eikenaar, C., Schaefer, J., Hessler, S., Packmor, F., & Schmaljohann, H. (2020). Diel variation in corticosterone and departure decision making in migrating birds. *Hormones and Behavior*, 122, Article 104746.
- Endriss, S. B., Vahsen, M. L., Bitume, E. V., Grey Monroe, J., Turner, K. G., Norton, A. P., & Hufbauer, R. A. (2019). The importance of growing up: Juvenile environment influences dispersal of individuals and their neighbours. *Ecology Letters*, 22, 45–55.
- Eng, M. L., Stutchbury, B. J., & Morrissey, C. A. (2019). A neonicotinoid insecticide reduces fueling and delays migration in songbirds. *Science*, 365(6458), 1177–1180.
- Engels, S., Schneider, N. L., Lefeldt, N., Hein, C. M., Zapka, M., Michalik, A., Elbers, D., Kittel, A., Hore, P. J., & Mouritsen, H. (2014). Anthropogenic electromagnetic noise disrupts magnetic compass orientation in a migratory bird. *Nature*, 509(7500), 353–356.
- Evans, D. R. (2018). *The post-fledging survival and movements of juvenile Barn Swallows (Hirundo rustica): An automated telemetry approach* (Master's thesis). The University of Western Ontario.
- Evans, D. R., Hobson, K. A., Kusack, J. W., Cadman, M. D., Falconer, C. M., & Mitchell, G. W. (2020). Individual condition, but not fledging phenology, carries over to affect post-fledging survival in a Neotropical migratory songbird. *Ibis*, 162(2), 331–344.
- Evans, D. R. (2018). The post-fledging survival and movements of juvenile Barn Swallows (*Hirundo rustica*): an automated telemetry approach. Master's thesis, The University of Western Ontario, Canada.
- Fattorini, N., Costanzo, A., Romano, A., Rubolini, D., Baillie, S., Bairlein, F., Spina, F., & Ambrosini, R. (2023). Eco-evolutionary drivers of avian migratory connectivity. *Ecology Letters*, 26(7), 1095–1107.
- Fiedler, W. (2009). New technologies for monitoring bird migration and behaviour. *Ringed & Migration*, 24(3), 175–179.
- Flack, A., Aikens, E. O., Kölzsch, A., Nourani, E., Snell, K. R., Fiedler, W., Linck, N., Bauer, H. G., Thorup, K., Partecke, J., Wikelski, M., & Williams, H. J. (2022). New frontiers in bird migration research. *Current Biology*, 32(20), R1187–R1199.
- Frick, W. F., Kingston, T., & Flanders, J. (2020). A review of the major threats and challenges to global bat conservation. *Annals of the New York Academy of Sciences*, 1469(1), 5–25.
- Gauld, J. G., Silva, J. P., Atkinson, P. W., Record, P., Acácio, M., Arkumarev, V., Blas, J., Bouten, W., Burton, N., Catry, I., & Champagnon, J. (2022). Hotspots in the grid: Avian sensitivity and vulnerability to collision risk from energy infrastructure interactions in Europe and North Africa. *Journal of Applied Ecology*, 59(6), 1496–1512.
- Gómez, C., Bayly, N. J., & Rosenberg, K. V. (2014). Fall stopover strategies of three species of thrush (*Catharus*) in northern South America. *Auk*, 131, 702–717.
- Gómez, C., Hobson, K. A., Bayly, N. J., Rosenberg, K. V., Morales-Rozo, A., Cardozo, P., & Cadena, C. D. (2021). Migratory connectivity then and now: A northward shift in breeding origins of a long-distance migratory bird wintering in the tropics. *Proceedings of the Royal Society B: Biological Sciences*, 288(1948), Article 20210188.
- González, A. M., Bayly, N. J., Wilson, S., & Hobson, K. A. (2021). Shade coffee or native forest? Indicators of winter habitat quality for a long-distance migratory bird in the Colombian Andes. *Ecological Indicators*, 131, Article 108115.
- Gottwald, J., Royauté, R., Becker, M., Geitz, T., Höchst, J., Lampe, P., ... Nauss, T. (2023). Classifying the activity states of small vertebrates using automated VHF telemetry. *Methods in ecology and evolution*, 14(1), 252–264.
- Goymann, W., Spina, F., Ferri, A., & Fusani, L. (2010). Body fat influences departure from stopover sites in migratory birds: Evidence from whole-island telemetry. *Biological Letters*, 6(4), 478–481.
- Gregory, K. A., Francesiaz, C., Jiguet, F., & Besnard, A. (2023). A synthesis of recent tools and perspectives in migratory connectivity studies. *Movement Ecology*, 11(1), Article 69.
- Griffin, A. S., Brown, C., Woodworth, B. K., Ballard, G. A., Blanch, S., Campbell, H. A., Crewe, T. L., Hansbro, P. M., Herbert, C. A., Hosking, T., Hoyer, B. J., Law, B., Leigh, K., Machovsky-Capuska, G. E., Rasmussen, T., McDonald, P. G., Roderick, M., Slade, C., Mackenzie, S. A., & Taylor, P. D. (2020). A large-scale automated radio telemetry network for monitoring movements of terrestrial wildlife in Australia. *Australian Zoologist*, 40(3), 379–391.
- Guilherme, J. L., Jones, V. R., Catry, I., Beal, M., Dias, M. P., Oppel, S., Vickery, J. A., Hewson, C. M., Butchart, S. H. M., & Rodrigues, A. S. (2023). Connectivity between countries established by landbirds and raptors migrating along the African–Eurasian flyway. *Conservation Biology*, 37(1), Article e14002.
- Hagelin, J. C., Hallworth, M. T., Barger, C. P., Johnson, J. A., DuBour, K. A., Pendelton, G. W., ... Taylor, A. (2021). Revealing migratory path, important stopovers and non-breeding areas of a boreal songbird in steep decline. *Animal Migration*, 8(1), 168–191.
- Hayes, S., Boyd, B. P., & Stutchbury, B. J. (2024). Why do juvenile Wood Thrushes make long-distance pre-migratory movements across a fragmented landscape? *Journal of Field Ornithology*, 95(2), Article 9.
- Hedenström, A., & Pettersson, J. (1987). Migration routes and wintering areas of willow warblers *Phylloscopus trochilus* (L.) ringed in Fennoscandia. *Ornis Fennica*, 64(4), 137–143.
- Hegemann, A., Alcalde Abril, P., Sjöberg, S., Muheim, R., Alerstam, T., Nilsson, J. Å., & Hasselquist, D. (2018). A mimicked bacterial infection prolongs

- stopover duration in songbirds—But more pronounced in short- than long-distance migrants. *Journal of Animal Ecology*, 87(6), 1698–1708.
- Herbert, J. A., Mizrahi, D., & Taylor, C. M. (2022). Migration tactics and connectivity of a Nearctic–Neotropical migratory shorebird. *Journal of Animal Ecology*, 91(4), 819–830.
- Hewson, C. M., Thorup, K., Pearce-Higgins, J. W., & Atkinson, P. W. (2016). Population decline is linked to migration route in the Common Cuckoo. *Nature Communications*, 7(1), Article 12296.
- Hirschhofer, S., Liechti, F., Ranacher, P., Weibel, R., & Schmid, B. (2024). High-intensity bird migration along Alpine valleys calls for protective measures against anthropogenically induced avian mortality. *Remote Sensing in Ecology and Conservation*, 10(3), 360–373.
- Holberton, R. L., Taylor, P. D., Tudor, L. M., O'Brien, K. M., Mittelhauser, G. H., & Breit, A. (2019). Automated VHF radiotelemetry revealed site-specific differences in fall migration strategies of Semipalmated Sandpipers on stopover in the Gulf of Maine. *Frontiers in Ecology and Evolution*, 7, Article 327.
- Horton, K. G., Buler, J. J., Anderson, S. J., Burt, C. S., Collins, A. C., Dokter, A. M., Guo, F., Sheldon, D., Tomaszewska, M. A., & Henebry, G. M. (2023). Artificial light at night is a top predictor of bird migration stopover density. *Nature Communications*, 14(1), Article 7446.
- Horton, K. G., La Sorte, F. A., Sheldon, D., Lin, T. Y., Winner, K., Bernstein, G., Maji, S., Hochachka, W. M., & Farnsworth, A. (2020). Phenology of nocturnal avian migration has shifted at the continental scale. *Nature Climate Change*, 10(1), 63–68.
- Howard, C., Stephens, P. A., Pearce-Higgins, J. W., Gregory, R. D., Butchart, S. H., & Willis, S. G. (2020). Disentangling the relative roles of climate and land cover change in driving the long-term population trends of European migratory birds. *Diversity and Distributions*, 26(11), 1442–1455.
- Iler, A. M., CaraDonna, P. J., Forrest, J. R., & Post, E. (2021). Demographic consequences of phenological shifts in response to climate change. *Annual Review of Ecology, Evolution, and Systematics*, 52, 221–245.
- Imlay, T. L., Saldanha, S., & Taylor, P. D. (2020). The fall migratory movements of bank swallows, *Riparia riparia*: Fly-and-forage migration? *Avian Conservation and Ecology*, 15, Article 2.
- Jaureguiberry, P., Titeux, N., Wiemers, M., Bowler, D. E., Coscieme, L., Golden, A. S., Guerra, C. A., Jacob, U., Takahashi, Y., Settele, J., & Díaz, S. (2022). The direct drivers of recent global anthropogenic biodiversity loss. *Science Advances*, 8(45), Article eabm9982.
- Jiguet, F., Robert, A., Lorrillière, R., Hobson, K. A., Kardynal, K. J., Arlettaz, R., Bairlein, F., Belik, V., Bernardy, P., Copete, J. L., & Moussy, C. (2019). Unravelling migration connectivity reveals unsustainable hunting of the declining ortolan bunting. *Science Advances*, 5(5), Article eaau2642.
- Karwinkel, T., Winkhofer, M., Allenstein, D., Brust, V., Christoph, P., Holland, R. A., Hüppop, O., Steen, J., Bairlein, F., & Schmaljohann, H. (2024). A refined magnetic pulse treatment method for magnetic navigation experiments with adequate sham control: A case study on free-flying songbirds. *Journal of the Royal Society Interface*, 21(214), Article 20230745.
- Karwinkel, T., Winkhofer, M., Janner, L. E., Brust, V., Hüppop, O., Bairlein, F., & Schmaljohann, H. (2022). A magnetic pulse does not affect free-flight navigation behaviour of a medium-distance songbird migrant in spring. *Journal of Experimental Biology*, 225(19), Article jeb244473.
- Klaassen, R. H. G., Hake, M., Strandberg, R., Koks, B. J., Trierweiler, C., Exo, K.-M., Bairlein, F., & Alerstam, T. (2014). When and where does mortality occur in migratory birds? Direct evidence from long-term satellite tracking of raptors. *Journal of Animal Ecology*, 83, 176–184.
- Knight, S. M., Pitman, G. M., Flockhart, D. T., & Norris, D. R. (2019). Radio-tracking reveals how wind and temperature influence the pace of daytime insect migration. *Biology Letters*, 15(7), Article 20190327.
- Lagerveld, S., de Vries, P., Harris, J., Parsons, S., Debusschere, E., Hüppop, O., Brust, V., & Schmaljohann, H. (2024). Migratory movements of bats are shaped by barrier effects, sex-biased timing and the adaptive use of winds. *Movement Ecology*, 12(1), Article 81.
- Lagerveld, S., Wilkes, T., van Puijenbroek, M. E., Noort, B. C., & Geelhoed, S. C. (2023). Acoustic monitoring reveals spatiotemporal occurrence of *Nathusius' pipistrelle* at the southern North Sea during autumn migration. *Environmental Monitoring and Assessment*, 195(9), Article 1016.
- Lees, A. C., & Gilroy, J. J. (2009). Vagrancy mechanisms in passerines and near-passerines. In R. Slack (Ed.), *Rare birds, where and when: An analysis of status and distribution in Britain and Ireland* (Vol. 1, pp. 1–23). Rare Bird Books.
- Lefevre, K. L., & Smith, A. D. (2020). Florida's strategic position for collaborative automated telemetry tracking of avian movements across the Americas. *Journal of Fish and Wildlife Management*, 11(1), 369–375.
- le Roux, C. E., & Nocera, J. J. (2021). Roost sites of chimney swift (*Chaetura pelagica*) form large-scale spatial networks. *Ecology & Evolution*, 11(9), 3820–3829.
- Linhart, R. C., Hamilton, D. J., Paquet, J., & Gratto-Trevor, C. L. (2023). Evidence of differing staging strategies between adult and juvenile Semipalmated Sandpipers highlights the importance of small staging sites in Atlantic Canada. *Ornithology*, 140(1), Article ukac056.
- Linscott, J. A., & Senner, N. R. (2021). Beyond refueling: Investigating the diversity of functions of migratory stopover events. *The Condor: Ornithological Applications*, 123, Article duaa074.
- López-López, P. (2016). Individual-based tracking systems in ornithology: Welcome to the era of big data. *Ardeola*, 63, 103–136.
- Loring, P. H., Paton, P. W. C., McLaren, J. D., Bai, H., Janaswamy, R., Goyert, H. F., Griffin, C. R., & Sievert, P. R. (2019). *Tracking offshore occurrence of common terns, endangered roseate terns, and threatened piping plovers with VHF arrays* (OCS Study BOEM 2019–017). US Department of the Interior, Bureau of Ocean Energy Management.
- Machado, R., Nabo, P., Cardia, P., Moreira, P., Nicolau, P., & Repas-Goncalves, M. (2024). *Bird Curtailment in Offshore Wind Farms: Application of curtailment in offshore wind farms at a sea basin level to mitigate collision risk for birds*. Birdlife Europe and Central Asia and STRIX.
- Marcacci, G., Briedis, M., Diop, N., Diallo, A. Y., Kebede, F., & Jacot, A. (2023). A roadmap integrating research, policy, and actions to conserve Afro-Palaearctic migratory landbirds at a flyway scale. *Conservation Letters*, 16(1), e12933.
- Marques, A. T., Batalha, H., & Bernardino, J. (2021). Bird displacement by wind turbines: Assessing current knowledge and recommendations for future studies. *Birds*, 2(4), 460–475.
- Martell, M. S., Archer, A. A., Burnette, A., Lee, M., Nicoletti, F., & Hall, K. A. (2023). Using the Motus system to track post-breeding dispersal of American Kestrels nesting in Minnesota, USA. *Journal of Raptor Research*, 57(2), 154–163.
- Matthysen, E., & Clobert, J. (2012). Multicausality of dispersal: A review. In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal ecology and evolution* (pp. 3–18). Oxford University Press.
- McKinnon, E. A., Laplante, M. P., Love, O. P., Fraser, K. C., Mackenzie, S., & Vézina, F. (2019). Tracking landscape-scale movements of snow buntings and weather-driven changes in flock composition during the temperate winter. *Frontiers in Ecology and Evolution*, 7, Article 329.
- McKinnon, E. A., & Love, O. P. (2018). Ten years tracking the migrations of small landbirds: Lessons learned in the golden age of bio-logging. *Auk: Ornithological Advances*, 135(4), 834–856.
- McLaren, J. D., Buler, J. J., Schreckengost, T., Smolinsky, J. A., Boone, M., Emiel van Loon, E., Dawson, D. K., & Walters, E. L. (2018). Artificial light at night confounds broad-scale habitat use by migrating birds. *Ecology Letters*, 21(3), 356–364.
- Mitchell, G. W., Woodworth, B. K., Taylor, P. D., & Norris, D. R. (2015). Automated telemetry reveals age specific differences in flight duration and speed are driven by wind conditions in a migratory songbird. *Movement Ecology*, 3, Article 19.
- Moore, F. R. (2018). Biology of landbird migrants: a stopover perspective. *The Wilson Journal of Ornithology*, 130(1), 1–12.
- Morales, A., Frei, B., Mitchell, G. W., Bégin-Marchand, C., & Elliott, K. H. (2022). Reduced diurnal activity and increased stopover duration by molting Swainson's Thrushes. *Ornithology*, 139(2), Article ukab083.
- Morrison, C. A., Robinson, R. A., Butler, S. J., Clark, J. A., & Gill, J. A. (2016). Demographic drivers of decline and recovery in an Afro-Palaearctic migratory bird population. *Proceedings of the Royal Society B: Biological Sciences*, 283(1842), Article 20161387.
- Mukhin, A., Kobylkov, D., Kishkinev, D., & Grinkevich, V. (2018). Interrupted breeding in a songbird migrant triggers development of nocturnal locomotor activity. *Scientific Reports*, 8(1), Article 5520.

- Mukhin, A., Kosarev, V., & Ktitorov, P. (2005). Nocturnal life of young songbirds well before migration. *Proceedings of the Royal Society B: Biological Sciences*, 272(1572), 1535–1539.
- Nadal, J., Ponz, C., Margalida, A., & Pennisi, L. (2020). Ecological markers to monitor migratory bird populations: Integrating citizen science and trans-boundary management for conservation purposes. *Journal of Environmental Management*, 255, Article 109875.
- Neima, S. G., Hamilton, D. J., Gratto-Trevor, C. L., & Paquet, J. (2020). Intra-and interannual regional fidelity of Semipalmated Sandpipers (*Calidris pusilla*) during migratory stopover in the upper Bay of Fundy, Canada. *Avian Conservation & Ecology*, 15(1).
- Niebuhr, B. B. S., Wosniack, M. E., Santos, M. C., Raposo, E. P., Viswanathan, G. M., Da Luz, M. G., & Pie, M. R. (2015). Survival in patchy landscapes: The interplay between dispersal, habitat loss and fragmentation. *Scientific Reports*, 5(1), Article 11898.
- Nilsson, C., Dokter, A. M., Verlinden, L., Shamoun-Baranes, J., Schmid, B., Desmet, P., Bauer, S., Chapman, J., Alves, J. A., Stepanian, P. M., Sapir, N., Wainwright, C., Boos, M., Górska, A., Menz, M. H. M., Rodrigues, P., Leijnse, H., Zehindjiev, P., Brabant, R., ... Liechti, F. (2019). Revealing patterns of nocturnal migration using the European weather radar network. *Ecography*, 42(5), 876–886.
- Nussbaumer, R., Gravey, M., Briedis, M., & Liechti, F. (2023). Global positioning with animal-borne pressure sensors. *Methods in Ecology and Evolution*, 14(4), 1104–1117.
- Packmor, F., Klinner, T., Woodworth, B. K., Eikenaar, C., & Schmaljohann, H. (2020). Departure decisions in songbirds following different migration strategies: Long-distance migrants depart earlier and more independently of weather conditions than medium-distance migrants. *Movement Ecology*, 8, Article 6.
- Perrow, M. (Ed.). (2019). *Wildlife and wind farms - Conflicts and solutions: Volume 3: Offshore: Potential effects*. Pelagic Publishing Ltd.
- Rigal, S., Dakos, V., Alonso, H., Aunipš, A., Benkő, Z., Brotons, L., Chodkiewicz, T., Chylarecki, P., De Carli, E., Del Moral, J. C., & Domşa, C. (2023). Farmland practices are driving bird population decline across Europe. *Proceedings of the National Academy of Sciences of the United States of America*, 120(21), Article e2216573120.
- Robinson, S. K. (2023). Radar ornithology, stopover hotspots, and the conservation of migratory landbirds. *Proceedings of the National Academy of Sciences of the United States of America*, 120(4), Article e2220980120.
- Ruegg, K. C., Anderson, E. C., Paxton, K. L., Apkenas, V., Lao, S., Siegel, R. B., DeSante, D. F., Moore, F., & Smith, T. B. (2014). Mapping migration in a songbird using high-resolution genetic markers. *Molecular Ecology*, 23(23), 5726–5739.
- Rüppel, G., Hüppop, O., Schmaljohann, H., & Brust, V. (2023). The urge to breed early: Similar responses to environmental conditions in short- and long-distance migrants during spring migration. *Ecology and Evolution*, 13(7), Article e10223.
- Sala, O. E., Stuart Chapin, F. I. I., Armesto, J. J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-Sanwald, E., Huenneke, L. F., Jackson, R. B., Kinzig, A., & Leemans, R. (2000). Global biodiversity scenarios for the year 2100. *Science*, 287(5459), 1770–1774.
- Sandercock, B. K., Murray, D. L., & Sandercock, B. K. (2020). Mark-recapture models for estimation of demographic parameters. In D. L. Murray & B. K. Sandercock (Eds.), *Population ecology in practice* (pp. 157–190). Wiley-Blackwell.
- Satterfield, D. A., Sillett, T. S., Chapman, J. W., Altizer, S., & Marra, P. P. (2020). Seasonal insect migrations: massive, influential, and overlooked. *Frontiers in Ecology and the Environment*, 18(6), 335–344.
- Saura, S., Bodin, Ö., & Fortin, M. J. (2014). EDITOR'S CHOICE: Stepping stones are crucial for species' long-distance dispersal and range expansion through habitat networks. *Journal of Applied Ecology*, 51(1), 171–182.
- Schaub, M., Jakober, H., & Stauber, W. (2013). Strong contribution of immigration to local population regulation: Evidence from a migratory passerine. *Ecology*, 94(8), 1828–1838.
- Schaub, M., Reichlin, T. S., Abadi, F., Kéry, M., Jenni, L., & Arlettaz, R. (2012). The demographic drivers of local population dynamics in two rare migratory birds. *Oecologia*, 168, 97–108.
- Schmaljohann, H., & Eikenaar, C. (2017). How do energy stores and changes in these affect departure decisions by migratory birds? A critical view on stopover ecology studies and some future perspectives. *Journal of Comparative Physiology A*, 203, 411–429.
- Schmaljohann, H., Eikenaar, C., & Sapir, N. (2022). Understanding the ecological and evolutionary function of stopover in migrating birds. *Biological Reviews*, 97(4), 1231–1252.
- Schmaljohann, H., & Klinner, T. (2020). A quasi-experimental approach using telemetry to assess migration-strategy-specific differences in the decision-making processes at stopover. *BMC Ecology*, 20, Article 36.
- Schmaljohann, H., Liechti, F., Bächler, E., Steuri, T., & Bruderer, B. (2008). Quantification of bird migration by radar—A detection probability problem. *Ibis*, 150(2), 342–355.
- Schmaljohann, H., & Naef-Daenzer, B. (2011). Body condition and wind support initiate the shift of migratory direction and timing of nocturnal departure in a songbird. *Journal of Animal Ecology*, 80(6), 1115–1122.
- Schwemmer, P., Mercker, M., Haecker, K., Kruckenberg, H., Kämpfer, S., Bocher, P., ... Garthe, S. (2023). Behavioral responses to offshore windfarms during migration of a declining shorebird species revealed by GPS-telemetry. *Journal of Environmental Management*, 342, 118131.
- Serratos, J., Oppel, S., Rotics, S., Santangeli, A., Butchart, S. H., Cano-Alonso, L. S., Tellería, J. L., Kemp, R., Nicholas, A., Kalvāns, A., & Galarza, A. (2024). Tracking data highlight the importance of human-induced mortality for large migratory birds at a flyway scale. *Biological Conservation*, 293, Article 110525.
- Sharma, A., Sur, S., Tripathi, V., & Kumar, V. (2023). Genetic control of avian migration: Insights from studies in latitudinal passerine migrants. *Genes*, 14(6), Article 1191.
- Sjöberg, S., Alerstam, T., Åkesson, S., Schulz, A., Weidauer, A., Coppack, T., & Muheim, R. (2015). Weather and fuel reserves determine departure and flight decisions in passerines migrating across the Baltic Sea. *Animal Behaviour*, 104, 59–68.
- Smetzer, J. R., & King, D. I. (2018). Prolonged stopover and consequences of migratory strategy on local-scale movements within a regional songbird staging area. *The Auk: Ornithological Advances*, 135(3), 547–560.
- Smith, R. A., Gagné, M., & Fraser, K. C. (2021). Pre-migration artificial light at night advances the spring migration timing of a trans-hemispheric migratory songbird. *Environmental Pollution*, 269, Article 116136.
- Snell, K. R. S., Stokke, B. G., Moksnes, A., Thorup, K., & Fosøy, F. (2018). From Svalbard to Siberia: Passerines breeding in the High Arctic also endure the extreme cold of the Western Steppe. *PLoS ONE*, 13(9), Article e0202114.
- Soulsbury, C., Gray, H., Smith, L., Braithwaite, V., Cotter, S., Elwood, R. W., Wilkinson, A., & Collins, L. M. (2020). The welfare and ethics of research involving wild animals: A primer. *Methods in Ecology and Evolution*, 11(10), 1164–1181.
- Spina, F., Baillie, S. R., Bairlein, F., Ambrosini, R., Fattorini, N., Fiedler, W., Franks, S., Funghi, C., & Thorup, K. (2022). *Eurasian African Bird Migration Atlas—Executive summary*. CMS Secretariat.
- Sutherland, W. J. (1998). Evidence for flexibility and constraint in migration systems. *Journal of Avian Biology*, 29(4), 441–446.
- Taylor, P. D., Crewe, T. L., Mackenzie, S. A., Lepage, D., Aubry, Y., Crysler, Z., Finney, G., Francis, C. M., Guglielmo, C. G., Hamilton, D. J., Holberton, R. L., Loring, P. H., Mitchell, G. W., Norris, D. R., Paquet, J., Ronconi, R. A., Smetzer, J. R., Smith, P. A., Welch, L. J., & Woodworth, B. K. (2017). The Motus Wildlife Tracking System: A collaborative research network. *Avian Conservation and Ecology*, 12(1), Article 8.
- Taylor, P. D., Mackenzie, S. A., Thorber, B. G., Calvert, A. M., Mills, A. M., McGuire, L. P., & Guglielmo, C. G. (2011). Landscape movements of migratory birds and bats reveal an expanded scale of stopover. *PLoS ONE*, 6(11), Article e27054.
- Thorup, K., Korner-Nievergelt, F., Cohen, E. B., & Baillie, S. R. (2014). Large-scale spatial analysis of ringing and re-encounter data to infer movement patterns: A review including methodological perspectives. *Methods in Ecology and Evolution*, 5(12), 1337–1350.
- Travis, J. M. J., & Dytham, C. (2013). Dispersal and climate change: A review of theory. In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal ecology and evolution* (pp. 337–348). Oxford University Press.
- Turbek, S. P., Scordato, E. S. C., & Safran, R. J. (2018). The role of seasonal migration in population divergence and reproductive isolation. *Trends in Ecology and Evolution*, 33(3), 164–175.

- United Nations Environment Programme (UNEP). (2020). *Convention on Migratory Species (CMS) Resolution 12.6 (Rev. COP13): Improving ways of addressing connectivity in the conservation of migratory species*. https://www.cms.int/sites/default/files/document/cms_cop13_res.12.26_rev.cop13_e.pdf
- van Doren, B. M., Willard, D. E., Hennen, M., Horton, K. G., Stuber, E. F., Sheldon, D., Sivakumar, A. H., Wang, J., Farnsworth, A., & Winger, B. M. (2021). Drivers of fatal bird collisions in an urban center. *Proceedings of the National Academy of Sciences of the United States of America*, 118(24), Article e2101666118.
- Vickery, J. A., Mallord, J. W., Adams, W. M., Beresford, A. E., Both, C., Cresswell, W., Diop, N., Ewing, S. R., Gregory, R. D., Morrison, C. A., Sanderson, F. J., Thorup, K., Van Wijik, R. E., & Hewson, C. M. (2023). The conservation of Afro-Palaearctic migrants: What we are learning and what we need to know? *Ibis*, 165(3), 717–738.
- Weisshaupt, N., Lehtiniemi, T., & Koistinen, J. (2021). Combining citizen science and weather radar data to study large-scale bird movements. *Ibis*, 163, 728–736.
- Wikelski, M., Kays, R. W., Kasdin, N. J., Thorup, K., Smith, J. A., & Swenson, G. W., Jr. (2007). Going wild: What a global small-animal tracking system could do for experimental biologists. *Journal of Experimental Biology*, 210(2), 181–186.
- Wilcove, D. S., & Wikelski, M. (2008). Going, going, gone: Is animal migration disappearing. *PLoS Biology*, 6(7), Article e188.
- Wilcox, A. A., Newman, A. E., Raine, N. E., Mitchell, G. W., & Norris, D. R. (2021). Effects of early-life exposure to sublethal levels of a common neonicotinoid insecticide on the orientation and migration of monarch butterflies (*Danaus plexippus*). *Journal of Experimental Biology*, 224(4), Article jeb230870.
- Wild, T. A., van Schalkwyk, L., Viljoen, P., Heine, G., Richter, N., Vorneweg, B., Koblit, J. C., Dechmann, D. K., Rogers, W., Partecke, J., & Linek, N. (2023). A multi-species evaluation of digital wildlife monitoring using the Sigfox IoT network. *Animal Biotelemetry*, 11(1), Article 13.
- Willmott, J. R., Forcey, G., & Vukovich, M. (2023). New insights into the influence of turbines on the behaviour of migrant birds: Implications for predicting impacts of offshore wind developments on wildlife. *Journal of Physics: Conference Series*, 2507(1), Article 012006.
- Woodworth, B. K., Mitchell, G. W., Norris, D. R., Francis, C. M., & Taylor, P. D. (2015). Patterns and correlates of songbird movements at an ecological barrier during autumn migration assessed using landscape- and regional-scale automated radiotelemetry. *Ibis*, 157(2), 326–339.
- Zaugg, S., Saporta, G., van Loon, E., Schmaljohann, H., & Liechti, F. (2008). Automatic identification of bird targets with radar via patterns produced by wing flapping. *Journal of the Royal Society Interface*, 5, 1041–1053.
- Zenzal, T. J., Ward, M. P., Diehl, R. H., Buler, J. J., Smolinsky, J., Deppe, J. L., Bolus, R. T., Celis-Murillo, A., & Moore, F. R. (2021). Retreat, detour or advance? Understanding the movements of birds confronting the Gulf of Mexico. *Oikos*, 130(5), 739–752.
- Zurell, D., Graham, C. H., Gallien, L., Thuiller, W., & Zimmermann, N. E. (2018). Long-distance migratory birds threatened by multiple independent risks from global change. *Nature Climate Change*, 8(11), 992–996.
- Züst, Z., Mukhin, A., Taylor, P. D., & Schmaljohann, H. (2023). Pre-migratory flights in migrant songbirds: The ecological and evolutionary importance of understudied exploratory movements. *Movement Ecology*, 11(1), Article 78.

How to cite this article: Mitchell, L., Brust, V., Karwinkel, T., Åkesson, S., Kishkinev, D., Norevik, G., Szep, T., Hedenström, A., Lagerveld, S., Helm, B., & Schmaljohann, H. (2025). Conservation-focused mapping of avian migratory routes using a pan-European automated telemetry network. *Conservation Biology*, e70017. <https://doi.org/10.1111/cobi.70017>