

Hirshfeld atoms in molecules revisited

*P. Bultinck**

Department of Inorganic and Physical Chemistry, Ghent University, Krijgslaan 281 (S3), 9000
Gent, Belgium.

e-mail: Patrick.Bultinck@UGent.be.

The well-known Hirshfeld [1] method for the atom in the molecule has been criticized by several authors [2-6] as being a quite arbitrary method to describe the atom in the molecule (AIM). Bultinck et al. [5-6] recently described a self-consistent Hirshfeld scheme where this arbitrariness was removed by letting the molecule decide its own promolecule. The weight function for the AIM A is given by:

$$w_A(\mathbf{r}, N_A^0) = \frac{\rho_A^0(\mathbf{r}, N_A^0)}{\sum_A \rho_A^0(\mathbf{r}, N_A^0)}$$

Where explicitly the dependence on the number of electrons N_A in AIM A is indicated. The AIM density $\rho_A(\mathbf{r})$ is then obtained by letting the weight act on the molecular density $\rho(\mathbf{r})$:

$$\rho_A(\mathbf{r}) = w_A(\mathbf{r}) \rho(\mathbf{r}) \quad \wedge \quad N_A = \int \rho_A(\mathbf{r}) d\mathbf{r}$$

The key feature of the Hirshfeld-I method is that the Hirshfeld process is repeated until $N_A^0 = N_A$. In this presentation the information theoretic context of Hirshfeld-I is sketched, performance of the method compared to regular Hirshfeld analysis discussed [5-7], its impact on the nature of atom condensed Fukui functions described [8] and its use for good quality monopole approximation electrostatic potentials shown [9].

[1] F.L. Hirshfeld, *Theor. Chim. Acta*, **44**, 129 (1977).

[2] E.R. Davidson, and S. Chakravorty, *Theor. Chim. Acta*, **83**, 319 (1992).

[3] R.F.W. Bader, and C.F. Matta *J. Phys. Chem. A*, **108**, 8385 (2004).

[4] C.F. Matta, and R.F.W. Bader, *J. Phys. Chem. A*, **110**, 6365 (2006).

[5] P. Bultinck, C. Van Alsenoy, P.W. Ayers, and R. J. Carbó-Dorca, *J. Chem. Phys.*, **126**, 144111 (2007).

[6] P. Bultinck, *P. Faraday Discussions*, **135**, 244 (2007).

[7] P. Bultinck, P.W. Ayers, S. Fias, K. Tiels, C. Van Alsenoy, *Chem. Phys. Lett.*, **444**, 205 (2007).

[8] P. Bultinck, P., S. Fias, C. Van Alsenoy, P.W. Ayers, and R. Carbó-Dorca, *J. Chem. Phys.*, **127**, 034102 (2007).

[9] S. Van Damme, P. Bultinck, and S. Fias, *J. Chem. Theory and Comput.*, **5**, 334 (2009).