

1 A time-integrated MODIS burn severity assessment using the multi-temporal differenced
2 Normalized Burn Ratio (dNBR_{MT})

3 S. VERAVERBEKE^{A*}, S. LHERMITTE^B, W.W. VERSTRAETEN^C AND R. GOOSSENS^A

4 A Department of Geography, Ghent University, Krijgslaan 281 S8, BE-9000 Ghent, Belgium

5 B Centro de Estudios Avanzados en Zonas Aridas, Universidad de la Serena, Campus A. Bello,
6 ULS, Chile

7 C Department of Biosystems, Katholieke Universiteit Leuven, Willem de Croylaan 34, BE-3001,
8 Belgium

9 *Corresponding author. Email: sander.veraverbeke@ugent.be (Tel. 0032 9 2644646, Fax. 0032 9
10 264 49 85)

11 **Abstract**

12 Burn severity is an important parameter in post-fire management. It incorporates both the direct
13 fire impact (vegetation depletion) and ecosystem responses (vegetation regeneration). From a
14 remote sensing perspective, burn severity is traditionally estimated using Landsat's differenced
15 Normalized Burn Ratio (dNBR). In this case study of the large 2007 Peloponnese (Greece)
16 wildfires, Landsat dNBR estimates correlated reasonably well with Geo Composite Burn Index
17 (GeoCBI) field data of severity ($R^2 = 0.56$). The usage of Landsat imagery is, however, restricted
18 by cloud cover and image-to-image normalization constraints. Therefore a multi-temporal burn
19 severity approach based on coarse spatial, high temporal resolution Moderate Resolution Imaging

20 Spectroradiometer (MODIS) imagery is presented in this study. The multi-temporal dNBR
21 (dNBR_{MT}) is defined as the one-year integrated difference between burned pixels and their
22 unique control pixels. These control pixels were selected based on time series similarity and
23 spatial context and reflect how burned pixels would have behaved in the case no fire had
24 occurred. Linear regression between downsampled Landsat dNBR and dNBR_{MT} estimates
25 resulted in a moderate-high coefficient of determination $R^2 = 0.54$. dNBR_{MT} estimates are
26 indicative for the change in vegetation productivity due to the fire. This change is considerably
27 higher for forests than for more sparsely vegetated areas like shrub lands. Although Landsat
28 dNBR is superior for spatial detail, MODIS-derived dNBR_{MT} estimates present a valuable
29 alternative for burn severity mapping at continental to global scale without image availability
30 constraints. This is beneficial to compare trends in burn severity across regions and time.
31 Moreover, thanks to MODIS's repeated temporal sampling, the dNBR_{MT} accounts for both first-
32 and second-order fire effects.

33 **Keywords:** differenced Normalized Burn Ratio, fire severity, burn severity, MODIS, Landsat
34 Thematic Mapper, Composite Burn Index, multi-temporal, vegetation regeneration

35 **1 Introduction**

36 Biomass burning is a major disturbance in almost all terrestrial ecosystems (Pausas, 2004; Riano
37 et al., 2007). At landscape level, wildland fires partially or completely remove the vegetation
38 layer and affect post-fire vegetation composition (Epting and Verbyla 2005). The fire-induced
39 vegetation depletion causes abrupt changes in carbon, energy and water fluxes at local scale
40 (Amiro et al., 2006a; Montes-Helu et al., 2009), thereby influencing species richness, habitats and
41 community composition (Moretti et al., 2002; Capitaino and Carcaillet, 2008). Accurate estimates

42 of post-fire effects are therefore of paramount importance. To name these post-fire effects the
43 terms fire severity and burn severity are often interchangeably used (Keeley, 2009) describing the
44 amount of damage (Chafer, 2008), the physical, chemical and biological changes (Lee et al.,
45 2008) or the degree of alteration (Eidenshink et al., 2007) that fire causes to an ecosystem. Some
46 authors, however, suggest a clear distinction between both terms by considering the fire
47 disturbance continuum (Jain et al., 2004), which addresses three different temporal fire effects
48 phases: before, during and after the fire. In this context, fire severity quantifies the short-term fire
49 effects in the immediate post-fire environment whereas burn severity quantifies both the short-
50 and long-term impact as it includes response processes (e.g. resprouting, delayed mortality)
51 (Lentile et al., 2006; Key, 2006). Figure 1 represents a summary of post-fire effects terminology.

52 FIGURE 1 HERE

53 In remote sensing studies burn severity is traditionally estimated using Landsat imagery (Key and
54 Benson, 2005; French et al., 2008). A popular approach, partly because of its conceptual
55 simplicity, can be found in ratioing band reflectance data. In this respect the Normalized Burn
56 Ratio (NBR) has become accepted as the standard spectral index to assess burn severity (Lopez-
57 Garcia and Caselles, 1991; Key and Benson, 2005; French et al., 2008, Veraverbeke et al.,
58 2010a). The NBR relates to vegetation moisture content by combining the near infrared (NIR)
59 and mid infrared (MIR) spectral regions. Generally, pre- and post-fire NBR images are bi-
60 temporally differenced, resulting in the differenced NBR (dNBR).

61 The dNBR method relies on Landsat imagery and thus depends on image availability, which is
62 limited to infrequent images over small areas due to Landsat's 16-day revisiting cycle and cloud
63 cover (Ju and Roy, 2008). Bi-temporal studies are even more hampered as they require an
64 effective image-to-image normalization (Coppin et al. 2004) including the removal of
65 phenological, atmospheric and bi-directional reflectance distribution function (BRDF) effects

66 (Verbyla et al., 2008; Veraverbeke et al., 2010b). As a result Landsat-based burn severity studies
67 have proven to be valuable for obtaining detailed information over specific fires, however, the
68 magnitude of the observed dNBR change heavily depends on assessment timing (Key, 2006;
69 Veraverbeke et al., 2010c). This temporal dissimilarity limits the comparison between bi-
70 temporal dNBR assessments of different fires (Eidenshink et al., 2007, Verbyla et al., 2008),
71 especially when a comparison between different ecoregions is required (Eidenshink et al., 2007,
72 French et al., 2008). The use of high temporal, coarse spatial resolution data possibly provides a
73 sound alternative to Landsat dNBR estimates. In addition, their repeated temporal sampling
74 allows quantifying both the direct fire impact and regeneration processes. To date few studies
75 have implemented coarse resolution time series to assess burn severity. In this context it is worth
76 mentioning the effort of Lhermitte et al. (2010a), who illustrated the potential of time series data
77 to account for inter- and intra-annual post-fire vegetation dynamics. In their method each burned
78 pixel is compared with an unburned control pixel. These control pixels were selected based on
79 pre-fire time series similarity and spatial context.

80 The aim of this study is to present a multi-temporal dNBR ($dNBR_{MT}$) burn severity assessment as
81 an alternative for traditional Landsat dNBR mapping. The method incorporates both the direct
82 fire impact and vegetation regeneration (Lentile et al., 2006). Moderate Resolution Imaging
83 Spectroradiometer (MODIS) time series are used over the large 2007 Peloponnese (Greece)
84 wildfires. $dNBR_{MT}$ estimates are compared with Landsat and field data.

85 **2 Data and study area**

86 **2.1 Study area**

87 The study area is situated at the Peloponnese peninsula, in southern Greece ($36^{\circ}30'$ - $38^{\circ}30'$ N,
88 21° - 23° E) (see figure 2). The topography is rugged with elevations ranging between 0 and 2404

89 m above sea level. The climate is typically Mediterranean with hot, dry summers and mild, wet
90 winters. For the Kalamata meteorological station (37°4' N, 22°1' E) the average annual
91 temperature is 17.8 °C and the mean annual precipitation equals 780 mm.

92 FIGURE 2 HERE

93 After a severe drought period several large wildfires of unknown cause have struck the area in the
94 2007 summer. The fires were the worst natural disaster of the last decades in Greece, both in
95 terms of human losses and the extent of the burned area. The fires consumed more than 175 000
96 ha, which consisted of 57% shrub land, 21% coniferous forest, 20% olive groves and 2%
97 broadleaved forest (Veraverbeke et al., 2010c).

98 **2.2 Field data**

99 150 Geo Composite Burn Index (GeoCBI) plots were sampled one year post-fire, in September
100 2008. The GeoCBI is a modification of the Composite Burn Index (CBI) (De Santis and
101 Chuvieco, 2009). It is an operational tool used in conjunction with the Landsat dNBR approach to
102 assess burn severity in the field (Key and Benson, 2005). The GeoCBI divides the ecosystem into
103 five different strata, one for the substrates and four vegetation layers. These strata are: (i)
104 substrates, (ii) herbs, low shrubs and trees less than 1 m, (iii) tall shrubs and trees of 1 to 5 m, (iv)
105 intermediate trees of 5 to 20 m and (v) big trees higher than 20 m. In the field form, 20 different
106 factors can be rated (e.g. soil and rock cover/color change, % LAI change, char height) but only
107 those factors present and reliably rateable, are considered. The rates are given on a continuous
108 scale between zero and three and the resulting factor ratings are averaged per stratum. Based on
109 these stratum averages, the GeoCBI is calculated in proportion to their corresponding fraction of
110 cover, resulting in a weighted average between zero and three that expresses burn severity. As the

111 field data were collected one year post-fire, it is an extended assessment. Additional information
112 on the field data can be found in Veraverbeke et al. (2010c).

113 **2.3 Landsat data**

114 For the traditional Landsat dNBR assessment two anniversary date Thematic Mapper (TM)
115 images (path/row 184/34) were used (23/07/2006 and 13/08/2008). In correspondence with the
116 timing of the field sampling, the post-fire image was acquired one year post-fire. The images
117 were acquired in the summer, minimizing effects of vegetation phenology and differing solar
118 zenith angles. The images were subjected to geometric, radiometric, atmospheric and topographic
119 correction.

120 The 2008 image was geometrically corrected using 34 ground control points (GCPs), recorded in
121 the field with a Garmin eTrex Vista GPS (15 m error in x and y (Garmin, 2005)). The resulting
122 Root Mean Squared Error (RMSE) was lower than 0.5 pixels. The 2006 and 2008 images were
123 co-registered within 0.5 pixels accuracy. The images were registered in UTM (zone 34S), with
124 the World Geodetic System 84 (WGS-84) as geodetic datum.

125 Raw digital numbers (DNs) were scaled to at-sensor radiance values (L_s) (Chander et al., 2007)
126 The radiance to reflectance conversion was performed using the COST method (Chavez, 1996):

$$127 \quad \rho_a = \frac{\pi(L_s - L_d)}{(E_o / d^2)(\cos \theta_z)^2} \quad (\text{Eq. 1})$$

128 where ρ_a is the atmospherically corrected reflectance at the surface; L_s is the at-sensor radiance
129 ($\text{Wm}^{-2}\text{sr}^{-1}$); L_d is the path radiance ($\text{Wm}^{-2}\text{sr}^{-1}$); E_o is the solar spectral irradiance (Wm^{-2}); d is the
130 earth-sun distance (astronomical units); and θ_z is the solar zenith angle. The COST method is a
131 dark object subtraction (DOS) approach that assumes 1% surface reflectance for dark objects
132 (e.g. deep water). After applying the COST atmospheric correction, pseudo-invariant features

133 (PIFs) such as deep water and bare soil pixels, were examined in the images. No further relative
 134 normalization between the images was required.

135 It was necessary to correct for different illumination effects due to topography as the common
 136 assumption that shading effects are removed in ratio-based analyses does not necessarily hold
 137 true (Verbyla et al., 2008; Veraverbeke et al., 2010b). This was done based on the modified C
 138 correction method (Veraverbeke et al., 2010b), a modification of the original C correction
 139 approach (Teillet et al., 1982), using a DEM and knowledge of the solar zenith and azimuth angle
 140 at the moment of image acquisition. Topographical slope and aspect data were derived from 90 m
 141 Shuttle Radar Topographic Mission SRTM elevation data (Jarvis et al., 2006) resampled and
 142 coregistered with the Landsat images. The illumination is modeled as:

$$143 \quad \cos \gamma_i = \cos \theta_p \cos \theta_z + \sin \theta_p \sin \theta_z \cos(\phi_a - \phi_o) \quad (\text{Eq. 2})$$

144 where γ_i is the incident angle (angle between the normal to the ground and the sun rays); θ_p is
 145 the slope angle; θ_z is the solar zenith angle; ϕ_a is the solar azimuth angle; and ϕ_o is the aspect
 146 angle. Then terrain corrected reflectance ρ_t is defined as:

$$147 \quad \rho_t = \rho_a \left(\frac{1 + c_k}{\cos \gamma_i + c_k} \right) \quad (\text{Eq. 3})$$

148 where c_k is a band specific parameter $c_k = b_k / m_k$ where b_k and m_k are the respective intercept
 149 and slope of the regression equation $\rho_a = b_k + m_k \cos \gamma_i$.

150 Finally, by inputting the NIR (TM4: centered at 830 nm) and MIR (TM7: centered at 2215 nm)
 151 bands NBR and dNBR images were generated:

$$152 \quad NBR = \frac{NIR - MIR}{NIR + MIR} \quad dNBR = NBR_{pre} - NBR_{post} \quad (\text{Eq. 4})$$

153 2.4 MODIS data

154 Level 2 daily Terra MODIS surface reflectance (500 m) tiles (MOD09GA) including associated
155 Quality Assurance (QA) layers were acquired from the National Aeronautics and Space
156 Administration (NASA) Warehouse Inventory Search Tool (WIST) (<https://wist.echo.nasa.gov>)
157 for the period 01/01/2006 till 31/12/2008. These products contain an estimate of the surface
158 reflectance for seven optical bands as it would have been measured at ground level as if there
159 were no atmospheric scattering or absorption (Vermote et al., 2002). The data preprocessing steps
160 included subsetting, reprojecting, compositing, creating continuous time series and indexing. The
161 study area was clipped and the NIR (centered at 858 nm), MIR (centered at 2130 nm) and QA
162 layers were reprojected into UTM with WGS 84 as geodetic datum. Subsequently, the daily NIR,
163 MIR and QA data were converted in 8-day composites using the minimum NIR criterion to
164 minimize cloud contamination and off-nadir viewing effects (Holben, 1986). The minimum NIR
165 criterion has proven to allow a more accurate discrimination between burned and unburned pixels
166 than traditional Maximum Value Composites (MVCs) (Chuvieco et al., 2005). After compositing
167 bad QA observations were replaced by a Savitzky-Golay filter as implemented in the TIMESAT
168 software (Jonsson and Eklundh, 2004). The TIMESAT program allows the inclusion of a
169 preprocessing mask that determines the uncertainty of data values. Cloud-affected observations
170 were identified using the internal cloud and cloud-adjacency algorithm flags of the QA layer.
171 These flags consist of binary layers which permit to assign a zero weight value to cloudy and
172 cloud-adjacent observations. Consequently, these data do not influence the filter procedure. Only
173 the values of the masked observations were replaced to retain as much as possible the original
174 NIR and MIR reflectance values. Finally, the NBR index was calculated as using equation 4.

175 **2.5 Control pixel data**

176 Control pixel data were retrieved making use of pre-fire time series similarity and spatial context
177 (Lhermitte et al., 2010b) as implemented in Veraverbeke et al. (2010c). The control pixel
178 selection procedure assigns a unique control pixel to each burned pixel. This is done based on
179 time series similarity between a burned pixel and its closest unburned neighbor pixels during a
180 pre-fire period. To quantify dissimilarity the averaged Euclidian distance dissimilarity criterion D
181 was used:

$$182 \quad D = \frac{\sqrt{\sum_{t=1}^N (NBR_t^f - NBR_t^x)^2}}{N} \quad (2)$$

183 where NBR_t^f and NBR_t^x are the respective burned focal and unburned candidate control pixel
184 time series, while N is the number of observations in pre-fire year ($N=46$). The Euclidian distance
185 metric has an intuitive appeal: it quantifies the straight line inter-point distance in a multi-
186 temporal space as distance measure. As a result, it is robust for both data space translations and
187 rotations. Consequently, it is a very useful metric to assess inter-pixel differences in time series
188 (Lhermitte et al., 2010b). In this approach the averaged time series from the four most similar out
189 of eight candidate pixels defines the control pixel time series. This setting accounts for both a
190 beneficial averaging effect and the advantage of spatial proximity (Veraverbeke et al. 2010c).
191 The resulting control pixels reflect the vegetation dynamics of each burned pixel in case that
192 there would not have occurred a fire. Additional information on the control plot selection
193 procedure can be found in Lhermitte et al. (2010b) and Veraverbeke et al. (2010c).

194 **3 Methodology**

195 Burn severity incorporates both short-and long-term post-fire effects on the environment (Lentile
196 et al., 2006). Consequently, burn severity is a combination of immediate fire impact and the

197 ecosystem's ability to regenerate. Based on these characteristics, we propose a multi-temporal
198 dNBR ($dNBR_{MT}$) that integrates the difference between the NBR values of a burned pixel and its
199 corresponding control pixel over time. Doing so the $dNBR_{MT}$ is defined as:

$$200 \quad dNBR_{MT} = \frac{\sum_{t=1}^N (NBR_t^f - NBR_t^c)}{N} \quad (3)$$

201 where NBR_t^f and NBR_t^c are the respective burned focal and unburned control pixel observations,
202 while N is the number of post-fire observations included in the study (here $N=46$ for one year)
203 and $t=1$ is the first post-fire observation. Figure 2 illustrates the principle of the $dNBR_{MT}$.
204 Dividing by the number of post-fire observations N normalizes the $dNBR_{MT}$ data to the same
205 range as bi-temporal dNBR assessments. $dNBR_{MT}$ estimates will show large positive values for
206 high burn severity. The application of an integral has been used to characterize vegetation
207 productivity (Reed et al., 1994; Heumann et al., 2007). The integrated change between NBR
208 values of control and burned pixels is therefore indicative for the change in vegetation
209 productivity caused by the fire. To evaluate the performance of the multi-temporal approach
210 comparison is made with a traditional Landsat TM dNBR assessment and GeoCBI field data.

211 FIGURE 3 HERE

212 **4 Results**

213 Figure 4A shows the result of the MODIS $dNBR_{MT}$ approach, while figure 4B details a specific
214 burned area framed in blue in figure 4A. Figure 4C displays the traditional Landsat dNBR, while
215 figure 4D also depicts the detailed subset. On a coarse scale the MODIS and Landsat assessments
216 reveal the same patterns of burn severity, however, it is trivial that Landsat estimates are

217 characterized by more spatial detail. This is also visible in figure 5. The scatter plot between
218 GeoCBI and Landsat dNBR estimates is given in figure 5A. The linear regression fit resulted in a
219 coefficient of determination $R^2 = 0.56$. Figure 5B presents the scatter plot between downsampled
220 Landsat data and corresponding $dNBR_{MT}$ estimates for the 150 field-sampled locations. The
221 vertical bars indicate the standard deviation (sd) of the Landsat pixels within one MODIS pixel.
222 Although the correlation between downsampled Landsat dNBR and MODIS $dNBR_{MT}$ estimates
223 is moderately high ($R^2 = 0.54$), it is clear that there exists considerable variation within one
224 MODIS pixel (sd of Landsat dNBR up to 0.25).

225 FIGURE 4 HERE

226 FIGURE 5 HERE

227 In figure 6 mean $dNBR_{MT}$ (sd) is plotted per land cover type. One can clearly see that the one-
228 year integrated change is higher for forests than for more sparsely vegetated covers. $dNBR_{MT}$
229 estimates are the highest for coniferous forest, followed by broadleaved forest. Shrub land and
230 olive groves have considerably lower $dNBR_{MT}$ estimates. Figure 7 examples temporal profiles of
231 eight pixels. These figures demonstrate that $dNBR_{MT}$ estimates account for both the direct fire
232 impact and the ability to recover.

233 FIGURE 6 HERE

234 FIGURE 7 HERE

235 **5 Discussion**

236 A major advantage of the multi-temporal burn severity approach is its combination of both the
237 immediate fire impact and vegetation regrowth. As such, it is more tightly connected to the

238 definition of burn severity. Key and Benson (2005) stated that burn severity encloses both first
239 and second order fire effects. The most important first order effect is the fire's vegetation
240 consumption, while vegetation regeneration and delayed mortality are substantial second order
241 effects. In that respect, Lentile et al. (2006) specified that burn severity relates to the amount of
242 time necessary to return to pre-fire level. As a consequence plots that experienced a high fire
243 severity and fast regeneration will result in similar $dNBR_{MT}$ outcomes as plots that were only
244 slightly affected by the fire but with slow recovery. While in some studies it can be important to
245 distinguish between first- and second-order effects, burn severity incorporates both (Lentile et al.,
246 2006; Keeley, 2009). The application of an integral has been used to characterize vegetation
247 productivity (Reed et al., 1994; Heumann et al., 2007). As such, the integrated change between
248 NBR values of control and burned pixels, as gauged by the $dNBR_{MT}$, reflects the change in
249 productivity due to the fire. Seasonality and recovery processes vary per land cover type (Reed et
250 al., 1994; White et al., 1996). As a result, $dNBR_{MT}$ estimates are clearly higher for forests than
251 for more sparsely vegetated areas (figures 6 and 7). Recovery in forests can take several decades
252 (Nepstad et al., 1999), whereas shrub species are typified by a relatively fast recovery (Keeley et
253 al., 2005). The $dNBR_{MT}$ incorporates this difference. Moreover, depending on the application and
254 the ecotype, one could decide to alter the integration period (one year in this study).

255 In corroboration with previous findings (French et al., 2008), Landsat $dNBR$ correlated
256 reasonably well with field data of severity. The correlation between GeoCBI and Landsat data
257 differed from previously published outcomes based on the same data (Veraverbeke et al. 2010a),
258 mainly because of some minor changes in satellite preprocessing and the exclusion of ten
259 unburned field plots. Multi-temporal MODIS burn severity estimates showed a moderate-high
260 correlation with the $dNBR$ of a traditional bi-temporal Landsat assessment ($R^2 = 0.54$). The slope

261 of the regression equation (0.77) was considerably lower than one. In contrast with the one-year
262 post-fire Landsat assessment, $dNBR_{MT}$ estimates also incorporate observations from the
263 immediate post-fire period. As a consequence $dNBR_{MT}$ estimates were slightly higher than the
264 Landsat $dNBR$. Despite of the coarse scale resemblance between Landsat and MODIS data,
265 Landsat data are superior to reveal spatial detail (Hilker et al., 2009). These data, however, fail to
266 comprehend the temporal dimension of burn severity. Moreover, the magnitude of change
267 measured with the traditional Landsat $dNBR$ highly depends on assessment timing (Key, 2006;
268 Veraverbeke et al., 2010c). Allen and Sorbel (2008), for example, found that initial and extended
269 assessments produced significantly different information with regards to burn severity for tundra
270 vegetation, while the timing of the assessment had no effect for back spruce forest, which was
271 attributed to the rapid tundra recovery. Verbyla et al. (2008) reported a seasonality effect that
272 resulted in large dissimilarities in $dNBR$ values for only slightly differing assessment timings,
273 probably due to a combined effect of senescing vegetation and changing illumination conditions.
274 Veraverbeke et al. (2010b) illustrated the necessity to correct for illumination effects, also in a
275 ratio-based NBR analysis, because these effects affected the performance of the $dNBR$, even for
276 bi-temporal acquisitions schemes that only slightly deviated from the ideal anniversary date
277 scheme. This timing constraint potentially hampers the comparison of Landsat $dNBR$ estimates
278 across region and time (Eidenshink et al., 2007; Verbyla et al., 2008). If the period of the
279 $dNBR_{MT}$'s integration remains the same for different fires, the multi-temporal approach truly has
280 the potential to allow a better comparison of burn severity either in time or space. Thus, where
281 fine resolution Landsat studies allow revealing high spatial detail, which is favorable for regional
282 studies, their usage is limited due cloud cover problems (Ju and Roy, 2008) and difficulties in
283 image-to-image normalization (Coppin et al., 2004; Verbyla et al., 2008; Veraverbeke et al.,
284 2010b). Therefore, the high temporal frequency of coarse resolution imagery can either be a vital

285 complement to traditional Landsat dNBR mapping of specific fires or an imperative alternative
286 for the assessment of burn severity at continental to global scales.

287 **6 Conclusions**

288 In this study a multi-temporal method to assess burn severity of the 2007 Peloponnese (Greece)
289 wildfires has been proposed. The approach introduces an alternative for traditional Landsat
290 dNBR mapping, which can be constrained due to cloud cover and image-to-image normalization
291 difficulties. The method is based on coarse spatial resolution with high temporal frequency
292 MODIS imagery. MODIS's daily MIR and NIR reflectance products were first composited in 8-
293 day periods and missing values were replaced. Subsequently, for each burned pixel a unique
294 control pixel has been retrieved based on time series similarity and spatial context. The $dNBR_{MT}$
295 was then calculated as the one-year post-fire integrated difference between the NBR of the
296 control and burned pixels, averaged by the total number of observations. $dNBR_{MT}$ estimates
297 reflect the change in vegetation productivity caused by the fire. This change is clearly higher for
298 forests than for shrub lands. By integrating over time, $dNBR_{MT}$ estimates account for both the
299 direct fire impact and ecosystem responses. As such the $dNBR_{MT}$ is more tightly connected to the
300 definition of burn severity compared to traditional bi-temporal Landsat dNBR mapping. $dNBR_{MT}$
301 estimates correlated reasonably well with the downsampled Landsat dNBR, which on its turn
302 showed a moderate-high correlation with GeoCBI field data. Although Landsat dNBR is superior
303 for spatial detail in regional scale studies, the $dNBR_{MT}$ presents a valuable alternative for burn
304 severity mapping at a regional to global scale. The approach also has potential to enhance
305 comparability of different fires across regions and time.

306 **Acknowledgements**

307 The study was financed by the Ghent University special research funds (BOF: Bijzonder
308 Onderzoeksfonds). The authors would like to thank the anonymous reviewers for their
309 constructive remarks.

310 **References**

311 Allen, J., Sorbel, B., 2008. Assessing the differenced Normalized Burn Ratio's ability to map
312 burn severity in the boreal forest and tundra ecosystems of Alaska's national parks *Int. J.*
313 *Wildland Fire*, 17, 463-475

314 Amiro, B., Barr, A., Black, T., Iwashita, H., Kljun, N., McCaughey, J., Morgenstern, K.,
315 Muruyama, S., Nesic, Z., Orchansky, A., Saigusa, N., 2006. Carbon, energy and water fluxes at
316 mature and disturbed forest sites, Saskatchewan, Canada. *Agric. and For. Meteorol.*, 136, 237-
317 251

318 Capitaino, R., Carcaillet, C., 2008. Post-fire Mediterranean vegetation dynamics and diversity: a
319 discussion of succession models. *For. Ecol. Manage.*, 255, 431-439

320 Chafer, C., 2008. A comparison of fire severity measures: an Australian example and
321 implications for predicting major areas of soil erosion. *Catena*, 74, 235-245

322 Chander G., Markham, L., Barsi, J., 2007. Revised Landsat-5 Thematic Mapper radiometric
323 calibration. *IEEE Geosc. Remote Sens. Lett.*, 4, 490-494

324 Chavez, P., 1996. Image-based atmospheric corrections – revisited and improved. *Photogramm.*
325 *Eng. Remote Sens.*, 6, 1025-1036

326 Chuvieco, E., Ventura, G., Martin, P., Gomez, I., 2005. Assessment of multitemporal
327 compositing techniques of MODIS and AVHRR images for burned land mapping. *Remote Sens.*
328 *Environ.*, 94, 450-462

329 Coppin, P., Jonckheere, I., Nackaerts, K., Muys, B., 2004. Digital change detection techniques in
330 ecosystem monitoring: a review. *Int. J. Remote Sens.*, 25, 1565-1595

331 De Santis, A., Chuvieco, E., 2009. GeoCBI: a modified version of the Composite Burn Index for
332 the initial assessment of the short-term burn severity from remotely sensed data. *Remote Sens.*
333 *Environ.*, 113, 554-562

334 Eidenshink, J., Schwind, B., Brewer, K., Zhu, Z., Quayle, B., Howard, S., 2007. A project for
335 monitoring trends in burn severity. *Fire Ecol.*, 3, 3-21

336 Epting, J., Verbyla, D., 2005. Landscape-level interactions of prefire vegetation, burn severity,
337 and postfire vegetation over a 16-year period in interior Alaska. *Can. J. For. Res.*, 35, 1367-1377

338 French, N., Kasischke, E., Hall, R., Murphy, K., Verbyla, D., Hoy, E., Allen, J., 2008. Using
339 Landsat data to assess fire and burn severity in the North American boreal forest region: an
340 overview and summary of results. *Int. J. Wildland Fire*, 17, 443-462

341 Garmin, 2005. Garmin eTrex Vista personal navigator. Owner's manual and reference guide.
342 Available from: [https://buy.garmin.com/shop/store/manual.jsp?product=010-00243-](https://buy.garmin.com/shop/store/manual.jsp?product=010-00243-00&cID=167&pID=163)
343 [00&cID=167&pID=163](https://buy.garmin.com/shop/store/manual.jsp?product=010-00243-00&cID=167&pID=163) (Last visited on 22/04/2010).

344 Heumann, B., Seaquist, J., Eklundh, L., Jonsson, P., 2007. AVHRR derived phenological change
345 in the Sahel and Soudan, Africa, 1982-2005. *Remote Sens. Environ.*, 108, 385-392

346 Hilker, T., Wulder, M., Coops, N., Linke, J., McDermid, G., Masek, J., Gao, F., White, J. 2009.
347 A new data fusion model for high spatial- and temporal-resolution mapping of forest disturbance
348 based on Landsat and MODIS. *Remote Sens. Environ.*, 113, 1623-1627

349 Holben, B., 1986. Characteristics of maximum-value composite images from temporal AVHRR
350 data. *Int. J. Remote Sens.*, 7, 1417-1434

351 Jain, T., Pilliod, D., Graham, R., 2004. Tongue-tied. *Wildfire*, 4, 22–26.

352 Jarvis, A., Reuter, H., Nelson, A., Guevara, E., 2006. Hole-filled seamless SRTM data V3.
353 Available from: <http://srtm.csi.cgiar.org> (Last visited on 22/04/2010)

354 Jonsson, P., Eklundh, L., 2004. TIMESAT-a program for analyzing time-series of satellite sensor
355 data. *Comput. Geosci.*, 30, 833-845

356 Ju, J., Roy, D., 2008. The availability of cloud-free Landsat ETM+ data over the conterminous
357 United States and globally. *Remote Sens. Environ.*, 112, 1196-1211

358 Keeley, J., 2009. Fire intensity, fire severity and burn severity: a brief review and suggested
359 usage. *Int. J. Wildland Fire*, 18, 116-126

360 Key, C., Benson, N., 2005. Landscape assessment: ground measure of severity; the Composite
361 Burn Index, and remote sensing of severity, the Normalized Burn Index, in: Lutes, D., Keane, R.,
362 Caratti, J., Key, C., Benson, N., Sutherland, S., Gangi, L. (Eds.), FIREMON: Fire effects
363 monitoring and inventory system, USDA Forest Service, Rocky Mountains Research Station,
364 General Technical Report RMRS-GTR-164-CD LA, pp. 1-51

365 Keeley, J., Fotheringham, C., Bear-Keeley, M. 2005. Factors affecting plant diversity during
366 post-fire recovery and succession of mediterranean-climate shrublands in California, USA.
367 *Divers Distrib*, 11, 525-537

368 Key, C., 2006. Ecological and sampling constraints on defining landscape fire severity. *Fire*
369 *Ecol.*, 2, 34–59

370 Lee, B., Kim, S., Chung, J., Park, P., 2008. Estimation of fire severity by use of Landsat TM
371 images and its relevance to vegetation and topography in the 2000 Samcheok forest fire. *J. For.*
372 *Res.*, 13, 197-204

373 Lentile, L., Holden, Z., Smith, A., Falkowski, M., Hudak, A., Morgan, P., Lewis, S., Gessler, P.,
374 Benson, N., 2006. Remote sensing techniques to assess active fire characteristics and post-fire
375 effects. *Int. J. Wildland Fire*, 15, 319-345

376 Lhermitte, S., Verbesselt, J., Verstraeten, W.W., Veraverbeke, S. Coppin, P., 2010a. Assessing
377 intra-annual vegetation regrowth after fire using the pixel based regeneration index. *ISPRS J.*
378 *Photogramm. Remote Sens.*, in review

379 Lhermitte, S., Verbesselt, J., Verstraeten, W.W., Coppin, P., 2010b. A pixel based regeneration
380 index using time series similarity and spatial context. *Photogramm. Eng. Remote Sens.*, 76, 673-
381 682

382 Lopez-Garcia, M., Caselles, V., 1991. Mapping burns and natural reforestation using Thematic
383 Mapper data. *Geocarto Int.*, 6, 31-37

384 Michalek, J., French, N., Kasischke, E., Johnson, R., Colwell, J., 2000. Using Landsat TM data to
385 estimate carbon release from burned biomass in an Alaskan spruce forest complex. *Int. J. Remote*
386 *Sens.*, 21, 323-338

387 Nepstad, D., Verssimo, A., Alencar, A., Nobre, C., Lima, E., Lefebvre, P., Schlesinger, P., Potter,
388 C., Moutinho, P., Mendoza, E., Cochrane, M., Brooks, V., 1999. Large-scale impoverishment of
389 Amazonian forests by logging and fire. *Nature*, 398, 505-508.

390 Pausas, J., 2004. Changes in fire and climate in the eastern Iberian peninsula (Mediterranean
391 Basin). *Clim. Change*, 63, 337-350

392 Reed, B., Brown, J., Vanderzee, D., Loveland, T., Merchant, J., Ohlen, D., 1994. Measuring
393 phenological variability from satellite imagery. *J. Veg. Sci.*, 15, 703-714

394 Riano, D., Moreno-Ruiz, J., Isidoros, D., Ustin, S., 2007. Global spatial patterns and temporal
395 trends of burned area between 1981 and 2000 using NOAA-NASA Pathfinder. *Glob. Chang.*
396 *Biol.*, 13, 40-50

397 Teillet, P., Guindon, B., Goodenough, D., 1982. On the slope-aspect correction of multispectral
398 scanner data. *Can. J. Remote Sens.*, 8, 84-106

399 Veraverbeke, S., Verstraeten, W., Lhermitte, S., Goossens, R., 2010a. Evaluation Landsat
400 Thematic Mapper spectral indices for estimating burn severity of the 2007 Peloponnese wildfires
401 in Greece. *Int. J. Wildland Fire*, in press.

402 Veraverbeke, S., Verstraeten, W.W., Lhermitte, S., Goossens, R., 2010b. Illumination effects on
403 the differenced Normalized Burn Ratio's optimality for assessing fire severity. *Int. J. Appl. Earth*
404 *Observation Geoinf.*, 12, 60-70

405 Veraverbeke, S., Lhermitte, S., Verstraeten, W.W., Goossens, R., 2010c. The temporal dimension
406 of differenced Normalized Burn Ratio (dNBR) fire/burn severity studies: the case of the large
407 2007 Peloponnese wildfires in Greece. *Remote Sens. Environ.*, in press.

408 Verbyla, D., Kasischke, E., Hoy, E., 2008. Seasonal and topographic effects on estimating fire
409 severity from Landsat TM/ETM+ data. *Int. J. Wildland Fire*, 17, 527-534

410 Vermote, E., El Saleous, N., Justice, C., 2002. Atmospheric correction of MODIS data in the
411 visible to middle infrared: first results. *Remote Sens. Environ.*, 83, 97-111

412 White, J., Ryan, K., Key, C., Running, S., 1996. Remote sensing of forest fire severity and
413 vegetation recovery. *Int. J. Wildland Fire*, 6, 125-136

414

415 Figure 1. Schematic representation of post-fire effects terminology (Veraverbeke et al. 2010a).

416 Figure 2. Pre-fire land cover types of the burned areas (Veraverbeke et al., 2010a). The locations of the example
417 pixels shown in figure 7 are also indicated (A-H).

418 Figure 3. Principle of the multi-temporal dNBR ($dNBR_{MT}$). The $dNBR_{MT}$ represents the averaged integrated
419 difference between the one-year post fire NBR time series of the control and focal pixels, as shown in the figure by
420 the shaded area.

421 Figure 4. MODIS $dNBR_{MT}$ map (A), subset MODIS $dNBR_{MT}$ map of the blue rectangle in A (B), Landsat dNBR
422 map (C) and subset Landsat dNBR map of the blue rectangle in C (D). The locations of the example pixels shown in
423 figure 7 are also indicated in A.

424 Figure 5. Scatter plot and regression line between Landsat dNBR and GeoCBI (A) and between MODIS $dNBR_{MT}$
425 and Landsat dNBR (B) ($n = 150$, $p < 0.001$). The vertical bars in B indicate the standard deviation of Landsat pixels
426 within one MODIS pixel.

427 Figure 6. Mean $dNBR_{MT}$ and standard deviation per land cover type.

428 Figure 7. Illustration of $dNBR_{MT}$ estimates (shaded area) for coniferous forest (A-B), shrub land (C-D), olive groves
429 (E-F) and broadleaved forest (G-H). The location of the pixels is given in figures 2 and 4A.