



A common founder for the 35delG *GJB2* gene mutation in connexin 26 hearing impairment

L Van Laer, P Coucke, R F Mueller, et al.

J Med Genet 2001 38: 515-518

doi: 10.1136/jmg.38.8.515

Updated information and services can be found at:

<http://jmg.bmj.com/content/38/8/515.full.html>

These include:

References

This article cites 29 articles, 8 of which can be accessed free at:

<http://jmg.bmj.com/content/38/8/515.full.html#ref-list-1>

Article cited in:

<http://jmg.bmj.com/content/38/8/515.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Genetic screening / counselling](#) (2131 articles)

[Molecular genetics](#) (2080 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://journals.bmj.com/cgi/ep>

A common founder for the 35delG *GJB2* gene mutation in connexin 26 hearing impairment

L Van Laer, P Coucke, R F Mueller, G Caethoven, K Flothmann, S D Prasad, G P Chamberlin, M Houseman, G R Taylor, C M Van de Heyning, E Fransen, J Rowland, R A Cucci, R J H Smith, G Van Camp

Abstract

Fifty to eighty percent of autosomal recessive congenital severe to profound hearing impairment result from mutations in a single gene, *GJB2*, that encodes the protein connexin 26. One mutation of this gene, the 35delG allele, is particularly common in white populations. We report evidence that the high frequency of this allelic variant is the result of a founder effect rather than a mutational hot spot in *GJB2*, which was the prevailing hypothesis. Patients homozygous for the 35delG mutation and normal hearing controls originating from Belgium, the UK, and the USA were genotyped for different single nucleotide polymorphisms (SNPs). Four SNPs mapped in the immediate vicinity of *GJB2*, while two were positioned up to 76 kb from it. Significant differences between the genotypes of patients and controls for the five SNPs closest to *GJB2* were found, with nearly complete association of one SNP allele with the 35delG mutation. For the most remote SNP, we could not detect any association. We conclude that the 35delG mutation is derived from a common, albeit ancient founder.

(J Med Genet 2001;38:515-518)

Keywords: connexin 26; *GJB2*; 35delG; founder effect

Congenital severe to profound hearing impairment affects 1 in 1000 neonates.¹ In half of these children, the hearing impairment is the result of genetic factors,² with autosomal recessive inheritance representing approximately 80% of this total.¹ Although 30 recessive genes have been localised (G Van Camp, R J H Smith. Hereditary Hearing Loss Homepage, <http://dnalab-www.uia.ac.be/dnalab/hhh>, March 2001), a large proportion of recessive severe to profound hearing impairment has been linked to the DFNB1 locus.³ In 1997, Kelsell *et al*⁴ showed that mutations in *GJB2* lead to hearing impairment at this locus. Mutations in the same gene also are responsible for autosomal dominant hearing impairment at the DFNA3 locus.⁵

GJB2 encodes connexin 26, a gap junction protein that is assumed to be a component of the potassium recycling pathway in the inner ear. Loss or malfunction of these gap junctions, as might be reflected by mutations in *GJB2*, may disrupt potassium movement from the hair cells through the supporting cell network

to the endolymph, leading to hearing impairment. The resultant hearing loss involves all frequencies, is of variable severity (from mild to profound) even within sibships, rarely progresses, and most frequently is symmetrical in the two ears.⁶⁻⁸

To date, 48 recessive and seven dominant disease causing *GJB2* mutations have been identified (R Rabionet, P Gasparini, X Estivill. Connexins and Deafness Homepage, <http://www.iro.es/cx26deaf.html>, March 2001). One mutation, the deletion of one guanosine residue from a stretch of six between nucleotide positions 30 and 35 (35delG), which results in a frameshift and premature protein truncation at codon 13, is the most common deafness causing allelic variant of *GJB2* in sporadic patients and autosomal recessive families from Italy, Spain, Portugal, France, UK, Israel, Lebanon, Tunisia, Algeria, New Zealand, and in white American families of northern and southern European origin.⁹⁻¹³

Carrier frequencies of the 35delG mutation have been determined in several countries (table 1), and in some populations this mutation is more frequent than the $\Delta F508$ cystic fibrosis mutation. However, in non-white populations, the 35delG mutation is either not found or is very rare, with other "common" mutations prevailing, like the 235delC in the Japanese and Koreans,¹⁴⁻¹⁶ the 167delT in the Ashkenazi Jewish,^{17 18} and the R143W mutation in an African village.¹⁹ Most investigators have assumed that the high frequency of the 35delG mutation reflects the presence of a mutational hot spot within *GJB2*,^{9 11 17 20 21} but others have suggested a founder effect.¹³ To resolve the issue, we performed genotyping in the immediate vicinity of the *GJB2* gene. From these data, we present evidence that the high frequency of the 35delG *GJB2* allelic variant in the white population is the result of a founder effect rather than a mutational hot spot.

Subjects and methods

SUBJECTS

In this study, the following subjects were genotyped: 35 Belgian, 30 British, and 49 American white patients with non-syndromic hearing impairment who were homozygous for the 35delG mutation, and 70 Belgian, 30 British, and 50 American white random normal hearing controls. The American control samples were screened and selected for the absence of the 35delG mutation. In the Belgian and British control samples, no 35delG genotyping was performed.

Department of Medical Genetics, University of Antwerp, Universiteitsplein 1, B-2610 Antwerp, Belgium

L Van Laer
P Coucke
G Caethoven
K Flothmann
C M Van de Heyning
E Fransen
G Van Camp

Molecular Medicine Unit and Yorkshire Regional DNA Laboratory, St James's University Hospital, Leeds, UK
R F Mueller
M Houseman
G R Taylor
J Rowland

Molecular Otolaryngology Research Laboratories, Department of Otolaryngology, Head and Neck Surgery, Iowa City, Iowa, USA
S D Prasad
G P Chamberlin
R A Cucci
R J H Smith

Correspondence to: Professor Van Camp, gvcamp@uia.ua.ac.be

Revised version received 26 April 2001
Accepted for publication 3 May 2001

Table 1 Carrier frequency of the 35delG mutation in various populations

Population	Subjects tested	Carriers 35delG	Carrier frequency (%)	Reference
<i>Europe</i>				
Norway	190	1	0.5	22
Denmark	95	2	2.1	22
Estonia	113	5	4.4	22
Germany	200	4	2.0	22
UK	119	0	0.0	22
Holland	89	2	2.2	22
Belgium	190	1	0.5	22
Belgium	360	9	2.5	23
France (Brittany)	96	1	1.0	22
France	200	1	0.5	22
France	68	0	0.0	9
France	512	14	2.7	24
Bulgaria	157	1	0.6	22
Czech Republic	195	4	2.1	22
Slovenia	182	1	0.5	22
Italy (Sardinia)	116	4	3.4	22
Italy	255	8	3.1	22
Italy	150	6	4.0	12
Spain	200	5	2.5	22
Spain	130	3	2.3	22
Portugal	179	4	2.2	22
Malta	144	4	2.8	22
Greece	400	12	3.0	22
Greece	395	14	3.5	25
Turkey	150	4	2.7	22
<i>North America</i>				
White Americans	173	1	0.6	17
White Americans	100	1	1.0	26
White Americans	96	2	2.0	11
White Americans	560	14	2.5	27
African Americans	173	0	0.0	17
African Americans	190	0	0.0	22
Asian Americans (Indian, Japanese, Koreans)	53	0	0.0	17
Ashkenazi Jewish Americans	551	4	0.7	17
<i>Other populations</i>				
Egyptians	95	0	0.0	22
Arabs	58	1	1.7	22
Koreans	100	1	1	15




Figure 1 (A) Positions of the SNPs relative to the *GJB2* gene. The *GJB2* open reading frame (ORF) is represented by the black box. The 3'UTR and the URR are represented by shaded boxes. The borders of the ORF, UTR, and URR, the positions of the six SNPs, and the 35delG mutation are indicated with their respective nucleotide numbers derived from the genomic sequence of clone RP11-264J4 (Genbank accession number AL138688). (B) The distance from each SNP to the 35delG mutation in the *GJB2* gene (given in bp).

SNP IDENTIFICATION

To identify SNPs (single nucleotide polymorphisms) in regions adjacent to the *GJB2* gene, the 3'UTR (untranslated region, Genbank accession number M86849), and the URR (upstream regulatory region, Genbank accession number AF091526) were sequenced in six 35delG carriers. Four SNPs were discovered, two in the 3'UTR (SNP 1 and 2) and two in the URR (SNP 3 and 4). The relative positions of

the SNPs with regard to the *GJB2* gene are depicted in fig 1, where the SNPs are placed on the complete nucleotide sequence of clone RP11-264J4 (Genbank accession number AL138688). Several SNPs more distant from the *GJB2* gene were chosen from the SNP Consortium LTD (<http://snp.cshl.org/>) and analysed in random subjects. Two informative SNPs, SNP 5 (TSC1102325) and SNP 6 (TSC0127952), were selected for further analysis and positioned on the nucleotide sequence of clone RP11-264J4 (fig 1).

SNP ANALYSIS

SNP analysis was performed using the ABI Prism SNaPshot ddNTP Primer Extension Kit (ABI) according to the manufacturer's guidelines with minor modifications. PCR was performed using standard conditions and the primers and annealing temperatures listed in table 2. PCR products were treated with 0.164 U/μl shrimp alkaline phosphatase (Promega) and 0.164 U/μl exonuclease I (New England Biolabs) for one hour at 37°C, followed by the deactivation of the enzymes for 15 minutes at 72°C. SNaPshot thermal cycling was performed with the respective SNaPshot primer (table 2), the matching PCR product, and the SNaPshot Ready Reaction Premix for 25 cycles of 10 seconds at 96°C denaturation, five seconds at 50°C annealing, and 30 seconds at 60°C extension. SNaPshot products were dephosphorylated by incubation at 37°C (one hour) with 0.15 U/μl calf intestine alkaline phosphatase (Gibco BRL Life Technologies). The enzyme was deactivated as described above and SNaPshot products were analysed on an ABI3100 automated DNA sequencer (ABI).

STATISTICAL ANALYSIS

Differences between groups were tested with χ^2 statistics, if at least 80% of the expected frequencies exceeded 5 and all the expected frequencies exceeded 1. If the expected values did not meet the criteria mentioned above, the numbers of the two least frequent alleles were added and Yates's correction for continuity for two by two tables was used. If the expected values of the two by two table still did not meet the criteria mentioned above, Fisher's exact test was used. Both the χ^2 test and Fisher's exact test were calculated using the SigmaStat 2.0 software (Jandel Corporation). All p values were taken to be significant at <0.05.

Results

In order to determine whether the high frequency of the 35delG allele variant of *GJB2* is the result of a mutational hot spot or a founder effect, SNPs adjacent to (SNPs 1 to 4, fig 1) and more distant from (SNPs 5 and 6, fig 1) *GJB2* were analysed. Table 3 summarises the genotypes observed in this study. SNP 1 is very informative and results, even with small sample numbers, in significantly different genotypes for controls and patients. SNPs 2 and 3 are less informative, as the allele that is linked to 35delG (G for SNP 2 as well as for SNP 3) is also the most frequent allele in the

Table 2 PCR and SNaPshot primers

SNP	Primer	Primer sequence	Annealing temperature (°C)	Length of PCR product (bp)
SNP 1	F	aggcactggtaactttgttc	59	589
	R	gtgtctggaatttgcacctgc		
SNP 2	SNaPshot	agctgtcaaggctcagtcgc	59	901
	F	caatgctattcttgacaacagg		
SNP 3	R	ccacgttaagggtgaacattgg	56	676
	SNaPshot	acctctttaaatcagcgctctttcc		
SNP 4	F	ctgtgtgatgttgtagtcg	56	502
	R	gacaccacagtggattttcc		
SNP 5	SNaPshot	ctgccctagaggagctctgagcctactgg	56	230
	F	gcttaggagtagtactgctgc		
SNP 6	R	cacgggtccattccatcagg	55	381
	SNaPshot	ctcattgacgtgcagagactgaccagatagagaag		
SNP 7	F	ctgagtagcaatctcaccac	55	381
	R	tgacacgggataataaccgc		
SNP 8	SNaPshot	gggccccttaaacatgaggc	55	381
	F	taacaaggccctccaagtgc		
SNP 9	R	gcttgccatcaagaaggag	55	381
	SNaPshot	tcagcgctcatctcttggactgca		

general population. However, when using larger sample numbers even these less informative SNPs show significant differences between controls and patients. SNP 4 is informative, giving significant differences even when using smaller sample numbers. SNP 6, which is approximately 63 kb away from the *GJB2* gene, only shows significant differences between patients and controls in the total test population (fig 1, table 1). For SNP 5, which is the most remote SNP we investigated (approximately 76 kb away from *GJB2*, fig 1), no significant association with the 35delG mutation could be found. We conclude that the high frequency of the 35delG mutation in *GJB2* is the result of a founder effect.

The 35delG mutation must have arisen many centuries ago as haplotype sharing is observed only in a small region. On the basis of the recombination frequency of a more distant SNP, we can estimate the age of the 35delG mutation. From one generation to the next, the probability of having a recombination between a gene and a nearby marker is equal to the recombination fraction θ . In one generation, a fraction θ of all sibs will have a recombination and a fraction $1-\theta$ will not. After N generations, the fraction of descendants that has no recombination equals $(1-\theta)^N$. When examining SNP 6 in the total test population and assuming A was the allele associated with the 35delG mutation

in the founder, 161 out of 224 chromosomes showed no recombination in a region of approximately 63 kb (assumed to correspond to $\theta=0.00063$). Consequently, the 35delG mutation is estimated to be about 500 generations, or approximately 10 000 years old. As we do not know the true recombination frequency of this region, it should be emphasised that this calculation represents no more than a rough estimation, and that the range of error for this calculation is undoubtedly large.

Discussion

In this study, we show that the high frequency of the 35delG mutation is the result of a founder effect rather than a mutational hot spot, although the latter hypothesis has been favoured by most authors.^{9 11 17 20 21} It has been argued that the deletion of one G from a stretch of six could have arisen frequently through polymerase slippage or the presence of a chi consensus motif in the immediate vicinity.^{9 11} A strong argument for a mutational hot spot is certainly the presence of this mutation in several geographically and ethnically distinct populations.²¹ However, its absence or low prevalence in other ethnic groups, such as Asians,^{14 15} argues against this hypothesis.

Morell *et al*¹⁷ have shown that several different haplotypes surround the 35delG mutation,¹⁷ another strong argument in favour of a mutational hot spot. Their study considered an interval of 2 cM, while we have focused on a region of less than 70 kb or more than 95% smaller. As we have shown, the 35delG mutation is ancient and therefore haplotype sharing will be observed only in a very small chromosomal interval.

The age of the mutation further explains its widespread occurrence across Europe, the Mediterranean region, and the United States. Possibly, the mutation originated somewhere in the Middle East and was spread throughout Europe along the two Neolithic population movement routes. The first route followed the coast of the Mediterranean sea to Italy and Spain, while the other followed the Danube and Rhine valleys to northern Europe. The 35delG mutation would have been brought to the United States by emigration from Europe.

Table 3 Results of the SNP analysis

SNP	Genotype	Belgian			British			American			Total		
		Patient	Control	p	Patient	Control	p	Patient	Control	p	Patient	Control	p
SNP 1	CC	0	47	<0.001	0	18	<0.001	0	31	<0.001	0	96	<0.001
	CT	0	13		2	12		2	18		4	43	
	TT	33	1		27	0		45	1		105	2	
SNP 2	GG	31	20	0.014	27	24	0.025	47	43	0.013	105	87	<0.001
	GA	0	5		0	6		0	7		0	18	
SNP 3	AA	0	0	0.053	0	0	0.052	0	0	0.003	0	0	<0.001
	GG	33	29		29	25		49	41		111	95	
	GA	0	5		0	5		0	9		0	19	
SNP 4	AA	0	0	0.008	0	0	<0.001	0	0	<0.001	0	0	<0.001
	TT	31	22		28	20		48	35		107	77	
	TG	0	6		0	10		0	14		0	30	
SNP 5	GG	0	0	0.611	0	0	0.472	0	0	0.613	0	0	0.358
	GA	26	21		22	20		35	31		83	72	
	AA	6	8		5	10		13	13		24	31	
SNP 6	AA	2	2	0.072	1	0	0.431	1	4	0.224	4	6	0.019
	GG	6	5		6	3		3	6		15	14	
	GA	10	17		6	13		17	22		33	52	
SNP 7	AA	19	9		17	13		28	22		64	44	
	GA												

The low carrier frequency seen in other populations, for example, in the Korean and Arab population, could also have originated from the same founder. On the other hand, however, it cannot be excluded that the 35delG mutation has arisen on more than one occasion.

The reason for the high frequency of 35delG remains to be elucidated. By comparison, the $\Delta F508$ cystic fibrosis mutation is by far the best known and studied ancient mutation. Its frequency varies in different populations and estimates of age of origin of the mutation range, from 2940 to 52 000 years, depending on the methods used.^{28 29} It has been postulated that $\Delta F508$ carriers have an increased resistance to chloride secreting diarrhoea resulting in selective advantage for heterozygous carriers.³⁰ At present, it is not known whether the carrier status for a mutation in *GJB2* is associated with increased reproductive fitness.

The authors wish to thank Dr Rudy D'Hooghe for statistical advice, Drs Declau, De Meirleir, Goffart, Govaerts, Haentjens, Janssens, Lissens, Mansbach, Messiaen, Mortier, Officiers, Raes, Van de Heyning, van den Ende, Verstreken, and Verysen for samples from congenital hearing impaired patients, and Sandra Wilcox for 35delG genotyping. This study was supported by grant G.0277.01 from the FWO (Flemish Fonds voor Wetenschappelijk Onderzoek) and grant RO1-DC02842 from the NIDCD (RJHS). LVL, EF, and GVC hold research positions with the FWO.

- Morton NE. Genetic epidemiology of hearing impairment. *Ann NY Acad Sci* 1991;630:16-31.
- Marazita ML, Ploughman LM, Rawlings B, Remington E, Arnos KS, Nance WE. Genetic epidemiological studies of early-onset deafness in the U.S. school-age population. *Am J Med Genet* 1993;46:486-91.
- Gasparini P, Estivill X, Volpini V, Totaro A, Castellvi-Bel S, Govea N, Mila M, Della Monica M, Ventruto V, De Benedetto M, Stanziale P, Zelante L, Mansfield ES, Sandkuijl L, Surrey S, Fortina P. Linkage of DFNB1 to non-syndromic neurosensory autosomal-recessive deafness in Mediterranean families. *Eur J Hum Genet* 1997;5:83-8.
- Kelsell DP, Dunlop J, Stevens HP, Lench NJ, Liang JN, Parry G, Mueller RF, Leigh IM. Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature* 1997;387:80-3.
- Denoyelle F, Lina-Granade G, Plauchu H, Bruzzone R, Chaib H, Levi-Acobas F, Weil D, Petit C. Connexin 26 gene linked to a dominant deafness. *Nature* 1998;393:319-20.
- Cohn ES, Kelley PM, Fowler TW, Gorga MP, Lefkowitz DM, Kuehn HJ, Schaefer GB, Gobar LS, Hahn FJ, Harris DJ, Kimberling WJ. Clinical studies of families with hearing loss attributable to mutations in the connexin 26 gene (GJB2/DFNB1). *Pediatrics* 1999;103:546-50.
- Denoyelle F, Marlin S, Weil D, Moatti L, Chauvin P, Garabedian EN, Petit C. Clinical features of the prevalent form of childhood deafness, DFNB1, due to a connexin-26 gene defect: implications for genetic counselling. *Lancet* 1999;353:1298-303.
- Murgia A, Orzan E, Polli R, Martella M, Vinanzi C, Leonardi E, Arslan E, Zaccchello F. Cx26 deafness: mutation analysis and clinical variability. *J Med Genet* 1999;36:829-32.
- Denoyelle F, Weil D, Maw MA, Wilcox SA, Lench NJ, Allen-Powell DR, Osborn AH, Dahl HHM, Middleton A, Houseman MJ, Dodé C, Marlin S, Boulila-ElGated A, Grati M, Ayadi H, BenArab S, Bitoun P, Lina-Granade G, Godet J, Mustapha M, Loiselet J, El-Zir E, Aubeis A, Joannard A, Levilliers J, Garabedian EN, Mueller RF, Gardner RJM, Petit C. Prelingual deafness: high prevalence of a 30delG mutation in the connexin 26 gene. *Hum Mol Genet* 1997;6:2173-7.
- Zelante L, Gasparini P, Estivill X, Melchionda S, D'Agruma L, Govea N, Mila M, Della Monica M, Lutfi J, Shohat M, Mansfield E, Delgrosso K, Rappaport E, Surrey S, Fortina P. Connexin26 mutations associated with the most common form of non-syndromic neurosensory autosomal recessive deafness (DFNB1) in Mediterraneans. *Hum Mol Genet* 1997;6:1605-9.
- Kelley PM, Harris DJ, Comer BC, Askew JW, Fowler T, Smith SD, Kimberling WJ. Novel mutations in the connexin 26 gene (GJB2) that cause autosomal recessive (DFNB1) hearing loss. *Am J Hum Genet* 1998;62:792-9.
- Estivill X, Fortina P, Surrey S, Rabionet R, Melchionda S, D'Agruma L, Mansfield E, Rappaport E, Govea N, Mila M, Zelante L, Gasparini P. Connexin-26 mutations in sporadic and inherited sensorineural deafness. *Lancet* 1998;351:394-8.
- Rabionet R, Zelante L, Lopez-Bigas N, D'Agruma L, Melchionda S, Restagno G, Arbones ML, Gasparini P, Estivill X. Molecular basis of childhood deafness resulting from mutations in the *GJB2* (connexin 26) gene. *Hum Genet* 2000;106:40-4.
- Abe S, Usami S, Shinkawa H, Kelley PM, Kimberling WJ. Prevalent connexin 26 gene (*GJB2*) mutations in Japanese. *J Med Genet* 2000;37:41-3.
- Park HJ, Hahn SH, Chun YM, Park K, Kim HN. Connexin26 mutations associated with nonsyndromic hearing loss. *Laryngoscope* 2000;110:1535-8.
- Kudo T, Ikeda K, Kure S, Matsubara Y, Oshima T, Watanabe K, Kawase T, Narisawa K, Takasaka T. Novel mutations in the connexin 26 gene (GJB2) responsible for childhood deafness in the Japanese population. *Am J Med Genet* 2000;90:141-5.
- Morell RJ, Kim HJ, Hood LJ, Goforth L, Friderici K, Fisher R, Van Camp G, Berlin CI, Oddoux C, Ostrer H, Keats B, Friedman TB. Mutations in the connexin 26 gene (*GJB2*) among Ashkenazi Jews with nonsyndromic recessive deafness. *N Engl J Med* 1998;339:1500-5.
- Sobe T, Erlich P, Berry A, Korostichevsky M, Vreugde S, Avraham KB, Bonne-Tamir B, Shohat M. High frequency of the deafness-associated 167delT mutation in the connexin 26 (*GJB2*) gene in Israeli Ashkenazim. *Am J Med Genet* 1999;86:499-500.
- Brobbly GW, Müller-Myhsok B, Horstmann RD. Connexin 26 R143W mutation associated with recessive nonsyndromic sensorineural deafness in Africa. *N Engl J Med* 1998;338:548-9.
- Carrasquillo MM, Zlotogora J, Barges S, Chakravarti A. Two different connexin 26 mutations in an inbred kindred segregating non-syndromic recessive deafness: implications for genetic studies in isolated populations. *Hum Mol Genet* 1997;6:2163-72.
- Lench NJ, Markham AF, Mueller RF, Kelsell DP, Smith RJH, Willems PJ, Schatteman I, Capon H, Van De Heyning PJ, Van Camp G. A Moroccan family with autosomal recessive sensorineural hearing loss caused by a mutation in the gap junction protein gene connexin 26 (GJB2). *J Med Genet* 1998;35:151-2.
- Gasparini P, Rabionet R, Barbujani G, Melchionda S, Petersen M, Brondum-Nielsen K, Metspalu A, Oitmaa E, Pisano M, Fortina P, Zelante L, Estivill X. The Genetic Analysis Consortium of *GJB2* 35delG. High carrier frequency of the 35delG deafness mutation in European populations. *Eur J Hum Genet* 2000;8:19-23.
- Storm K, Wilcox S, Flothmann K, Van Camp G. Determination of the carrier frequency of the common GJB2 (connexin-26) 35delG mutation in the Belgian population using an easy and reliable screening method. *Hum Mutat* 1999;14:263-6.
- Lucotte G, Bathelier C, Champenois T. PCR test for diagnosis of the common GJB2 (connexin 26) 35delG mutation on dried blood spots and determination of the carrier frequency in France. *Mol Cell Probes* 2001;15:57-9.
- Antoniadi T, Rabionet R, Kroupis C, Aperis GA, Economides J, Petmezakis J, Economou-Petersen E, Estivill X, Petersen MB. High prevalence in the Greek population of the 35delG mutation in the connexin 26 gene causing prelingual deafness. *Clin Genet* 1999;55:381-2.
- Scott DA, Kraft ML, Carmi R, Ramesh A, Elbedour K, Yairi Y, Srisailapathy CRS, Rosengren SS, Markham AF, Mueller RF, Lench NJ, Van Camp G, Smith RJH, Sheffield VC. Identification of mutations in the connexin 26 gene that cause autosomal recessive nonsyndromic hearing loss. *Hum Mutat* 1998;11:387-94.
- Green GE, Scott DA, McDonald JM, Woodworth GG, Sheffield VC, Smith RJH. Carrier rates in the midwestern United States for GJB2 mutations causing inherited deafness. *JAMA* 1999;281:2211-16.
- Morral N, Bertranpetit J, Estivill X, Nunes V, Casals T, Gimenez J, Reis A, Varon-Mateeva R, Macek MJ, Kalaydjieva L, Angelicheva D, Dancheva R, Romeo G, Russo MP, Garnerone S, Restagno G, Ferrari M, Magnani C, Claustres M, Desgeorges M, Schwartz M, Schwarz M, Dalapiccola B, Novelli G, Ferec C, de Arce M, Nemeti M, Kere J, Anvret M, Dahl N, Kadasi L. The origin of the major cystic fibrosis mutation (delta F508) in European populations. *Nature* 1994;7:169-75.
- Serre JL, Simon-Bouy B, Mornet E, Jaume-Roig B, Balassopoulou A, Schwartz M, Taillandier A, Boue J, Boue A. Studies of RFLP closely linked to the cystic fibrosis locus throughout Europe lead to new considerations in populations genetics. *Hum Genet* 1990;84:449-54.
- Romeo G, Devoto M, Galietta LJ. Why is the cystic fibrosis gene so frequent? *Hum Genet* 1989;84:1-5.