

Accepted Manuscript

Behavioral Interventions in Attention-Deficit/Hyperactivity Disorder: A Meta-Analysis of Randomized Controlled Trials across Multiple Outcome Domains

David Daley, PhD Saskia van der Oord, PhD Maite Ferrin, MD, PhD Marina Danckaerts, MD, PhD Manfred Doepfner, PhD Samuele Cortese, MD, PhD Edmund J.S. Sonuga-Barke, PhD

PII: S0890-8567(14)00408-0

DOI: [10.1016/j.jaac.2014.05.013](https://doi.org/10.1016/j.jaac.2014.05.013)

Reference: JAAC 1027

To appear in: *Journal of the American Academy of Child & Adolescent Psychiatry*


Received Date: 13 November 2013

Revised Date: 19 May 2014

Accepted Date: 20 May 2014

Please cite this article as: Daley D, van der Oord S, Ferrin M, Danckaerts M, Doepfner M, Cortese S, Sonuga-Barke EJS, , Behavioral Interventions in Attention-Deficit/Hyperactivity Disorder: A Meta-Analysis of Randomized Controlled Trials across Multiple Outcome Domains, *Journal of the American Academy of Child & Adolescent Psychiatry* (2014), doi: 10.1016/j.jaac.2014.05.013.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



<LRH>Daley *et al.*

<RRH>Behavioral interventions for ADHD

<DOC>New research

<AT>Behavioral Interventions in Attention-Deficit/Hyperactivity Disorder: A Meta-Analysis of Randomized Controlled Trials across Multiple Outcome Domains

<AU>David Daley, PhD, Saskia van der Oord, PhD, Maite Ferrin, MD, PhD, Marina Danckaerts, MD, PhD, Manfred Doepfner, PhD, Samuele Cortese, MD, PhD, Edmund J.S. Sonuga-Barke, PhD, on behalf of the European ADHD Guidelines Group

<ARTFN>This article is discussed in an editorial by Dr. Linda Pfiffner on page xxx.

Supplemental material cited in this article is available online.

<BEGIN ABSTRACT>Objective: Behavioral interventions are recommended as attention-deficit/hyperactivity disorder (ADHD) treatments. However, a recent meta-analysis found no effects on core ADHD symptoms when raters were probably blind to treatment allocation. The present analysis is extended to a broader range of child and parent outcomes. Method: A systematic search in PubMed, Ovid, Web of Knowledge, ERIC, and CINAHAL databases (up to February 5, 2013) identified published randomized controlled trials measuring a range of patient and parent outcomes for children and adolescents diagnosed with ADHD (or who met validated cutoffs on rating scales). Results: Thirty-two of 2,057 nonduplicate screened records were analyzed. For assessments made by individuals closest to the treatment setting (usually unblinded), there were significant improvements in parenting quality (standardized mean difference [SMD] for positive parenting 0.68; SMD for negative parenting 0.57), parenting self-concept (SMD 0.37), and child ADHD (SMD 0.35), conduct problems (SMD 0.26), social skills (SMD 0.47), and academic performance (SMD 0.28). With probably blinded assessments, significant effects persisted for parenting (SMD for positive parenting 0.63; SMD for negative parenting 0.43) and conduct problems (SMD 0.31). Conclusion: In contrast to the lack of blinded evidence of ADHD symptom decrease, behavioral interventions have positive effects on a range of other outcomes when used with patients with ADHD. There is blinded evidence that they improve parenting and decrease childhood conduct problems. These effects also may feed through into a more positive parenting self-concept but not improved parent mental well-being. *J. Am. Acad. Child Adolesc. Psychiatry*, 2014;16(x):xx-xx. Key Words: ADHD, parenting, intervention, conduct<END ABSTRACT>

Attention-deficit/hyperactivity disorder (ADHD) is characterized by age-inappropriate, persistent, and pervasive inattention and/or overactivity/impulsiveness that impairs daily functioning¹ and is associated with substantial long-term burden on patients, families, and health and educational services.^{2,3} Multimodal treatment approaches are recommended.⁴ Medication is typically used as the first-line intervention, especially for severe cases.⁵ Despite robust evidence of medium-term symptom control,⁶ medication has some limitations. A proportion of patients shows partial or no response.⁶ Long-term effectiveness remains to be established.^{7,8} Important aspects of functioning may not improve (e.g., academic achievement⁹). Adverse effects on sleep, appetite, and growth, although rarely serious and generally manageable, are common and may not be well tolerated.¹⁰ Treatment compliance can be low, especially during adolescence.¹¹ Parents and clinicians can have reservations about medication use¹² and may prefer nonpharmacologic approaches.¹³

Interventions using behavioral techniques also are recommended and commonly used as ADHD treatments.¹⁴ Systematic reviews of treatment trials have provided evidence to support their efficacy.¹⁵⁻¹⁷ However, these reviews can be difficult to interpret, because they sometimes include nonrandomized controlled trials (RCTs), mix individuals with and without ADHD, and have not always drawn clear boundaries between ADHD-specific and other outcomes. Furthermore, outcome assessment is often made unblinded by individuals taking an active part in the intervention (e.g., parents receiving parent training), which is likely to inflate efficacy estimates.¹⁸ Sonuga-Barke *et al.*¹⁹ published a meta-analysis of RCTs of behavioral interventions. Stringent inclusion and exclusion criteria addressed some limitations of previous meta-analyses. There was a moderate, statistically significant, positive effect on ADHD core symptoms for assessments made by individuals most proximal to the therapeutic setting—typically unblinded parent ratings. However, these effects were not corroborated by probably blinded measurements made by observers or raters unaware of treatment allocation when the effect size decreased to near 0 and became nonsignificant. A similar, although less marked, decrease was found for neural feedback and cognitive training. There are some possible explanations for these findings. First, that unblinded raters are biased and overestimate treatment effects.²⁰ Second, that interventions increase parental tolerance for ADHD or their ability to cope with its negative impact rather than decreasing symptom levels.¹⁹ Third, that probably blinded measurements were less valid than most proximal measurements.²¹ Fourth, that intervention effects did not generalize from the therapeutic setting (e.g., the home) to other settings (e.g., school).²² The authors concluded that more evidence from studies with blinded assessments is required before behavioral interventions can be supported as treatments for core ADHD symptoms.

The limited effects of behavioral approaches on blinded core ADHD measurements may be explicable if one considers the treatment models on which many are based. For instance, although most treatments in the trials included in the meta-analysis of Sonuga-Barke *et al.*¹⁹ were implemented to target

ADHD symptoms, they were initially developed and have been used extensively for children with oppositional and conduct problems.²³⁻²⁵ For many of these, the rationale is that children's challenging behavior develops because of coercive interactional cycles that, over time, co-reinforce noncompliant and oppositional behaviors in the child and negative and inappropriate responses from significant adults (usually the parent but also potentially teachers and other caregivers).²⁶ During intervention, the adult is taught to apply behavior modification techniques to reinforce appropriate and discourage inappropriate child behaviors, enhance effective and enjoyable adult-child interactions,²⁷ and so transform negative into positive interactional cycles. Such interventions, it could be argued, are unlikely to be effective as treatments for core ADHD symptoms because ADHD does not emerge along a similar environmentally mediated route as conduct problems and therefore is less likely to respond to the modification of environmental contingencies.²⁸

However, the value of behavioral interventions does not rest exclusively on their potential effects on ADHD symptoms. Patients with ADHD often have conduct problems²⁹ and other comorbidities³⁰ in addition to poor social and organizational skills and low academic achievement. Their parents can have poor parenting self-concept and mental health problems.³¹ These associated features of the disorder are important treatment targets in and of themselves, because each is associated with substantial burden to the child, the child's family, and society through the criminal justice, social, and health systems.^{22,32} Behavioral interventions may have an important role in treating these problems whether or not they decrease core ADHD symptoms. Indeed, behavioral treatments used with patients with ADHD have targeted ADHD-related but nonspecific aspects of impairment rather than ADHD symptoms themselves (e.g., social skills,³³ organizational skills,³⁴ and academic achievement³⁵).

In this article, the authors build on the previous meta-analysis¹⁹ to address the broader impact of behavioral interventions for children with ADHD. They address 3 related questions. First, given that most, although not all, interventions are implemented by changes in the behavior of responsible adults (typically parents or teachers), do behavioral interventions improve adult responses to children with ADHD? Second, do they improve the sense of efficacy and competence and decrease the mental health problems of adults working with children with ADHD? Third, do they decrease levels of child oppositional behavior and other comorbidities and other aspects of impairment such as social skills and academic performance? To address these questions, *most proximal* and *probably blinded* assessments were contrasted.

<H1>Method

Please see the registered protocol CRD42011001393 (link deleted to blind the identity of the authors) for more details.

<H2>Inclusion Criteria

Only published peer-reviewed RCTs were included, although the authors acknowledge that many well-designed studies using single-subject research designs examining the effects of behavioral interventions have been published. Following the recommendation of the Cochrane group, the search was limited to published trials to ensure a level of methodologic adequacy and rigor among included trials and to avoid the inevitable problems with securing access to a full set of unpublished trials and the bias that this would introduce.³⁶ Participants needed to be 3 to 18 years old and have an ADHD diagnosis (any subtype) or have met accepted cutoffs on validated ADHD rating scales. Trials involving only rare comorbid disorders (e.g., fragile X syndrome) were excluded. Acceptable control conditions were “treatment as usual,” “wait list,” or “active” controls. “Treatment as usual” could include medication, but trials were excluded if the behavioral intervention was an adjunct to medication or if pharmacologic and behavioral interventions were combined into 1 therapeutic arm as part of the study design. For the present extended review, trials could be included despite not having an ADHD-related outcome (as required in the original protocol).

<H2>Search Strategy

The search was updated to February 5, 2013. Drs. Cortese and Ferrin blindly conducted and cross-checked the updated search using the same databases, search strategy, and search terms as used previously¹⁹ (see protocol). The searches were conducted for records included from the inception of the databases. Behavioral interventions were defined as those interventions directed at changing behaviors (increasing desired and decreasing undesired behaviors). They encompass classic contingency management, behavior therapy (mainly through mediators such as parents or teachers), and cognitive behavior therapy (such as verbal self-instruction, problem-solving strategies, or social skills training). The treatment search terms covered a wide variety of intervention types with the aim of including trials involving any form of behaviorally based therapies, implemented in any setting (home or school), and indirectly by an adult or directly to the child (see protocol).

<H2>Outcome Measurements

To increase analytical robustness, outcome domains were only considered if at least 5 RCTs were available. Outcome measurements meeting this criterion were pre- to post-treatment changes in positive and negative parenting, parent mental health (e.g., anxiety, depression) and parenting self-concept (e.g., sense of competence and efficacy), child ADHD, conduct problems (i.e., negative and noncompliant behavior including symptoms of oppositional defiance and conduct disorders), social skills, and academic achievement. There were too few RCTs ($n < 5$) to examine changes in teacher behavior and well-being,

child impairment, internalizing problems, executive/organizational skills, or more general measurements of family functioning.

<H2>Study Selection

Article titles and abstracts were screened. Final inclusion was based on the full text. Trials were blindly double-coded for eligibility. Study quality was assessed by 2 independent raters according to the criteria of Jadad *et al.*³⁷ (Table 1).^{24,25,33-35,40-66} These provide a rating for each trial in terms of standard definitions for randomization, blinding, and treatment of missing data defined by Jadad *et al.*³⁷ Jadad scores for blinding were adapted for use with multiple outcomes so that studies with at least 1 blinded outcome yielded a score of 1 on this dimension. A score of at least 3 is regarded as indicating acceptable quality. Initial disagreements ($n = 4$) were resolved by the coders through discussion without recourse to an independent arbitrator.

<H2>Data Extraction and Statistical Analysis

Trial information was entered into RevMan 5.1 (<http://ims.cochrane.org/revman>).³⁸ Data extraction was independently rated by 2 authors. The standardized mean difference (SMD), namely the mean pre- to post-treatment change minus the mean pre- to post-treatment control group change divided by the pooled pretest standard deviation with a bias adjustment, was calculated.³⁹ SMDs for trials in each domain were combined using the inverse-variance method.³⁶ Given the inherent heterogeneity of studies, random-effects models were used. The I^2 statistic was calculated, a posteriori, to estimate between-trial SMD heterogeneity. Most proximal and probably blinded analyses were performed for all domains except parent mental health, parenting self-concept, child social skills, and academic outcomes for which insufficient trials with probably blinded measurements were available. When multiple measurements were available for an outcome, the one most frequently reported across included trials was included. For the most proximal analysis, parent ratings, if available, were used, except for teacher-based interventions, when teacher ratings or direct observations were preferred. Probably blinded assessments were made by an individual likely to be blind to allocation. In trials in which more than 1 such measurement was available, the best blinded measurement was selected. This affected only trials with a home-based element where direct observations by an independent researcher and teacher ratings were the probably blinded measurement. In such cases, direct observation was selected over teacher ratings. Sensitivity analyses examined the impact of background ADHD medication use in trial samples for which at least 3 trials had fewer than 30% of participants receiving medications (i.e., were no-/low-medication trials) and the effects of outliers identified using funnel plots within RevMan 5.1.³⁸ Meta-regression tested whether effect sizes were larger in lower-quality trials according to Jadad *et al.*³⁷ For 1 study,³⁴ 2 active treatment arms were suitable for analysis. Parents and Teachers Helping Children Organize and Organizational Skills

Treatment yielded similar results; so only 1 arm, Parents and Teachers Helping Children Organize, which was considered the more standard behavioral intervention, was included in the final analysis. For another study,⁴⁰ outcomes were taken from 3 publications.⁴⁰⁻⁴²

<H1>Results

Thirty-two studies met the entry criteria and had data amenable to analysis. Eight had a Jadad study quality rating of at least 3.³³ Thirty-one studies had a parent-based component implemented at home. Four had an additional school-based, teacher-focused element. Fourteen included direct intervention with the child. Although most used standard behavioral principles and targeted children's externalizing behaviors (ADHD, oppositional defiance disorder, conduct disorder), some implemented specialized social ($n = 3$), organizational ($n = 3$), or academic ($n = 1$) skills. Figure 1 presents the trial selection flowchart. Tables 1 and 2 present information about included trials and the measurements used in each analysis, respectively.

<H2>Do Behavioral Interventions Improve Adult Responses to Children with ADHD?

No trials measured responses to children with ADHD by adults other than parents (i.e., no teachers). Nine trials measured positive parenting. Three most proximal measurements were parent rated; 5 were observational. Eight trials had probably blinded measurements, all observational. For positive parenting, most proximal effects were significant (SMD 0.68; 95% confidence interval [CI] 0.27-1.09; Figure 2 shows forest plots). Effects were similar for probably blinded outcomes (SMD 0.63, 95% CI 0.47-0.78). Heterogeneity was significant in the 2 analyses (χ^2 for most proximal = 63.55, $I^2 = 87\%$, $p < .01$; χ^2 for probably blinded = 40.58, $I^2 = 83\%$, $p < .01$). Effects were unaffected by limiting the analyses to no-/low-medication trials ($n = 5$, SMD for most proximal 1.23, 95% CI 0.26-2.20; $n = 4$, SMD for probably blinded 0.89, 95% CI 0.65-2.13), although heterogeneity remained high (χ^2 for most proximal = 28.35, $I^2 = 89\%$, $p < .01$; χ^2 probably blinded = 23.29, $I^2 = 87\%$, $p < .01$). Removing outliers decreased effect sizes ($n = 2$, SMD most proximal 0.32, 95% CI -0.06 to 0.58; $n = 1$, SMD probably blinded 0.44, 95% CI 0.14-0.75). Heterogeneity was no longer significant (χ^2 most proximal = 3.46, $I^2 = 0\%$, $p = .48$; χ^2 for probably blinded = 0.83, $I^2 = 0\%$, $p = .66$). Fourteen trials measured negative parenting. The 9 most proximal measurements were parent ratings (4 observations and 1 speech sample). Eight studies met criteria for probably blinded assessments (7 observations and 1 speech sample). For negative parenting, effects were significant for the 2 analyses (SMD for most proximal 0.57, 95% CI 0.37-0.78; SMD for probably blinded 0.43, 95% CI 0.24-0.62). Heterogeneity also was significant (χ^2 for most proximal = 32.7, $I^2 = 60\%$, $p < .01$; χ^2 for probably blinded = 19.8, $I^2 = 65\%$, $p < .01$). Most proximal effects were unchanged when no-/low-medication trials were analyzed ($n = 7$, SMD 0.57, 95% CI 0.22-0.92, $\chi^2 = 7.81$, $I^2 = 49\%$, $p < .01$), but effects became nonsignificant for probably blinded outcomes ($n = 6$, SMD 0.42,

95% CI -0.07 to 0.91, $\chi^2 = 14.25$, $I^2 = 72\%$, $p < .01$). When outliers were excluded, effects remained significant for the 2 outcomes ($n = 2$, SMD for most proximal 0.45, 95% CI 0.31-0.58; $n = 1$, SMD for probably blinded 0.36, 95% CI 0.12-0.60). Heterogeneity was decreased for one (χ^2 for most proximal = 15.45, $I^2 = 35\%$, $p = .12$) but not the other (χ^2 for probably blinded = 15.03, $I^2 = 73\%$, $p < .01$).

<H2>Do Behavioral Interventions Improve the Self-Concept and Mental Health of Adults Working with Children with ADHD?

No trials measured the self-concept and mental health of adults other than parents. Seven trials included self-ratings of parental self-concept (6 parenting efficacy/sense of competence, 1 parenting self-esteem). There was a small but significant improvement in self-concept after treatment (SMD 0.37, 95% CI 0.03-0.70). Heterogeneity was significant ($\chi^2 = 19.03$, $I^2 = 68\%$, $p < .01$). Effects increased slightly in no-/low-medication trials ($n = 5$, SMD 0.68, 95% CI 0.22-0.92), whereas heterogeneity decreased substantially ($\chi^2 = 0.99$, $I^2 = 0\%$, $p = .61$). When outliers were removed ($n = 1$), effects and their heterogeneity were decreased to nonsignificant levels (SMD 0.30, 95% CI -0.07 to 0.65, $\chi^2 = 5.24$, $I^2 = 4\%$, $p = .39$). Measurements of self-rated parent mental health were included in 9 studies (7 depression/anxiety, 2 more general well-being). There were no significant effects of treatment (SMD 0.09, 95% CI -0.09 to 0.23; no-/low-medication analysis, $n = 6$, SMD 0.09, 95% CI -0.19 to 0.37, $p = .13$; $\chi^2 = 5.63$, $I^2 = 47\%$, $p < .01$). No outliers were identified.

<H2>Do Behavioral Interventions Decrease Child Psychopathology and Improve Functioning?

Nineteen studies had most proximal ADHD measurements (4 in addition to the trials included in the study by Sonuga-Barke *et al.*²¹). The treatment effect was significant (SMD 0.35, 95% CI 0.19-0.50). Heterogeneity was significant ($\chi^2 = 32.63$, $I^2 = 45\%$, $p < .05$). Restricting the analysis to no-/low-medication enhanced the effect ($n = 11$, SMD 0.50, 95% CI 0.24-0.76), but heterogeneity remained significant ($\chi^2 = 20.33$, $I^2 = 51\%$, $p < .05$). Effects were decreased when outliers ($n = 3$) were removed but remained significant (SMD 0.23, 95% CI 0.12-0.35). Heterogeneity was no longer significant ($\chi^2 = 16.86$, $I^2 = 23\%$, $p = .21$). None of the trials added since the study by Sonuga-Barke *et al.*¹⁹ had a probably blinded measurement, so the prior finding of no treatment effect for ADHD remained unchanged (overall SMD 0.02, 95% CI -0.30 to 0.34).

Fifteen trials had most proximal measurements of conduct problems (10 parent ratings, 3 teacher ratings, and 2 clinic observations). Eight trials had probably blinded measurements (1 teacher rating and 7 observational ratings). Most proximal and probably blinded effects were significant (SMD for most proximal 0.26, 95% CI 0.14-0.37; SMD for probably blinded 0.31, 95% CI -0.05 to 0.57). Heterogeneity was significant for most proximal ($\chi^2 = 25.87$, $I^2 = 46\%$, $p = .03$) but not for probably blinded ($\chi^2 = 14.28$, $I^2 = 51\%$, $p = .05$) effects. Low-/no-medication studies yielded significant effects for the 2 analyses

($n = 7$, SMD for most proximal 0.54, 95% CI 0.32-0.76; $n = 6$, SMD for probably blinded 0.27, 95% CI = 0.02-0.51). Heterogeneity was not significant for either of these analyses (χ^2 for most proximal = 5.96, $I^2 = 37\%$, $p = .43$; χ^2 for probably blinded = 3.77, $I^2 = 0\%$, $p = .44$). Most proximal effects remained significant when outliers ($n = 2$) were removed (SMD 0.26, 95% CI = 0.12-0.46, $\chi^2 = 14.16$, $I^2 = 22\%$, $p = .22$). There were no probably blinded outliers.

Nine trials included most proximal social skills outcomes (4 parent and 5 teacher ratings). Four of these included a core specialized social skills training component. The effect was significant (SMD 0.47, 95% CI 0.15-0.78), as was heterogeneity ($\chi^2 = 22.98$, $I^2 = 65\%$, $p < .01$). Restricting the analysis to the no-/low-medication studies increased heterogeneity ($n = 5$, $\chi^2 = 22.08$, $I^2 = 86\%$, $p < .01$), so that although the effect size increased, it was no longer significant (SMD 0.67, 95% CI -0.20 to 1.55). When outliers ($n = 1$) were removed, the effect remained (SMD 0.30, 95% CI 0.07-0.52) and heterogeneity was nonsignificant ($\chi^2 = 4.04$, $I^2 = 0\%$, $p = .54$).

Academic achievement was measured in 9 trials (6 parent or teacher questionnaire-based measurements and 3 objective assessments). Seven studies had a specialized component to specially target this aspect of impairment. Most proximal effects were significant, and heterogeneity was nonsignificant (SMD 0.28, 95% CI 0.06-0.50, $\chi^2 = 14.55$, $I^2 = 45\%$, $p = .07$) and persisted when outliers ($n = 2$) were removed (SMD 0.16, 95% CI 0.01-0.31) and heterogeneity was nonsignificant ($\chi^2 = 7.95$, $I^2 = 37\%$, $p = .16$). There were insufficient no-/low-medication studies to conduct a sensitivity analysis (Figure 3).

There was no association between larger effect sizes and lower Jadad ratings. In fact, for most proximal ADHD ($t = 2.52$, $p = .02$), higher-quality trials yielded larger effects. Meta-regression did show larger effect sizes in trials with younger children for most proximal positive parenting ($t = -2.63$, $p = .03$), most proximal ADHD ($t = -2.09$, $p = .05$), and most proximal conduct problems ($t = -2.46$, $p = .03$).

<H1>Discussion

The present meta-analyses found blinded evidence that behavioral interventions used to treat children and adolescents with ADHD had beneficial effects on important aspects of child and parent functioning. This finding contrasts with the lack of blinded evidence relating to ADHD symptoms reported by Sonuga-Barke *et al.*¹⁹ Specifically, in this analysis, behavioral interventions improved parenting, decreasing negative and increasing positive parenting, and decreased children's comorbid conduct problems.

Although improving parent functioning was rarely their primary therapeutic goal, nearly all behavioral interventions included parent training as a core therapeutic mechanism. Positive effects on these outcomes are therefore expectable and, in some ways, a necessary condition for subsequent impacts on children's behavior. Given the need to pool across very diverse and often rather general parenting

assessments, a fine-grained analysis of these effects was not possible. For instance, positive parenting analyses pooled studies with general assessments of overall increases in positive behaviors/strategies (e.g., praise, encouragement⁶¹), together with measurements of their appropriate use,²⁵ according to therapeutic models. This is important because, according to most models,²⁸ simply increasing levels of positive parenting may not be sufficient to produce positive changes in children's behavior. Objective effects on parenting were reflected in improved parenting self-concept: an empowering effect important in the process of breaking negative parent-child interaction cycles.³¹ Increased parenting confidence could come from working with experienced therapists who validate parents' approaches and/or implementing behavioral strategies and seeing their positive effects. It also may be an effect of psychoeducation that emphasizes the power parents have to influence developmental outcomes.⁶⁷ Given these parenting-specific positive effects, it was perhaps surprising that no beneficial effects were seen on parent mental health more generally. It would seem that the high levels of mental health problems that often affect parents of children with ADHD are not solely the result of issues around parenting morale, but rather reflect a more deep-seated, pervasive psychological aspect, perhaps reflecting shared genetic risk for mental health problems within families.^{68,69} There also was blinded evidence that behavioral interventions decreased conduct problems in children with ADHD; benefits of behavioral interventions for children seen with conduct disorder extend to those with a full ADHD diagnosis.²⁵ Given the limited number of studies with probably blinded measurements and the fact that few had objective measurements of child behavior measured at home and school, the authors were unable to assess the generalization of child effects across settings. According to most proximal measurements, children's academic performance and social skills also were improved—an effect that is perhaps not surprising, because trials included in these analyses often specifically targeted these aspects of impairment with specialized packages. If these effects could be corroborated with blinded ratings, this would be encouraging given the inconsistent evidence regarding the effects of medication on these outcomes.^{44,55}

When interpreting the present analysis, it is important to take some factors into account; all meta-analyses are constrained by the quality and diversity of the studies available for inclusion. First, for nearly all analyses, there was significant SMD heterogeneity among studies that may reflect the variety of intervention and outcome types analyzed. In most, although not all, cases, removing outliers identified using funnel plots decreased this heterogeneity while leaving treatment effects significant. Second, the authors could not establish how many parents had definite parenting or mental health problems, or how many children had clinical levels of conduct problems. Because participants were specifically selected for ADHD in childhood, it is likely that substantial numbers of individuals were unaffected by these additional problems. As a consequence, effect sizes seen for these outcomes, given the smaller room for improvement, are likely to be an underestimate of the true effects of behavioral interventions. Third,

Sonuga-Barke *et al.*¹⁹ raised questions about the status of the probably blinded measurements as valid outcomes. This is because they were often based on relatively small snapshots of behavior or a rating by a teacher who may not know the child well. However, in the present study, these measurements proved to be sensitive to treatment-related change. Fourth, this evaluation did not explore the impact on long-term outcomes, where one might predict a more robust impact of behavioral interventions. Fifth, although the authors established that parenting improved on rather artificial trial-based assessments, these may not have been able to capture improvements in everyday life situations. Sixth, there were insufficient studies to explore whether interventions specifically tailored for ADHD⁶⁴ were more effective than parenting interventions designed to treat more general childhood behavior problems or other aspects of impairment (social and academic skills). Seventh, the pool of trials meeting inclusion criteria was dominated by parenting interventions. This meant that the present analysis says little specifically about the value of child- or teacher-focused interventions. Eighth, in a related manner, no trials included measurements of changes in adult responses to children with ADHD other than parents. Ninth, the authors decided not to conduct or report an analysis of publication bias using techniques such as funnel plots because their interpretation is equivocal when based on a small number of studies, with difficulty in particular distinguishing between the effects of study heterogeneity and publication bias with sparse data.³⁶

In summary, although more evidence is required before behavioral interventions can be supported as a front-line treatment for core ADHD symptoms, the authors found evidence that they do have beneficial effects on parenting and parents' sense of empowerment and independently corroborated effects on conduct problems in children with ADHD. Initial evidence from proximal outcomes relating to academic achievement and social skills needs to be confirmed by probably blinded analyses and greater exploration is needed on the moderating impact of child age on intervention outcome.

<TXBX>Accepted June 3, 2014.

This article was reviewed under and accepted by Deputy Editor Stephen V. Faraone, PhD.

Dr. Daley is with the School of Medicine, University of Nottingham, UK, and the Centre for ADHD and Neurodevelopmental Disorders Across the Lifespan, Institute of Mental Health, University of Nottingham. Dr. van der Oord is with KU Leuven, Belgium, and the University of Amsterdam, the Netherlands. Dr. Ferrin is with Kings College London, Institute of Psychiatry, UK, and the Centro de Salud Mental de Estella, Navarra, Spain. Dr. Danckaerts is KU Leuven. Dr. Doepfner is with the University of Cologne, Germany. Dr. Cortese is with the Cambridgeshire and Peterborough Foundation Trust, and the School of Medicine and the Centre for ADHD and Neurodevelopmental Disorders Across the Lifespan, Institute of Mental Health, University of Nottingham. Dr. Sonuga-Barke is with the Developmental Brain-Behaviour Laboratory, School of Psychology, University of Southampton, UK; Ghent University, Belgium; and Aarhus University, Denmark.

This work was not directly funded. However support for meetings was received from Brain Products GmbH, Janssen-Cilag, Eli Lilly and Co., Medice, Shire, and Vifor. No honoraria were received, and funders had no input in the review and meta-analysis process or the writing of the article.

The European ADHD Guidelines Group is a workgroup of the European Network for Hyperkinetic Disorder (Eunethydis) and consists of the following members and associates co-opted to work on this review (listed in alphabetical order): T. Banaschewski, MD, PhD; J. Buitelaar, MD, PhD; D. Brandeis, PhD; D. Coghill, MD; S. Cortese, MD, PhD; D. Daley, PhD; M. Danckaerts, MD, PhD; R.W. Dittmann, MD, PhD; M. Döpfner, PhD; B. Falissard, MD, PhD; M. Ferrin, MD, PhD; J. Graham, MD; R. Hamilton, MD; C. Hollis, MD, PhD; M. Holtmann, MD, PhD; M. Hulpke-Wette, MD; M. Lecendreux, MD; E. Konofal, MD, PhD; P. Santosh, MD; E. Rosenthal, MD; A. Rothenberger, MD; J.A. Sergeant, PhD; E. Simonoff, MD; E.J. Sonuga-Barke, PhD; H.-Ch. Steinhausen, MD, PhD; J. Stevenson, PhD; A. Stringaris, MD, PhD; E. Taylor, MD; M. Thompson, MD; S. van der Oord, PhD; I. Wong, PhD; and A. Zuddas, MD.

The authors thank the following individuals who supplied information and advice: Howard Abikoff, PhD, New York University; Russell Barkley, PhD, Medical University of South Carolina; Bazian Ltd., UK; Steve Evans, PhD, Ohio University; Greg Fabiano, PhD, Buffalo State University; Philip Firestone, PhD, University of Ottawa, Canada; Nicholas Ialongo, PhD, Johns Hopkins University; Linda Pfiffner, PhD, University of California, San Francisco; Thomas Power, PhD, University of Pennsylvania; Mary V. Solanto, PhD, Mount Sinai School of Medicine; Jim Stevenson, PhD, University of Southampton, UK; Jim Swanson, PhD, University of California, Irvine; Barbara van den Hoofdakker, PhD, University of Groningen, Netherlands; and the anonymous reviewers who provided invaluable feedback. The authors also are grateful for the support given by Jana de Vos, MSc, and Stephanie Kristensen, MSc, of KU Leuven, and Jo Lockwood, BA, University of Nottingham.

Disclosure: Dr. Daley has provided educational talks for Eli Lilly and Co. and Shire, has attended an advisory board for Eli Lilly and Co., has received support for educational travel from Eli Lilly and Co., Shire, and HP Pharma, and currently holds funding from Shire. Dr. van der Oord has been involved in the development, implementation, and trialing of Braingame Brian, an executive functioning game training for children with ADHD, and Zelf Plannen, a cognitive behavioral planning intervention for adolescents with ADHD. She has no financial interests in either of these interventions. She has been a paid consultant for Janssen Pharmaceuticals, for the development and evaluation of the game Healseeker, which is aimed at training cognitive functions. Dr. Ferrin is a grant recipient for travel expenses from Alicia Koplowitz Foundation, Instituto de Salud Carlos III (ETS 07/90902, BAE 09/90088), and Gobierno de Navarra (Beca Jeronimo de Ayaz 2011/12). Dr. Danckaerts has served on the speaker's bench for Janssen-Cilag, Eli Lilly and Co., Shire, Novartis, and Medice. She has received funding for clinical trials from Eli Lilly and Co. and Shire and an educational grant from Shire. She has been involved in the development and dissemination of an ADHD Toolkit for teachers in primary school, distributed to all primary schools in Belgium by the Minister of Education, and has been a consultant for Janssen Pharmaceuticals for the development and evaluation of the game Healseeker, which is aimed at training cognitive functions. Dr. Döpfner has received research support from and served on the advisory or speaker's boards of Janssen-Cilag, Medice, Vifor, Shire, Eli Lilly and Co., and Novartis. He has been involved in the development, evaluation, and dissemination of the German

Therapieprogramm für Kinder mit hyperkinetischem und oppositionellem Problemverhalten (THOP) and Präventionsprogramm für Expansives Problemverhalten (PEP), which are behavioral interventions for children with ADHD. He is also head, supervisor, and lecturer at the School of Child Behavior Therapy at the University of Cologne. He has received royalties for treatment manuals (Beltz, Hogrefe Publisher) and as supervisor and lecturer and as consultant of the German Kassenärztliche Bundesvereinigung for the evaluation of behavior therapy. He has received support from Vifor Pharma for an ongoing trial on the effects of omega-3/6 fatty acids and from the German Research Foundation, German Ministry of Education and Research, Shire, Eli Lilly and Co. for trials on the effects of behavioral interventions for children with ADHD. Dr. Cortese has served as scientific consultant for Shire. He has received support to attend meetings from Eli Lilly and Co. and Shire. Dr. Sonuga-Barke has been involved in the development, implementation, and trialing of the New Forest Parenting Programme for preschool children with ADHD and has received royalties from sales of a New Forest Parent Training self-help manual. He has served on the speaker's board for Shire, UCB Pharma, Janssen-Cilag, and Medice. He has been a consultant for UCB Pharma and Shire. He has received research and conference support from Shire and has served on the advisory boards of Shire, Flynn Pharma, and AstraZeneca.

<COR>Correspondence to Edmund J.S. Sonuga-Barke, PhD, Academic Unit of Psychology, University of Southampton, Southampton SO17 1BJ, UK; e-mail: ejb3@soton.ac.uk. David Daley, PhD, Division of Psychiatry and Applied Psychology, School of Medicine, University of Nottingham, Nottingham, UK; e-mail: David.Daley@nottingham.ac.uk</TXBX>

References

1. Swanson JM, Sergeant JA, Taylor E, Sonuga-Barke EJS, Jensen PS. Attention-deficit hyperactivity disorder and hyperkinetic disorder. *Lancet*. 1998;351:429-433.
2. Biederman J, Monuteaux MC, Mick E *et al*. Young adult outcome of attention deficit hyperactivity disorder: a controlled 10-year follow-up study. *Psychol Med*. 2006;36:167-179.
3. Pelham WE, Foster EM, Robb JÁ. The economic impact of attention-deficit/hyperactivity disorder in children and adolescents. *J Pediatr Psychol*. 2007;32:711-727.
4. Taylor E, Döpfner M, Sergeant J *et al*. European clinical guidelines for hyperkinetic disorder: first upgrade. *Eur Child Adolesc Psychiatry*. 2004;13:I7-I30.
5. Zuvekas SH, Vitiello B. Stimulant medication use in children: a 12-year perspective. *Am J Psychiatry*. 2012;169:160-166.
6. Faraone SV, Biederman J, Spencer TJ, Aleardi M. Comparing the efficacy of medications for ADHD using meta-analysis. *Med Gen Med*. 2006;8:4.
7. van de Loo-Neus GH, Rommelse N, Buitelaar JK. To stop or not to stop? How long should medication treatment of attention deficit hyperactivity disorder be extended? *Eur Neuropsychopharmacol*. 2011;21:584-599.
8. Molina BS, Hinshaw SP, Swanson JM *et al*. MTA at 8 years: prospective follow-up of children treated for combined-type ADHD in a multisite study. *J Am Acad Child Adolesc Psychiatry*. 2009;48:484-500.

9. Van der Oord S, Prins PJM, Oosterlaan J, Emmelkamp PMG. Efficacy of methylphenidate, psychosocial treatments and their combination in school-aged children with ADHD: a meta-analysis. *Clin Psychol Rev.* 2008;28:783-800.
10. Graham J, Banaschewski T, Buitelaar J *et al.* European Guidelines Group: European guidelines on managing adverse effects of medication for ADHD. *Eur Child Adolesc Psychiatry.* 2011;20:17-37.
11. Adler LD, Nierenberg AA. Review of medication adherence in children and adults with ADHD. *Postgrad Med.* 2010;122:184-191.
12. Kovshoff H, Vrijens M, Thompson M *et al.* What influences clinicians' decisions about ADHD medication? Initial data from the Influences on Prescribing for ADHD Questionnaire (IPAQ). *Eur Child Adolesc Psychiatry.* 2013;22:533-542.
13. Fiks AG, Mayne S, DeBartolo E, Power TJ, Guevara JP. Parental preferences and goals regarding ADHD treatment. *Pediatrics.* 2013;132:692.
14. American Academy of Pediatrics. ADHD: clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics.* 2011;128:1007-1022.
15. Fabiano GA, Pelham WE Jr, Coles EK, Gnagy EM, Chronis-Tuscano A, O'Connor BC. A meta-analysis of behavioral treatments for attention-deficit/hyperactivity disorder. *Clin Psychol Rev.* 2009;29:129-140.
16. McGiloway S, Ni Mhaille G, Bywater T *et al.* A Parenting Intervention for childhood behavioral problems: a randomized controlled trial in disadvantaged community-based settings. *Consult Clin Psychol.* 2012;80:116-127.
17. Charach A, Carson P, Fox S, Ali M, Beckett J, Lim G. Interventions for preschool children at high risk for ADHD: a comparative effectiveness review. *Pediatrics.* 2013;131:E1584-E1604.
18. Day D, Altman G. Blinding in clinical trials and other studies. *BMJ.* 2000;321:504-504.
19. Sonuga-Barke E, Brandeis D, Cortese S *et al.* Non-pharmacological interventions for attention-deficit/hyperactivity disorder: systematic review and meta-analyses of randomised controlled trials of dietary and psychological treatments. *Am J Psychiatry.* 2013;170:275-289.
20. Polit DF, Gillespie BM, Griffin R. Deliberate ignorance: a systematic review of blinding in nursing clinical trials. *Nurs Res.* 2011;60:9-16.
21. Sonuga-Barke E, Brandeis D, Cortese S *et al.* Response to Chronis-Tuscano et al and Arns and Strehl A. *Am J Psychiatry.* 2013;170:800-802.
22. Chronis-Tuscano A, Chacko A, Barkley R. Key issues relevant to the efficacy of behavioral treatment for ADHD: a response to the Sonuga-Barke et al.'s meta-analysis of non-pharmacological treatments for ADHD. *Am J Psychiatry.* 2013;170:799.
23. Hutchings J, Bywater T, Daley D *et al.* A pragmatic randomised control trial of a parenting intervention in sure start services for children at risk of developing conduct disorder. *BMJ.* 2007;334:678-682.

24. Bor W, Sanders M, Markie-Dadds C. The effects of the Triple P-Positive Parenting Programme on preschool children with co-occurring disruptive behavior and attentional/hyperactive difficulties. *J Abnorm Child Psychol*. 2002;30:571-587.
25. Webster-Stratton CH, Reid MJ, Beauchaine T. Combining parent and child training for young children with ADHD. *J Clin Child Adolesc Psychol*. 2011;40:191-203.
26. Granic I, Patterson GR. Towards a comprehensive model of antisocial development: a dynamic systems approach. *Psychol Rev*. 2006;113:101-131.
27. Kazdin AF. Parent management training: evidence, outcomes and issues. *J Am Acad Psychiatry*. 1997;36:1349-1356.
28. Mash EJ, Barkley RA. *Treatment of Childhood Disorders*. 3rd ed. New York: Guilford; 2006.
29. O'Conner DF, Doerfler LA. ADHD with comorbid oppositional defiant disorder or conduct disorder: discrete or nondistinct disruptive behaviour disorders. *J Atten Disord*. 2008;12:126-134.
30. Yoshimasu K, Barbaresi WJ, Colligan RC, *et al*. Childhood ADHD is strongly associated with a broad range of psychiatric disorders during adolescence: a population based birth cohort study. *J Child Psychol Psychiatry*. 2012;53:1036-1043.
31. Johnston C. Parent characteristics and parent-child interactions in families of non-problem children and ADHD children with higher and lower levels of oppositional-defiant behavior. *J Abnorm Child Psychol*. 1996;24:85-104.
32. Deault LC. A systematic review of parenting in relation to the development of comorbidities and functional impairments in children with attention-deficit/hyperactivity disorder (ADHD). *Child Psychiatry Hum Dev*. 2010;41:168-192.
33. Pfiffner LJ, McBurnett K. Social skills training with parent generalization: treatment effects for children with attention deficit disorder. *J Consult Clin Psychol*. 1997;5:749-757.
34. Abikoff H, Gallagher R, Wells KC *et al*. Remediating organizational functioning in children with attention deficit hyperactivity disorder: immediate and long term effects from a randomised controlled trial. *J Consult Clin Psychol*. 2013;81:113.
35. Langberg JM, Epstein JN, Becker SP, Girio-Herrera E, Vaughn AJ. Evaluation of the homework, organization and planning (HOPS) intervention for middle school students with attention deficit hyperactivity disorder as implemented by school mental health providers. *Sch Psychol Q*. 2012;41:342-364.
36. Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*, Version 5.1.0. Cochrane Collaboration. <http://www.cochrane-handbook.org/>. Published March 2011.
37. Jadad AR, Moore RA, Carroll D *et al*. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials*. 1996;17:1-12.
38. RevMan, Version 5.1. Copenhagen: Nordic Cochrane Centre, Cochrane Collaboration; 2011.
39. Morris SB. Estimating effect sizes from pretest-posttest-control group designs. *Organ Res Methods*. 2008;11:364-386.

40. MTA Cooperative Group. A 14 month randomised clinical trial of treatment strategies for attention deficit/hyperactivity disorder. *Arch Gen Psychiatry*. 1999;56:1073-1086.
41. Wells KC, Epstein JN, Hinshaw SP *et al*. Parenting and family stress treatment outcomes in attention deficit/hyperactivity disorder (ADHD): an empirical analysis in the MTA study. *J Abnorm Child Psychol*. 2000;28:543-553.
42. Wells KC, Chi TC, Hinshaw SP *et al*. Treatment-related changes in objectively measured parenting behaviors in the multimodal treatment study of attention deficit hyperactivity disorder. *J Consult Clin Psychol*. 2006;74:649-657.
43. Antshell KM, Remer R. Social skills training in children with attention deficit/hyperactivity disorder: a randomised controlled clinical trial. *J Clin Child Adolesc Psychol*. 2003;32:153-165.
44. Bloomquist M, August G, Ostrander R. Effects of a school-based cognitive-behavioral intervention for ADHD children. *J Abnorm Child Psychol*. 1991;19:591-605.
45. Brown RT, Wynne ME, Borden KA, *et al*. Methylphenidate and cognitive therapy in children with attention deficit disorder: a double blind trial. *J Dev Behav Pediatr*. 1986;7:163-170.
46. Chacko A, Wymbs BT, Wymbs FA *et al*. Enhancing traditional parent training for single mothers of children with ADHD. *J Clin Child Adolesc Psychol*. 2009;38:206-218.
47. Evans SW, Schultz BK, DeMars CE, Davis H. Effectiveness of the Challenging Horizons afterschool programme for young adolescents with ADHD. *Behav Ther*. 2011;42:462-474.
48. Fabiano GA, Vujnovic RK, Pelham WE *et al*. Enhancing the effectiveness of special education programming for children with attention deficit hyperactivity disorder using a daily report card. *Sch Psychol Rev*. 2010;39:219-239.
49. Fabiano GA, Pelham WE, Cunningham CE *et al*. A waitlist controlled trial of behavioral parent training for fathers of children with ADHD. *J Clin Child Adolesc Psychol*. 2012;41:337-345.
50. Fehlings DL, Roberts W, Humphries T, Dawe G. Attention deficit/hyperactivity disorder: does cognitive behavioral therapy improve home behaviour? *J Dev Behav Pediatr*. 1991;2:222-228.
51. Hoath F, Sanders MR. A feasibility study of enhanced group Triple P-Positive Parenting Programme for parents of children with attention deficit/hyperactivity disorder. *Behav Change*. 2002;4:191-206.
52. Horn WF, Ialongo NS, Pascoe JM *et al*. Additive effects of psychostimulants, parent training, and self-control therapy with ADHD. *J Am Acad Child Adolesc Psychiatry*. 1991;30:233-234.
53. Jones K, Daley D, Hutchings J, Bywater T, Eames C. Efficacy of the Incredible Years basic parent training programme as an early intervention for children with conduct problems and ADHD. *Child Care Health Dev*. 2008;34:380-390.
54. Kapalka GM. Avoiding repetitions reduces ADHD children's management problems in the classroom. *Emot Behav Disord*. 2005;10:269-279.
55. Langberg JM, Epstein JN, Urbanowicz CM, Simon JO, Graham AJ. Efficacy of an organizational skill intervention to improve the academic functioning of students with attention deficit/hyperactivity disorder. *Sch Psychol Q*. 2008;23:407-417.

56. Mikami AY, Lerner MD, Griggs MS, McGrath A, Calhoun CD. Parental influence on children with attention deficit hyperactivity disorder: II results of a pilot intervention training parents as friendship coaches for children. *J Abnorm Child Psychol.* 2010;38:737-749.
57. Molina BSG, Flory K, Bukstein OG *et al.* Feasibility and preliminary efficacy of an afterschool programme for middle schoolers with ADHD: a randomised trial in a large public school. *J Atten Disord.* 2007;12:207-217.
58. Pfiffner LJ, Mikami AY, Hunag-Pollock C *et al.* Randomised controlled trial of integrated home-school behavioural treatment for ADHD, predominantly inattentive type. *J Am Acad Child Adolesc Psychiatry.* 2007;46:1041-1050.
59. Pisterman S, McGrath P, Firestone P *et al.* Outcome of parent mediated treatment of pre-schoolers with attention deficit with hyperactivity. *J Consult Clin Psychol.* 1989;57:628-635.
60. Pisterman S, Firestone P, McGrath P *et al.* The role of parent training in the treatment of preschoolers with ADDH. *Am J Orthopsychiatry.* 1992;62:397-408.
61. Power JT, Mautone JA, Soffer SL *et al.* A family-school intervention for children with ADHD: results of a randomised control clinical trial. *J Consult Clin Psychol.* 2012;80:611-623.
62. Sonuga-Barke EJS, Daley D, Thompson M, Laver-Bradbury C, Weeks A. Parent based therapies for preschool attention-deficit/hyperactivity disorder: a randomized, controlled trial with a community sample. *J Am Acad Child Adolesc Psychiatry.* 2001;40:402-408.
63. Sonuga-Barke EJS, Daley D, Thompson M, Laver-Bradbury C. Parent training for pre-school attention deficit/hyperactivity disorder: is it effective as part of routine primary care? *Br J Clin Psychol.* 2004;43:449-457.
64. Thompson MJ, Laver-Bradbury C, Ayres M *et al.* A small-scale randomized controlled trial of the revised New Forest Parenting Package for preschool children with attention deficit hyperactivity disorder. *Eur Child Adolesc Psychiatry.* 2009;18:605-616.
65. Tracey I, Tripp G. Stress management for attention deficit hyperactivity disorder. *Behav Ther.* 2005;36:223-233.
66. Van den Hoofdakker BJ, Van der Veen-Mulders L, Sytema S, Emmelkamp PMG, Minderaa RB, Nauta MH. Effectiveness of behavioral parent training for children with ADHD in routine clinical practice: a randomised controlled study. *J Am Acad Child Adolesc Psychiatry.* 2007;46:1263-1271.
67. Montoya A, Colom F, Ferrin M. Is psychoeducation for parents and teachers of children and adolescents with ADHD efficacious? A systematic literature review. *Eur Psychiatry.* 2011;26:166-175.
68. Goodman SH, Rouse MH, Connell AM *et al.* Maternal depression and child psychopathology: a meta-analytic review. *Clin Child Fam Psychol.* 2011;14:1-27.
69. Sonuga-Barke E, Halperin JM. Developmental phenotypes and causal pathways in attention deficit/hyperactivity disorder: potential targets for early intervention? *J Child Psychol Psychiatry.* 2010;51:368-389.

Figure 1 Flowchart showing the selection of trials. See Supplement 1 (available online) for specific reasons for exclusion.

Figure 2 Parental outcomes. Note: CI = confidence interval; MPROX = most proximal rating; MTA = National Institute of Mental Health Multimodal Treatment Study of Attention-Deficit/Hyperactivity Disorder; PBLIND = probably blinded rating; SMD = standardized mean difference.

Figure 3 Child outcomes. Note: ADHD = attention-deficit/hyperactivity disorder; CI = confidence interval; MPROX = most proximal rating; MTA = National Institute of Mental Health Multimodal Treatment Study of Attention-Deficit/Hyperactivity Disorder; PBLIND = probably blinded rating; SMD = standardized mean difference.

Table 1 Characteristics of Included Studies

Trial	Treatment		Control	Jadad Rating	Sample Size		Age Range (y)	Boys Medicated, % for ADHD, %	
	Delivery	Type			T	C			
Abikoff <i>et al.</i> (2012) ³⁴	parent and teacher	behavioral training	wait list	2	61	33	8-11	69	36
Antshel and Remer (2003) ⁴³	child and parent	social skills training	wait list	2	80	40	8-12	75	93
Bloomquist <i>et al.</i> (1991) ⁴⁴	child, parent, and teacher	CBT	wait list	2	20	16	8.58 mean	69	0 in analysis
Bor <i>et al.</i> (2002) ²⁴	parent	behavioral training	wait list	3	26	37	3.6 mean	73	0
Brown <i>et al.</i> (1986) ⁴⁵	child	self-control training	attention control	2	10	8	5-13	85	0
Chacko <i>et al.</i> (2009) ⁴⁶	parent	behavioral training	wait list	2	40	40	5-12	73	38.75
Evans <i>et al.</i> (2011) ⁴⁷	parent and child	behavioral and social skills training	TAU	1	31	18	11-13	71	57
Fabiano <i>et al.</i> (2010) ⁴⁸	children	daily report card	TAU	2	33	30	6-12	86	52
Fabiano <i>et al.</i> (2012) ⁴⁹	parent	behavioral training	wait list	2	27	28	6-12	87	54
Fehlings <i>et al.</i> (1991) ⁵⁰	parent and child	CBT	attention control	2	13	13	8-11	100	0

Hoath and Sanders (2002) ⁵¹	parent	behavioral training	wait list	1	9	11	5-9	76	70
Horn <i>et al.</i> (1991) ⁵²	parent and child	behavioral and self-control training	placebo	2	16	16	7-11	no info	0
Jones <i>et al.</i> (2008) ⁵³	parent	behavioral training	wait list	3	50	29	3.8 mean	68	0
Kapalka (2005) ⁵⁴	parent	behavioral training	wait list	0	45	41	5-10	100	no info
Langberg <i>et al.</i> (2008) ⁵⁵	child and parent	organizational skills training	wait list	1	24	13	grades 4-7	83	43
Langberg <i>et al.</i> (2012) ³⁵	child and parent	organizational skills training	wait list	2	23	24	grades 6-8	74	66
Mikami <i>et al.</i> (2011) ⁵⁶	parent	parent coaching on social skills	wait list	2	32	30	6-10	68	64.5
Molina <i>et al.</i> (2008) ⁵⁷	child and parent	organizational skills training	community	2	11	9	grades 6-8	75	31
MTA (1999, 2000, 2006) ⁴⁰⁻⁴²	child, parent, and teacher	Multicomponent : home, school, and camp	TAU	4	144	146	8.33 mean	80	47
Pfiffner and McBurnett (1997) ³³	child and parent	behavioral training and CBT	wait list	2	9	9	8-10	72	44
Pfiffner <i>et al.</i> (2007) ⁵⁸	child and parent	behavioral training and CBT	wait list	2	36	33	7-11	66	3
Pisterman <i>et al.</i> (1989) ⁵⁹	parent	behavioral training	wait list	3	23	23	4.1 mean	80	11
Pisterman <i>et al.</i> (1992) ⁶⁰	parent	behavioral training	wait list	3	23	22	4.1 mean	91	9
Power <i>et al.</i> (2012) ⁶¹	child and parent	behavioral and academic skills training	attention	2	100	99	grades 2-6	68	43
Sonuga-Barke <i>et al.</i> (2001) ⁶²	parent	behavioral training	counseling	3	30	28	2-4	62	0
Sonuga-Barke <i>et al.</i> (2004) ⁶³	parent	behavioral training	wait list	2	59	30	2-4	no info	0
Thompson <i>et al.</i> (2009) ⁶⁴	parent	behavioral training	wait list	4	21	20	2-6	73	0
Tracey and Tripp (2005) ⁶⁵	parent	stress management	wait list	2	20	20	6-15	88	88

Van den Hoofdakker <i>et al.</i> (2007) ⁶⁶	parent	behavioral training	TAU	2	48	48	4-12	76	40
Webster-Stratton <i>et al.</i> (2011) ²⁵	parent and child	behavioral training	wait list	3	49	50	6.4 mean	75	13

Note: ADHD = attention-deficit/hyperactivity disorder; C = control; CBT = cognitive behavioral therapy; grade = school year; MTA = National Institute of Mental Health Multimodal Treatment Study of Attention-Deficit/Hyperactivity Disorder; no info = no information provided in article; T = treatment; TAU = treatment as usual.

Table 2 Measurements Used in Each Trial for the Different Outcomes

		Child				Parent			
		ADHD	CP	SS	AS	MH	SC	PP	NP
Abikoff <i>et al.</i> ³⁴	MPROX				APRS ^a				COSS ^a
Antshel and Remer ⁴³	PBLIND								
	MPROX			SSRS ^a					
Bloomquist <i>et al.</i> ⁴⁴	PBLIND								
	MPROX	CTRS ^b		SSCSA ^a					
Bor <i>et al.</i> ²⁴	PBLIND								
	MPROX	ECBI ^a	ECBI ^a			DASS ^a	PSOC ^a		PS ^a
Brown <i>et al.</i> ⁴⁵	PBLIND		FOSR ^c						FOSR ^c
	MPROX	CPRS ^a	ACTeRS ^b	ACTeRS ^b	WRAT ^d				
Chacko <i>et al.</i> ⁴⁶	PBLIND	ACRS ^b							
	MPROX	DBD ^a	DBD ^a	IRS ^a		BDI ^a		DPICS ^c	DPICS ^c
Evans <i>et al.</i> ⁴⁷	PBLIND							DPICS ^c	DPICS ^c
	MPROX	ADHDRS ^a		IRS ^b	IRS ^b				
Fabiano <i>et al.</i> ⁴⁸	PBLIND								
	MPROX	DBD ^b	DBD ^b		APRS ^a				
Fabiano <i>et al.</i> ⁴⁹	PBLIND								
	MPROX		ECBI ^a					DPICS ^c	DPICS ^c
Fehlings <i>et al.</i> ⁵⁰	PBLIND							DPICS ^c	DPICS ^c
	MPROX	WWAS ^a							
Hoath and Sanders ⁵¹	PBLIND								
	MPROX	CAPS ^a	ECBI ^a			DASS ^a	PSBC ^a		PS ^a
Horn <i>et al.</i> ⁵²	PBLIND								
	MPROX	CAPS ^b	SESBI ^b						
Jones <i>et al.</i> ⁵³	PBLIND								
	MPROX	CPRS ^a							
	PBLIND								

Kapalka ⁵⁴	MPROX		SSQ ^b					
	PBLIND							
Langberg <i>et al.</i> ⁵⁵	MPROX				APRS ^a			
	PBLIND							
Langberg <i>et al.</i> ⁵⁵	MPROX	VADPRS ^a			HPC ^a			COSS ^a
	PBLIND							
Mikami <i>et al.</i> ⁵⁶	MPROX			SSRS ^a			PBIPC ^c	PBIPC ^c
	PBLIND						PBIPC ^c	PBIPC ^c
Molina <i>et al.</i> ⁵⁷	MPROX		ACPS ^a		SGR			
	PBLIND							
MTA ⁴⁰⁻⁴²	MPROX	SNAP ^a	SNAP ^a		WIAT ^d	BDI ^a	PCRQ ^a	PCRQ ^a
	PBLIND	classob ^c	classob ^c				OBS	
Pfiffner and McBurnett ³³	MPROX			SSRS ^a				
	PBLIND							
Pfiffner <i>et al.</i> ⁵⁸	MPROX			SSRS ^a				
	PBLIND							
Pisterman <i>et al.</i> ⁵⁹	MPROX		clinob ^c				clinob ^c	clinob ^c
	PBLIND		clinob ^c				clinob ^c	clinob ^c
Pisterman <i>et al.</i> ⁶⁰	MPROX	clinob ^c	clinob ^c				clinob ^c	clinob ^c
	PBLIND	clinob ^c	clinob ^c				clinob ^c	clinob ^c
Power <i>et al.</i> ⁶¹	MPROX	SNAP ^a	SNAP ^a		APRS ^b		PES ^a	PCRQ ^a
	PBLIND							
Sonuga-Barke <i>et al.</i> ⁶²	MPROX	PACS ^a	PACS ^a			GHQ ^a	PSOC ^a	
	PBLIND							
Sonuga-Barke <i>et al.</i> ⁶³	MPROX	homeob ^c	PACS			GHQ ^a	PSOC ^a	
	PBLIND							
Thompson <i>et al.</i> ⁶⁴	MPROX	PACS	PACS			GHQ ^a	GIPCI ^c	EE ^e
	PBLIND	homeob ^c	GIPCI ^c				GIPCI ^c	EE ^e
Tracey and Tripp ⁶⁵	MPROX					BDI ^a	PLOC ^a	PS ^a
	PBLIND							
Van den Hoofdakker <i>et al.</i> ⁶⁶	MPROX	CPRS ^a				PSI ^a	PSI ^a	
	PBLIND							
Webster-Stratton <i>et al.</i> ²⁵	MPROX	CPRS ^a	CPRS ^a	SCS ^a			PPI ^a	PPI ^a
	PBLIND							
		COCA-R					DPICS ^c	DPICS ^c

,
DPICS¹
7-OBS

Note: Superscript letter codes who provided the assessment; superscript number gives full name of test. ACPS = Aggression and Conduct Problem Scale; ACRS = Abbreviated Conners Rating Scale; ACTeRS = ACTeRS ADD-H Comprehensive Teachers Rating Scale; ADHD = attention-deficit/hyperactivity disorder; ADHDRS = Attention Deficit Hyperactivity Disorder Rating Scale; APRS = Academic Performance Rating Scale; AS = academic skills; BDI = Beck Depression Inventory; CAPS = Child Attention Problem Rating Scale; classob = classroom observation; clinob = clinic observation; COCA-R = Coder Observation of Classroom Adaption-Revised: Cognitive Concentration; COSS = Children's Organizational Skills Scale; CP = conduct problems; CPRS = Conners Parent Rating Scale-Hyperactivity; CTRS = Conners Teachers Rating Scale; DASS = Depression Anxiety and Stress Scale; DBD = Disruptive Disorders Rating Scale; DPICS = Dyadic Parent-Child Interaction Coding System; ECBI = Eyberg Child Behavior Inventory; EE = Expressed Emotion measured using the Pre-school Five Minute Speech Sample; FOSR = Family Observation Schedule; GHQ = General Health Questionnaire; GIPCI = Global Impressions of Parent Child Interaction Revised; homeob = home observation of on-task behavior; HPC = Homework Problem Checklist; IRS = Impairment Rating Scale; MH = mental health; MPROX = most proximal rating; MTA = National Institute of Mental Health Multimodal Treatment Study of Attention-Deficit/Hyperactivity Disorder; NP = negative parenting; OBS = observed constructive parenting; PACS = Parent Account of Childhood Symptoms Interview; PBIPC = Parental Behavior in Parent-Child Interaction; PBLIND = probably blinded rating; PCRQ = Parent Child Relationship Questionnaire; PES = Parent as Educator; PLOC = parents locus of control; PP = positive parenting; PPI = ??; PS = Parenting Scale; PSBC = Problem Setting and Behavior Checklist; PSI = Parenting Stress Inventory; PSOC = Parenting Sense of Competence Scale; SC = self-concept; SCS = Social Competence Scale; SESBI = Sutter-Eyberg Student Behavior Inventory Revised; SGR = school grade records; SNAP = Pelham Swanson and Nolan Questionnaire; SS = social skills; SSCSA = Walker-McConnel Scale of Social Competence and School Adjustment; SSQ = Social Situations Questionnaire; SSRS = Social Skills Rating Scale; VADPRS = Vanderbilt ADHD Diagnostic Parent Rating Scale; WIAT = Wechsler Individual Achievement Test; WRAT = Wide Range Achievement Test; WWAS = Werry Weiss Activity Scale.

^aParent.

^bTeacher.

^cObservation.

^dAcademic skills test.

^eSpeech sample.

Supplement 1 Information on Excluded Studies

Study	Reasons for Exclusion
Abikoff and Gittelman (1985) ¹	not a behavioral intervention; study explores cognitive training
Abikoff <i>et al.</i> (2004) ²	no appropriate control group; behavioral intervention adjunctive to medication
Abikoff <i>et al.</i> (2004) ³	no appropriate control group; behavioral intervention adjunctive to medication
Altemeier and Horwitz (1997) ⁴	not randomized
Anastopoulos <i>et al.</i> (1993) ⁵	not randomized
Arnold <i>et al.</i> (2003) ⁶	MTA study that did not add additional outcomes to the included studies
Barkley <i>et al.</i> (1992) ⁷	no appropriate control group; this study is a comparison of 3 active therapies
Barkley <i>et al.</i> (1996) ⁸	reports violation of randomization
Barkley <i>et al.</i> (2000) ⁹	reports violation of randomization in 8 children
Barkley <i>et al.</i> (2001) ¹⁰	no appropriate control group; this study compares 3 different combinations of intervention but does not have a control condition
Carlson <i>et al.</i> (1992) ¹¹	not randomized
Chacko <i>et al.</i> (2003) ¹²	not a behavioral intervention; this is a medication study
Christensen and Sprague (1973) ¹³	not randomized
Christensen (1975) ¹⁴	no appropriate control group; this is a within-subject design
Cohen <i>et al.</i> (1981) ¹⁵	not randomized

- Cunningham *et al.* (1995)¹⁶
 Döpfner *et al.* (2004)¹⁷
 Dubey and O'Leary (1983)¹⁸
 Epstein *et al.* (2007)¹⁹
 Evans *et al.* (2005)²⁰
 Fabiano *et al.* (2009)²¹
 Firestone *et al.* (1981)²²
 Firestone *et al.* (1986)²³
 Frankel *et al.* (1997)²⁴
 Gerber-von Müller *et al.* (2009)²⁵
 Gonzalez and Sellers (2002)²⁶
 Hanisch *et al.* (2010)²⁷
 Hantson *et al.* (2012)²⁸
 Hechtman *et al.* (2004)²⁹
 Hechtman *et al.* (2004)³⁰
 Hinshaw *et al.* (1984)³¹
 Hinshaw *et al.* (1984)³²
 Hinshaw *et al.* (2000)³³
 Horn *et al.* (1987)³⁴
 Horn *et al.* (1990)³⁵
 Hupp *et al.* (2002)³⁶
 Iolango *et al.* (1993)³⁷
 Jensen *et al.* (2001)³⁸
 Jensen *et al.* (2004)³⁹
 Jensen *et al.* (2007)⁴⁰
 Kapalka (2004)⁴¹
 Kapalka (2005)⁴²
 Kern *et al.* (2007)⁴³
 Kienle *et al.* (2009)⁴⁴
 Klein and Abikoff (1997)⁴⁵
 Klein *et al.* (2004)⁴⁶
 Lerner *et al.* (2011)⁴⁷
 Lloyd *et al.* (2010)⁴⁸
 McGrath *et al.* (2011)⁴⁹
 McNeil *et al.* (1991)⁵⁰
 Meyer and Kelley (2007)⁵¹
 Mikami *et al.* (2013)⁵²
 Miranda *et al.* (2002)⁵³
 Odom (1996)⁵⁴
 Osterberg and Rydell (2012)⁵⁵
 O'Leary *et al.* (1976)⁵⁶
 Papazian *et al.* (2009)⁵⁷
 Pariseau *et al.* (2010)⁵⁸
 Pelham (1977)⁵⁹
 Pelham *et al.* (1993)⁶⁰
 Pelham (1999)⁶¹
 Pelham and Gnagy (1999)⁶²
 Pelham *et al.* (2000)⁶³
- not a specific ADHD sample
 no appropriate control group; this study applies an adaptive and individually tailored approach to treatment
 randomization unclear; groups were assigned so that age and gender were evenly distributed
 not a behavioral intervention
 not randomized
 no appropriate control group; this study is a comparison of 2 behavioral interventions
 no control group
 no control group and follow-up of a previously excluded study
 children with and without ADHD included in sample
 no appropriate treatment control group
 ADHD status of children unclear
 not a specific ADHD sample
 not randomized
 no appropriate control group; behavioral intervention adjunctive to medication
 no appropriate control group; behavioral intervention adjunctive to medication
 no appropriate control group; behavioral intervention adjunctive to medication
 no appropriate control group; behavioral intervention adjunctive to medication
 MTA study that did not add any additional outcomes
 no appropriate control group; this study is a comparison of 2 behavioral interventions
 no appropriate control group; this study is a comparison of 2 behavioral interventions
 case study
 same dataset as Horn *et al.*,³⁵ which was excluded
 MTA study that did not add additional outcomes
 MTA study that did not add additional outcomes
 MTA study that did not add additional outcomes
 not a behavioral intervention
 weak randomization procedure
 no appropriate control group; this study is a comparison of 2 behavioral interventions
 not randomized
 no appropriate control group; this study explores combinations of medication and behavioral intervention
 methodology article on design of study by Hechtman^{29,30}; no control group
 randomization unclear
 not a behavioral intervention
 study met criteria for inclusion, but data were not amenable to analysis, and the investigators were unable to provide the data
 randomization unclear
 no appropriate control; this study compares parent with self-monitoring for homework problems
 no appropriate control; this study compares 2 behavioral interventions
 randomization unclear
 randomization unclear
 randomization unclear; not an entirely ADHD sample
 insufficient detail in summary statistics to allow calculation of SMD
 not a behavioral intervention
 no appropriate control group; this is a within-subject design
 case study
 no appropriate control group; this is a within-subject design
 review article; no data
 review article; no data
 no control group; same sample as MTA comparing behavioral and combined groups

Pfeiffer <i>et al.</i> (2008) ⁶⁴	not a specific ADHD sample and not a behavioral intervention
Pfeiffer <i>et al.</i> (2007) ⁶⁵	not fully randomized
Poulsen <i>et al.</i> (2010) ⁶⁶	not fully randomized
Presentación Herrero <i>et al.</i> (2010) ⁶⁷	not randomized
Rapport <i>et al.</i> (1982) ⁶⁸	case study (2)
Rieppi <i>et al.</i> (2002) ⁶⁹	MTA study that did not add additional outcomes
Rosén <i>et al.</i> (1984) ⁷⁰	not randomized
Rutter and Sroufe (2000) ⁷¹	review article; no data
Sanders <i>et al.</i> (2007) ⁷²	not a specific ADHD sample
Sayal <i>et al.</i> (2010) ⁷³	not a behavioral intervention
Shafro and Sulzbacher (1977) ⁷⁴	case study
Schumann <i>et al.</i> (1998) ⁷⁵	not a specific ADHD sample
Scott <i>et al.</i> (2010) ⁷⁶	not a specific ADHD sample
Shaffer <i>et al.</i> (2001) ⁷⁷	not a behavioral intervention
So <i>et al.</i> (2008) ⁷⁸	no appropriate control group; behavioral intervention adjunctive to medication
Springer and Reddy (2010) ⁷⁹	not randomized
Strayhorn and Weidman (1989) ⁸⁰	not a specific ADHD sample
Strayhorn and Bickel (2002) ⁸¹	no appropriate control group; comparison of 2 behavioral approaches
Storebø <i>et al.</i> (2012) ⁸²	behavioral intervention adjunctive to medication
Swanson <i>et al.</i> (2001) ⁸³	reanalysis of MTA study; no new outcomes added
Thorell (2009) ⁸⁴	not adequately randomized; allocation to condition influenced by other factors
Thurston (1979) ⁸⁵	randomization unclear
Tutty <i>et al.</i> (2003) ⁸⁶	no appropriate control group; behavioral intervention adjunctive to medication
Van der Oord <i>et al.</i> (2007) ⁸⁷	no appropriate control group; behavioral intervention adjunctive to medication
Waxmonsky <i>et al.</i> (2008) ⁸⁸	no appropriate control group; behavioral intervention adjunctive to medication
Waxmonsky <i>et al.</i> (2010) ⁸⁹	no appropriate control group; behavioral intervention adjunctive to medication
Wolraich <i>et al.</i> (1978) ⁹⁰	no appropriate control group
Wulbert and Dries (1977) ⁹¹	case study

Note: ADHD = attention-deficit/hyperactivity disorder; MTA = National Institute of Mental Health Multimodal Treatment Study of Attention-Deficit/Hyperactivity Disorder; SMD = standardized mean difference.

Supplementary References

1. Abikoff H, Gittelman R. Hyperactive children treated with stimulants: is cognitive training a useful adjunct? *Arch Gen Psychiatry*. 1985;42:953-961.
2. Abikoff H, Hechtman L, Klein RG *et al.* Symptomatic improvement in children with ADHD treated with long-term methylphenidate and multimodal psychosocial treatment. *J Am Acad Child Adolesc Psychiatry*. 2004;43:802-811.
3. Abikoff H, Hechtman L, Klein RG *et al.* Social functioning in children with ADHD treated with long-term methylphenidate and multimodal psychosocial treatment. *J Am Acad Child Adolesc Psychiatry*. 2004;43:820-829.
4. Altemeier WA, Horwitz E. The role of the school in the management of attention deficit hyperactivity disorder. *Pediatr Ann*. 1997;26:737-744.

5. Anastopoulos AD, Shelton TL, DuPaul GJ, Guevremont DC. Parent training for attention-deficit hyperactivity disorder: its impact on parent functioning. *J Abnorm Child Psychol*. 1993;21:581-596.
6. Arnold LM, Elliott M, Sachs L *et al*. Effects of ethnicity on treatment attendance, stimulant response/dose, and 14-month outcome in ADHD. *J Consult Clin Psychol*. 2003;71:713-727.
7. Barkley RA, Guevremont DC, Anastopoulos AD, Fletcher KF. A comparison of three family therapy programs for treating family conflicts in adolescents with attention-deficit hyperactivity disorder. *J Consult Clin Psychol*. 1992;60:450-462.
8. Barkley RA, Terri L, Shelton TL *et al*. Preliminary findings of an early intervention programme with aggressive hyperactive children. *Ann N Y Acad Sci*. 2003;794:277-287.
9. Barkley RA, Terri L, Shelton TL *et al*. Multi-method psycho-educational intervention for pre-school children with disruptive behaviour: preliminary results at post treatment. *J Child Psychol Psychiatry*. 2000;41:319-332.
10. Barkley RA, Edwards G, Laneri M, Fletcher K, Metevia L. The efficacy of problem-solving communication training alone, behavior management training alone, and their combination for parent-adolescent conflict in teenagers with ADHD and ODD. *J Consult Clin Psychol*. 2001;69:926-941.
11. Carlson CL, Pelham WE Jr, Milich R, Dixon J. Single and combined effects of methylphenidate and behavior therapy on the classroom performance of children with attention-deficit hyperactivity disorder. *J Abnorm Child Psychol*. 1992;20:213-32.
12. Chacko A, Pelham WE, Gnagy EM *et al*. Stimulant medication effects in a summer treatment program among young children with attention-deficit/hyperactivity disorder. *J Am Acad Child Psychiatry*. 2005;44:249-257.
13. Christensen DE, Sprague RL. Reduction of hyperactive behavior by conditioning procedures alone and combined with methylphenidate (Ritalin). *Behav Res Ther*. 1973;11:331-334.
14. Christensen DE. Effects of combining methylphenidate and a classroom token system in modifying hyperactive behavior. *Am J Ment Defic*. 1975;80:266-276.
15. Cohen NJ, Sullivan J, Minde K, Novak C, Helwig C. Evaluation of the relative effectiveness of methylphenidate and cognitive behavior modification in the treatment of kindergarten-aged hyperactive children. *J Abnorm Child Psychol*. 1981;9:43-54.
16. Cunningham CE, Bremner R, Boyle R. Large group community based parenting programmes for families of preschoolers at risk for disruptive behaviour disorders: utilization, cost effectiveness and outcomes. *J Child Psychol Psychiatry*. 1995;36:1141-1159.
17. Döpfner M, Breuer D, Schürmann S, Metternich TW, Rademacher C, Lehmkuhl G. Effectiveness of an adaptive multimodal treatment in children with attention-deficit hyperactivity disorder—global outcome. *Eur Child Adolesc Psychiatry*. 2004;13(suppl 1):I117-I129.
18. Dubbey DR, O'Leary SG. Training parents of hyperactive children in child management: a comparative outcome study. *J Abnorm Child Psychol*. 1983;11:229-246.


19. Epstein JN, Rabiner D, Johnson DE *et al.* Improving attention-deficit/hyperactivity disorder treatment outcomes through use of a collaborative consultation treatment service by community-based pediatricians: a cluster randomized trial. *Arch Pediatr Adolesc Med.* 2007;161:835-840.
20. Evans SW, Langberg J, Raggi V, Allen J, Buvinger EC. Development of a school-based treatment program for middle school youth with ADHD. *J Atten Disord.* 2005;9:343-353.
21. Fabiano GA, Chacko A, Pelham WE Jr *et al.* A comparison of behavioral parent training programs for fathers of children with attention-deficit/hyperactivity disorder. *Behav Ther.* 2009;40:190-204.
22. Firestone P, Kelly MJ, Goodman JT, Davey J. Differential effects of parent training and stimulant medication with hyperactives: a progress report. *J Am Acad Child Psychiatry.* 1981;20:135-147.
23. Firestone P, Crowe D, Goodman JT, McGrath P. Vicissitudes of follow-up studies: differential effects of parent training and stimulant medication with hyperactives. *Am J Orthopsychiatry.* 1986;5:184-194.
24. Frankel F, Myatt R, Cantwell DP, Feinberg DT. Parent-assisted transfer of children's social skills training: effects on children with and without attention-deficit hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry.* 1997;36:1056-1064.
25. Gerber-von Müller G, Petermann U, Petermann F *et al.* ADHD summer camp: development and evaluation of a multimodal intervention program. *Kindheit Entwicklung.* 2009;18:162-172.
26. Gonzalez LO, Sellers EW. The effects of a stress-management program on self-concept, locus of control, and the acquisition of coping skills in school-age children diagnosed with attention deficit hyperactivity disorder. *J Child Adolesc Psychiatr Nurs.* 2002;15:5-15.
27. Hanisch C, Freund-Braier I, Hautmann C. Detecting effects of the indicated prevention Programme for Externalizing Problem behaviour (PEP) on child symptoms, parenting, and parental quality of life in a randomized controlled trial. *Behav Cogn Psychother.* 2010;38:95-112.
28. Hantson J, Wang PP, Grizenko-Vida M *et al.* Effectiveness of a therapeutic summer camp for children with ADHD: phase i clinical intervention trial. *J Atten Disord.* 2012;16:610-617.
29. Hechtman L, Abikoff H, Klein RG *et al.* Academic achievement and emotional status of children with ADHD treated with long-term methylphenidate and multimodal psychosocial treatment. *J Am Acad Child Adolesc Psychiatry.* 2004;43:812-819.
30. Hechtman L, Abikoff H, Klein RG *et al.* Children with ADHD treated with long-term methylphenidate and multimodal psychosocial treatment: impact on parental practices. *J Am Acad Child Adolesc Psychiatry.* 2004;43:830-838.
31. Hinshaw SP, Henker B, Whalen CK. Self-control in hyperactive boys in anger-inducing situations: effects of cognitive-behavioral training and of methylphenidate. *J Abnorm Child Psychol.* 1984;12:55-77.
32. Hinshaw SP, Henker B, Whalen CK. Cognitive-behavioral and pharmacologic interventions for hyperactive boys: comparative and combined effects. *J Consult Clin Psychol.* 1984;52:739-749.
33. Hinshaw SP, Owens EB, Wells KC. Family processes and treatment outcome in the MTA: negative/ineffective parenting practices in relation to multimodal treatment. *J Abnorm Child Psychol.* 2000;28:555-568.


34. Horn WF, Ialongo N, Popovich S, Peradotto D. Behavioral parent training and cognitive-behavioral self-control therapy with ADD-H children: comparative and combined effects. *J Abnorm Child Psychol.* 1987;16:57-68.
35. Horn WF, Ialongo NS, Greenberg G, Packard T, Smith-Winberry C. Additive effects of behavioral parent training and self-control therapy with attention deficit hyperactivity disorder children. *J Clin Child Psychol.* 1990;19:98-110.
36. Hupp SD, Reitman D, Northup J, O'Callaghan P, LeBlanc M. The effects of delayed rewards, tokens, and stimulant medication on sportsmanlike behavior with ADHD-diagnosed children. *Behav Modif.* 2002;26:148-162.
37. Ialongo NS, Horn WF, Pascoe JM *et al.* The effects of a multimodal intervention with attention-deficit hyperactivity disorder children: a 9-month follow-up. *J Am Acad Child Adolesc Psychiatry.* 1993;32:182-189.
38. Jensen PS, Hinshaw SP, Swanson JM *et al.* Findings from the NIMH Multimodal Treatment Study of ADHD (MTA): implications and applications for primary care providers. *J Dev Behav Pediatr.* 2001;22:60-73.
39. Jensen PS, Eaton Hoagwood K, Roper M *et al.* The services for children and adolescents-parent interview: development and performance characteristics. *J Am Acad Child Adolesc Psychiatry.* 2004;43:1334-1344.
40. Jensen PS, Arnold LE, Swanson JM *et al.* 3-Year follow-up of the NIMH MTA study. *J Am Acad Child Adolesc Psychiatry.* 2007;46:989-1002.
41. Kapalka GM. Longer eye contact improves ADHD children's compliance with parents' commands. *J Atten Disord.* 2004;8:17-23.
42. Kapalka GM. Avoiding repetitions reduces ADHD children's management problems in the classroom. *Emot Behav Disord.* 2005;10:269-279.
43. Kern L, DuPaul GJ, Volpe RJ *et al.* Multisetting assessment-based intervention for young children at risk for attention deficit hyperactivity disorder: initial effects on academic and behavioral functioning. *Sch Psychol Rev.* 2007;36:237-255.
44. Kienle X, Koerber S, Karch D. [Effectiveness of parent training for children with ADHD in routine clinical practice]. *Prax Kinderpsychol Kinderpsychiatr.* 2009;58:16-33.
45. Klein RG, Abikoff H. Behavior therapy and methylphenidate in treatment of children with ADHD. *J Atten Disord.* 1997;2:89-114.
46. Klein RG, Abikoff H, Hechtman L, Weiss G. Design and rationale of controlled study of long-term methylphenidate and multimodal psychosocial treatment in children with ADHD. *J Am Acad Child Adolesc Psychiatry.* 2004;43:792-801.
47. Lerner MD, Mikami AY, McLeod BD. The alliance in a friendship coaching intervention for parents of children with ADHD. *Behav Ther.* 2011;42:449-461.
48. Lloyd A, Brett D, Wesnes K. Coherence training in children with attention-deficit hyperactivity disorder: cognitive functions and behavioural changes. *Altern Ther Health Med.* 2010;16:34-42.

49. McGrath PJ, Lingley-Pottie P, Thurston C *et al.* Telephone-based mental health interventions for child disruptive behaviour or anxiety disorder: randomised trials and overall analysis. *J Am Acad Child Adolesc Psychiatry.* 2011;50:1162-1172.
50. McNeil CB, Eyberg S, Eisenstadt TH, Newcomb K, Funderburk B. Parent-child interaction therapy with behaviour problem children: generalization of treatment effects to the school setting. *J Clin Child Psychol.* 1991;20:140-151.
51. Meyer K, Kelley ML. Improving homework in adolescents with attention-deficit/hyperactivity disorder: self vs. parent monitoring of homework behavior and study skills. *Child Fam Behav Ther.* 2007;29:25-42.
52. Mikami AY, Griggs MS, Lerner MD *et al.* A randomized trial of a classroom intervention to increase peers' social inclusion of children with attention-deficit/hyperactivity disorder. *J Consult Clin Psychol.* 2013;81:100.
53. Miranda A, Presentación MJ, Soriano M. Effectiveness of a school-based multicomponent program for the treatment of children with ADHD. *J Learn Disabil.* 2002;35:546-562.
54. Odom SE. Effects of an educational intervention on mothers of male children with attention deficit hyperactivity disorder. *J Community Health Nurs.* 1996;13:207-220.
55. Ostberg M, Rydell AM. An efficacy study of a combined parent and teacher management training programme for children with ADHD. *Nord J Psychiatry.* 2012;66:123-130.
56. O'Leary KD, Pelham WE, Rosenbaum A, Price GH. Behavioral treatment of hyperkinetic children. An experimental evaluation of its usefulness. *Clin Pediatr.* 1976;15:510-515.
57. Papazian O, Alfonso I, Luzondo RJ, Araguez N. [Training of executive function in preschool children with combined attention deficit hyperactivity disorder: a prospective, controlled and randomized trial]. *Rev Neurol.* 2009;48(suppl 2):S119-S122.
58. Pariseau ME, Fabiano G, Massetti GM, Hart KC, Pelham WE. Extended time on academic assignments: Does increased time lead to improved performance for children with ADHD? *Sch Psychol Q.* 2010;25:236-248.
59. Pelham WE. Withdrawal of a stimulant drug and concurrent behavioral intervention in the treatment of a hyperactive child. *Behav Ther.* 1977;8:473-479.
60. Pelham WE Jr, Carlson C, Sams SE, Vallano G, Dixon MJ, Hoza B. Separate and combined effects of methylphenidate and behavior modification on boys with attention deficit-hyperactivity disorder in the classroom. *J Consult Clin Psychol.* 1993;61:506-515.
61. Pelham WE Jr. The NIMH multimodal treatment study for attention-deficit hyperactivity disorder: just say yes to drugs alone? *Can J Psychiatry.* 1999;44:981-990.
62. Pelham WE, Gnagy EM. Psychosocial and combined treatments for ADHD. *Ment Retard Dev Disabil Res Rev.* 1999;5:225-236.
63. Pelham WE, Gnagy EM, Greiner AR *et al.* Behavioral versus behavioral and pharmacological treatment in ADHD children attending a summer treatment program. *J Abnorm Child Psychol.* 2000;28:507-525.


64. Pfeiffer B, Henry A, Miller S, Witherell S. Effectiveness of Disc 'O' Sit cushions on attention to task in second-grade students with attention difficulties. *Am J Occup Ther.* 2008;62:274-281.
65. Pfiffner LJ, Yee Mikami A, Huang-Pollock C, Easterlin B, Zalecki C, McBurnett K. A randomized, controlled trial of integrated home-school behavioral treatment for ADHD, predominantly inattentive type. *J Am Acad Child Adolesc Psychiatry.* 2007;46:1041-1050.
66. Poulsen AA, Horswill MS, Wetton MA, Hill A, Lim SM. A brief office-based hazard perception intervention for drivers with ADHD symptoms. *Aust N Z J Psychiatry.* 2010;44:528-534.
67. Presentación Herrero MJ, Siegenthaler Hierro R, Jara Jiménez P, Miranda Casas A. Psychosocial intervention follow-up in children with ADHD: effects on academic, emotional and social functioning. *Psicothema.* 2010;22:778-783.
68. Rapport MD, Murphy HA, Bailey JS. Ritalin vs. response cost in the control of hyperactive children: a within-subject comparison. *J Appl Behav Anal.* 1982;15:205-216.
69. Rieppi R, Greenhill LL, Ford RE *et al.* Socioeconomic status as a moderator of ADHD treatment outcomes. *J Am Acad Child Adolesc Psychiatry.* 2002;41:269-277.
70. Rosén LA, O'Leary SG, Joyce SA, Conway G, Pfiffner LJ. The importance of prudent negative consequences for maintaining the appropriate behavior of students. *J Abnorm Child Psychol.* 1984;12:581-604.
71. Rutter M, Sroufe A. Developmental psychopathology: concepts and challenges. *Dev Psychopathol.* 2000;12:265-296.
72. Sanders MR, Bor W, Morawska A. Maintenance of treatment gains: a comparison of enhanced, standard, and self-directed Triple P-Positive Parenting Program. *J Abnorm Child Psychol.* 2007;35:983-998.
73. Sayal K, Owen V, White K, Merrell C, Tymms P, Taylor E. Impact of early school-based screening and intervention programs for ADHD on children's outcomes and access to services: follow-up of a school-based trial at age 10 years. *Arch Pediatr Adolesc Med.* 2010;164:462-469.
74. Shafto F, Sulzbacher S. Comparing treatment tactics with a hyperactive preschool child: stimulant medication and programmed teacher intervention. *J Appl Behav Anal.* 1977;10:13-20.
75. Schumann EM, Foote RC, Eyberg SM, Boggs SR, Algina J. Efficacy of parent-child interaction therapy: interim report of a randomized trial with short-term maintenance. *J Clin Child Psychol.* 1998;27:34-45.
76. Scott S, Sylva K, Doolan M *et al.* Randomised controlled trial of parent groups for child antisocial behaviour targeting multiple risk factors: the SPOKES project. *J Child Psychol Psychiatry.* 2010;51:48-57.
77. Shaffer RJ, Jacokes LE, Cassily JF, Greenspan SI, Tuchman RF, Stemmer PJ Jr. Effect of interactive metronome training on children with ADHD. *Am J Occup Ther.* 2001;5:155-162.
78. So CY, Leung PW, Hung SF. Treatment effectiveness of combined medication/behavioural treatment with Chinese ADHD children in routine practice. *Behav Res Ther.* 2008;46:983-992.
79. Springer C, Reddy LR. Measuring parental treatment adherence in a multimodal treatment program for children with ADHD: a preliminary investigation. *Child Fam Behav Ther.* 2010;32:272-290.

80. Strayhorn JM, Weidman CS. Reduction of attention deficit and internalizing symptoms in preschoolers through parent-child interaction training. *J Am Acad Child Adolesc Psychiatry*. 1989;28:888-896.
81. Strayhorn JM Jr, Bickel DD. Reduction in children's symptoms of attention deficit hyperactivity disorder and oppositional defiant disorder during individual tutoring as compared with classroom instruction. *Psychol Rep*. 2002;91:69-80.
82. Storebø OJ, Gluud C, Winkel P, Simonsen E. Social-skills and parental training plus standard treatment versus standard treatment for children with ADHD—the randomised SOSTRA trial. *PLoS One*. 2012;7:e37280.
83. Swanson JM, Kraemer HC, Hinshaw SP *et al*. Clinical relevance of the primary findings of the MTA: success rates based on severity of ADHD and ODD symptoms at the end of treatment. *J Am Acad Child Adolesc Psychiatry*. 2001;40:168-179.
84. Thorell LB. The Community Parent Education Program (COPE): treatment effects in a clinical and a community-based sample. *Clin Child Psychol Psychiatry*. 2009;14:373-387.
85. Thurston LP. Comparison of the effects of parent training and of Ritalin in treating hyperactive children. *Int J Ment Health*. 1979;8:121-128.
86. Tutty S, Gephart H, Wurzbacher K. Enhancing behavioral and social skill functioning in children newly diagnosed with attention-deficit hyperactivity disorder in a pediatric setting. *J Dev Behav Pediatr*. 2003;24:51-57.
87. van der Oord S, Prins PJ, Oosterlaan J, Emmelkamp PM. Does brief, clinically based, intensive multimodal behavior therapy enhance the effects of methylphenidate in children with ADHD? *Eur Child Adolesc Psychiatry*. 2007;16:48-57.
88. Waxmonsky J, Pelham WE, Gnagy E *et al*. The efficacy and tolerability of methylphenidate and behavior modification in children with attention-deficit/hyperactivity disorder and severe mood dysregulation. *J Child Adolesc Psychopharmacol*. 2008;18:573-588.
89. Waxmonsky JG, Waschbusch DA, Pelham WE, Draganac-Cardona L, Rotella B, Ryan L. Effects of atomoxetine with and without behavior therapy on the school and home functioning of children with attention-deficit/hyperactivity disorder. *J Clin Psychiatry*. 2010;71:1535-1551.
90. Wolraich M, Drummond T, Salomon MK, O'Brien ML, Sivage C. Effects of methylphenidate alone and in combination with behavior modification procedures on the behavior and academic performance of hyperactive children. *J Abnorm Child Psychol*. 1978;6:149-161.
91. Wulbert M, Dries R. The relative efficacy of methylphenidate (Ritalin) and behavior-modification techniques in the treatment of a hyperactive child. *J Appl Behav Anal*. 1977;10:21-31.







Overall SMD=0.68;95%CI=0.27-1.09




Overall SMD=0.63;95%CI=0.47-0.7




Overall SMD=0.57;95%CI=0.37-0.78




Overall SMD=0.43;95%CI=0.24-0.62




Overall SMD=0.37;95%CI=0.03-0.7




Overall SMS=0.09;95%CI=0.06-0.23




Overall SMD=0.35;95%CI=0.19–0.50




Overall SMD=0.02;95%CI=-0.30-0.34




Overall SMD=0.26;95%CI=0.14-0.37



Overall SMD=0.31;95%CI=0.05–0.57



Overall SMD=0.47;95%CI=0.15-0.78



Overall SMD=0.28; 95%CI=0.06-0.59