


# Predictive value of urinary osmolality in the early morning for desmopressin response

DOSSCHE L.<sup>1, 3</sup>, RAES A.<sup>1,3</sup>, MAUEL R.<sup>1, 2,3</sup>, DE BRUYNE P.<sup>3</sup>, DEBRUYN N.<sup>1</sup>, CORNELIS A.S.<sup>1</sup>, VANDE WALLE J.<sup>1, 3</sup> <sup>1</sup>Department of Paediatric Nephrology, Ghent University Hospital, Ghent, Belgium <sup>2</sup>Department of Paediatric Nephrology, Vrije Universiteit Brussel, Belgium <sup>3</sup>On behalf of the SAFEPEDRUG consortium, http://safepedrug.eu

#### Introduction

**Desmopressin (dDAVP),** a synthetic vasopressin analogue, is a level 1, grade A treatment of monosymptomatic enuresis nocturna (MNE).



Although evidence based results show superiority of dDAVP above placebo, only 30 % of patients reach complete continence.

Surrogate parameters predicting the dDAVP response would be useful in individualizing therapy in children with nocturnal enuresis. They have been an issue for debate and research in the past decades.

Urinary osmolality in the morning was proposed by several authors, but without convincing results.


#### Aim

The aim of this study was to explore the potential value of urinary osmolality in an early morning sample as a surrogate parameter for dDAVP response.

### Method

Fig.1: Urinary osmolality in the early morning did not correlate with dDAVP response rate.

Full response: Reduction of at least 90% in the number of wet nights Non response: Reduction less than 50% in the number of wet nights



Study population

**45 children (>** 5 years old) with MNE. 33/ 12. Severe enuresis = wet at least 5/7 nights.


*Retrospective study* 

- 14 days diary for nocturnal enuresis episodes and diuresis-volume
- Morning sample collection at time of waking up Subsequent treatment with dDAVP 2x200µg tablets in the evening for minimum 6 weeks

## Results

- 13/45 patients were **full responder**
- 15/45 patients partial responder
- Urinary osmolality did not correlate with dDAVP response-rate (Fig.1)

Fig.2: Nocturnal diuresis volume < 100% of expected bladder capacity for age (EBC\*) correlated strongly with desmopressin resistance.



- Nocturnal diuresis volume < 100% of expected bladder capacity for age
- correlated strongly with **dDAVP resistance (Fig.2)** •Urinary osmolality-range was 675 to 1023 mosmol/l (Fig.3)

Fig.3: Urinary osmolality-range was 675 to 1023 mosmol/l. There is no difference in osmolality-range between the 3 different groups.

# <u>Conclusion</u>: Urinary osmolality in the morning has no predicting value in dDAVP - response

**CONTACT:** lien.dossche@ugent.be, johan.vandewalle@ugent.be Department of Paediatric Nephrology, Ghent University Hospital, Ghent, Belgium **ACKNOWLEDGEMENTS:** Supported by the Agency for Innovation by Science and Technology in Flanders (IWT) through the SAFEPEDRUG project (IWT/SBO 130033).

