

Linear-Optical Generation of Eigenstates of the Two-Site XY Model

Stefanie Barz,^{1,*} Borivoje Dakić,^{1,2} Yannick Ole Lipp,¹ Frank Verstraete,¹ James D. Whitfield,¹ and Philip Walther¹

¹*University of Vienna, Faculty of Physics, Boltzmannngasse 5, 1090 Vienna, Austria*

²*Institute for Quantum Optics and Quantum Information (IQOQI), Austrian Academy of Sciences, Boltzmannngasse 3, A-1090 Vienna, Austria*

(Received 30 September 2014; revised manuscript received 18 December 2014; published 24 April 2015)

Much of the anticipation accompanying the development of a quantum computer relates to its application to simulating dynamics of another quantum system of interest. Here, we study the building blocks for simulating quantum spin systems with linear optics. We experimentally generate the eigenstates of the XY Hamiltonian under an external magnetic field. The implemented quantum circuit consists of two CNOT gates, which are realized experimentally by harnessing entanglement from a photon source and applying a CPHASE gate. We tune the ratio of coupling constants and the magnetic field by changing local parameters. This implementation of the XY model using linear quantum optics might open the door to future studies of quenching dynamics using linear optics.

DOI: [10.1103/PhysRevX.5.021010](https://doi.org/10.1103/PhysRevX.5.021010)

Subject Areas: Optics, Quantum Physics

I. INTRODUCTION

In 1982, Feynman proposed the idea for the efficient simulation of quantum systems [1]. Complex systems, whose properties cannot easily be computed with classical computers, can be simulated by other well-controllable quantum systems. In this way, an easily accessible system can be used for reproducing the dynamics and the quantum state of another system of study. The insight of having one controllable quantum system simulate another is what forms the foundation of quantum simulation. There are two different approaches for simulating quantum systems that have been implemented experimentally: analog and digital simulation. Analog quantum simulators are designed to mimic a quantum system by reproducing its evolution in a faithful manner [2,3]. Alternatively, the effect of the unitary evolution of a quantum system may be regarded as that of a quantum circuit acting on some initial state. This inspires the approach of a digital quantum simulator where the state of the system is encoded into qubits and processed via quantum logic gates [4–7]. The main challenge—apart from providing a sufficiently powerful quantum computer—lies in finding a way to decompose the Hamiltonian into a suitable form. Experimentally, basic quantum simulations of both types have been demonstrated as proof-of-concept

experiments on several quantum architectures, including trapped ions [8–11], optical lattices [12,13], nuclear magnetic resonance [14–18], and photons [19–21].

Here, we exploit a scalable approach for digital quantum simulation for strongly interacting Hamiltonians, which was suggested in Ref. [22]. The general idea of Ref. [22] is to construct the explicit finite quantum circuits that transform the Hamiltonian into one corresponding to noninteracting particles. In this work, we apply this method to the XY Hamiltonian for two spins in a magnetic field. We develop a quantum circuit that transforms product input states to the eigenstates of this Hamiltonian. Our approach allows us to recover the whole spectrum of certain quantum many-body problems—a distinct advantage of our implementation. We experimentally implement this circuit in a linear optical setup and generate ground and excited states for the two-qubit Heisenberg XY model in a transverse external field. Our circuit consists of two CNOT gates, where one of the gates is absorbed in the state generation and the other is implemented physically.

II. THEORY

Our work focuses on the simulation of the two-qubit XY Hamiltonian in a transverse external field:

$$H = J_x \sigma_x \otimes \sigma_x + J_y \sigma_y \otimes \sigma_y + \frac{1}{2} B (\sigma_z \otimes \mathbb{1} + \mathbb{1} \otimes \sigma_z), \quad (1)$$

with J_x and J_y the coupling constants, B the magnetic field with unit magnetic moment, and σ_i the Pauli matrices that represent the particles' spin in the x , y , or z direction, respectively.

*Corresponding author.

barz@physics.ox.ac.uk

Present address: Clarendon Laboratory, Department of Physics, University of Oxford, Parks Road, Oxford OX1 3PU, United Kingdom.

Published by the American Physical Society under the terms of the *Creative Commons Attribution 3.0 License*. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Our work here focuses on preparing eigenstates. Specifically, our goal is to find a unitary U that transforms the Hamiltonian H into one corresponding to noninteracting quasiparticles \tilde{H} , hence diagonalizing it as

$$\tilde{H} = UHU^\dagger = \omega_1 \sigma_z \otimes \mathbb{1} + \omega_2 \mathbb{1} \otimes \sigma_z, \quad (2)$$

where $\omega_1 = (E_1 + E_2)/2$ and $\omega_2 = (E_1 - E_2)/2$ are the quasiparticle energies, and E_1, E_2 are the eigenenergies. It is not difficult to verify that the desired unitary U is given by

$$U = \begin{pmatrix} \cos \frac{w}{2} & 0 & 0 & \sin \frac{w}{2} \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\sin \frac{w}{2} & 0 & 0 & \cos \frac{w}{2} \end{pmatrix}, \quad (3)$$

with $\tan w = (J_x - J_y)/B$ and that $E_1 = \sqrt{B^2 + (J_x - J_y)^2} = -E_4$, $E_2 = J_x + J_y = -E_3$.

By applying U^\dagger to the computational basis states, the eigenstates of \tilde{H} , we obtain the eigenstates of the Hamiltonian H :

$$|\psi_1\rangle = U^\dagger|00\rangle = \cos \frac{w}{2}|00\rangle + \sin \frac{w}{2}|11\rangle, \quad (4)$$

$$|\psi_2\rangle = U^\dagger|01\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle), \quad (5)$$

$$|\psi_3\rangle = U^\dagger|10\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle), \quad (6)$$

$$|\psi_4\rangle = U^\dagger|11\rangle = -\sin \frac{w}{2}|00\rangle + \cos \frac{w}{2}|11\rangle. \quad (7)$$




FIG. 1. The quantum circuit for the generation of the eigenstates of the XY Hamiltonian. (a) The decomposition of the unitary U^\dagger that transforms the eigenstates of a noninteracting Hamiltonian to the eigenstates of the XY Hamiltonian. Full control of the system parameter w that specifies a particular Hamiltonian is granted by local operations. (b) The local unitaries $(YH) \otimes X$ together with the first CNOT transform the computational basis states into the four Bell states and we obtain the circuit shown in (b). The asterisk (*) is to remind us that this simplification is valid only for certain inputs.

We directly construct the circuit; the more general case, an arbitrary length XY spin chain, was presented in Ref. [22]. The steps for the general case are (1) identify spins with fermionic modes, (2) Fourier transform the fermionic modes, and (3) perform a Bogoliubov transformation to diagonalize the free fermions. The first step requires only relabeling while the second and third steps require actual transformations. The Fourier transform over L sites can be done in $L \log L$ steps. Additional L^2 gates are needed to account for the antisymmetry in the fermionic basis [22]. Finally, the Bogoliubov transformation requires only mixing positive and negative momenta pairwise. The parameters of this mixing depend on J_x, J_y, B and the momenta of the two Fourier modes being mixed. The full Bogoliubov transformation can be done with $L/2$ gates, which can all be done in parallel. The combined procedure implements a diagonalizing unitary similar to Eq. (2) in polynomial cost in L . Using photonic systems, the implementation of a heralded entangling gate requires two additional ancilla photons per two-qubit gate.

III. EXPERIMENT AND RESULTS

In our experiment, we generate the eigenstates using linear optics. To this end, we experimentally realize a




FIG. 2. Schematic of the experimental setup implementing U^\dagger [Fig. 1(b)]. A pair of photons—polarization-encoded qubits—is both created and initialized to a desired state via spontaneous parametric down-conversion. Different Hamiltonians can flexibly be simulated by tuning the parameter $w = \arctan[(J_x - J_y)/B]$ through local unitaries (LU) wrapped in between two CNOT operations. The former CNOT is thus merged with the SPDC process, while the latter is implemented by a combination of two-photon interference and polarization-dependent beam splitters [25,26,28].

flexible optical circuit (Fig. 2) that may implement any unitary $U^\dagger(w)$ corresponding to the two-qubit XY Hamiltonian [Eq. (1)] in a transverse field with the system parameter w . Our circuit consists of two CNOT gates and local operations, which allow the manipulation of w . Figure 1(a) shows that the first CNOT gate together with the preceding unitaries transforms the four product state inputs $|00\rangle, |01\rangle, |10\rangle, |11\rangle$ into one of the Bell states $|\psi^\pm\rangle = (|01\rangle \pm |10\rangle)/\sqrt{2}$, $|\phi^\pm\rangle = (|00\rangle \pm |11\rangle)/\sqrt{2}$. These Bell states can naturally be obtained by exploiting the entanglement of a spontaneous parametric down-conversion (SPDC) source. Thus, we integrate the first CNOT into the state preparation process [see Fig. 1(b)].

The input register in Fig. 1(b) originates from a type-II SPDC source, where a β -barium borate (BBO) crystal is pumped with a femtosecond-pulsed laser (394.5 nm, 76 MHz) to emit pairs of correlated photons at a wavelength of 789 nm [23]. In our implementation, $|0\rangle$ and $|1\rangle$ correspond to horizontal and vertical polarization, respectively. In our experiment, we generate entangled photon pairs in the four different Bell states and input them in the subsequent circuit. In combination with narrow-bandwidth filters of 3 nm, this procedure yields state fidelities for the input states of $97 \pm 1\%$.

Adjusting the subsequent local operations using a set of quarter-wave (QWPs) and half-wave plates (HWP) allows




FIG. 3. Characterization of the eigenstates of the XY Hamiltonian. We show the measurement of the three components of the two-qubit XY Hamiltonian separately. (a) Correlations when measuring the qubits in the basis $\sigma_x \otimes \sigma_x$. (b) Correlations when measuring the qubits in the basis $\sigma_y \otimes \sigma_y$. (c) Correlations when measuring the qubits in the basis $(\sigma_z \otimes \mathbb{1} + \mathbb{1} \otimes \sigma_z)/2$. This separate measurement of each part of the Hamiltonian makes it possible to see the measured data for each part separately and thus enables a more detailed analysis of each part of the Hamiltonian. The panels (a)–(c) show that an uncertainty in w would lead to only a small change in the measured correlation. (d) Fidelities of the generated states. In all panels, the lines show a fit of the experimental data. The error bars are smaller than the point size and hence not shown.

us to tune the system parameter w . The circuit for U^\dagger is completed by applying another CNOT gate. In our experiment, this destructive CNOT gate uses a polarization-dependent beam splitter (PDBS) that has a different transmission coefficient T for horizontally polarized light ($T_H = 1$) than for vertically polarized light ($T_V = 1/3$) [24]. If two vertically polarized photons are reflected at this PDBS, they acquire a phase shift of π . Two successive PDBSs with opposite splitting ratios then equalize the output amplitudes. This setup, in combination with two HWPs (see Fig. 2), implements a destructive CNOT gate, where the success of the operation is determined by postselection on a coincidence detection of the final output photons [24–26]. For this CNOT gate, we experimentally achieve a process fidelity [27] of $86.0 \pm 0.3\%$.

Using this setup we are able to prepare both ground and excited states at arbitrary values of the system parameter w by tuning the local unitaries in each input mode of the main polarization-dependent beam splitter. Figure 3 shows different correlation measurements to characterize these states for several choices of w . In the Supplemental Material [29], we show the reconstructed density matrices of all measured states; Fig. 3(d) shows the state fidelities as obtained from the density matrices.

Our demonstration shows that the main features of the XY Hamiltonian can be reproduced. The obtained fidelities lie between 0.75 and 0.9; these are the expected values when considering the fidelities of the entangled input states of 0.97 and a process fidelity of 0.86. Since the state fidelities of the experimental states are nonperfect, the measured data deviate from the theoretically expected values. Furthermore, the experimental eigenstates show weaker correlations as one would expect theoretically (see Fig. 3). However, as one can see in Fig. 3, the obtained states show the same behavior as one would expect from the theoretical eigenstates. Furthermore, the change of the eigenstates with respect to the system parameter is smooth and continuous. This means that small changes or imperfections in the parameter only lead to small changes in the results. In order to obtain data even closer to the values, one would need to increase the fidelity of the entangled input states, which are mainly limited by higher-order emissions in the current setup and can be increased using lower pump powers. Another limitation is the process fidelity of 0.86, which is mainly due to the nonperfect interference in our second CNOT gate. This interference could be improved by making the photons spectrally and spatially indistinguishable. In summary, the current state fidelities are mainly limited due to technical challenges, which can be overcome.

IV. CONCLUSION

We demonstrate the preparation of the eigenstates for the XY Hamiltonian under an external magnetic field. In the original proposal [22], it was pointed out that the same approach can also be applied to prepare thermal states and

simulate the dynamical evolution of any integrable model. Other examples of integrable models are the Kitaev honeycomb lattice [30], the 1D Hubbard model, and the Heisenberg models. Similar ideas can be used to simulate the models of Levin and Wen [31,32] in order to examine exotic features of topologically nontrivial models. This will be the basis for future research investigations.

We end this paper with a discussion on the extension to dynamical studies. The importance of generating eigenstates is underlined in the context of quenching, where the Hamiltonian of a system is instantaneously changed and the dynamics of a quantum system is examined. Recently, this problem has attracted significant interest [33,34], and it is a difficult task to simulate the quantum dynamics classically. In the quantum setting, utilizing algorithms such as the one implemented in the current work, initial states can be prepared and one could then perform evolution under a different Hamiltonian and observe the quenching properties for polynomial costs with a quantum computer. Since the XY model exhibits critical phases and quantum phase transitions, both adiabatic quenches through a phase transition and quenched dynamics can be studied using the present work as a starting point. While we did not explore dynamics in the present work, future work might begin with the preparation of eigenstates and proceed to break integrability, e.g., by including an additional magnetic field in the X direction and observing the dynamics of various observables. This will require the subsequent application of further entangling gates. However, currently the maximum number of subsequent photonic gates that have been demonstrated experimentally is two [35], which can be increased to three [36] when using entangling input states as demonstrated here. In that spirit, the technical challenges that need to be overcome for scaling up [37] our present scheme are the development of bright heralded photon sources [38–40], high-quality circuits implementing entangling gates with high quality, preferably using waveguide technology [41–44], and detection units with almost unity efficiency [45,46]. An ideal future implementation would be fully integrated with photons being directly generated, manipulated, and detected on chip, which allows us to achieve long-term stability and low losses. The generation of photons on chip has been demonstrated, with 6–8 photons being achievable in the near future. Current integrated waveguide circuits allow the implementation of complex quantum circuits with a high stability and also the integration of adaptive components such as phase shifters [47]. The limitations currently are the relatively high losses, which lead to relatively low counts rates and limit the complexity of the experiments that can be implemented. However, this is a technical limitation that can be overcome. Integrated detectors with high detection efficiency have been demonstrated and the extension of this work to higher photon numbers is a feasible engineering task [45].

ACKNOWLEDGMENTS

This work was supported by the European Commission, Q-ESSENCE (No. 248095), QUILMI (No. 295293), EQUAM (No. 323714), PICQUE (No. 608062), GRASP (No. 613024), and the ERA-Net CHISTERA project QUASAR, the John Templeton Foundation, the Ford Foundation, the Vienna Center for Quantum Science and Technology (VCQ), the Austrian Nano-initiative NAP Platon, the Austrian Science Fund (FWF) through the SFB FoQuS (F4006-N16), START (Y585-N20) and the doctoral program CoQuS, the Vienna Science and Technology Fund (WWTF, Grant No. ICT12-041), and the U.S. Air Force Office of Scientific Research (FA8655-11-1-3004).

- [1] R. Feynman, *Simulating Physics with Computers*, *Int. J. Theor. Phys.* **21**, 467 (1982).
- [2] I. Buluta and F. Nori, *Quantum Simulators*, *Science* **326**, 108 (2009).
- [3] A. Aspuru-Guzik and P. Walther, *Photonic Quantum Simulators*, *Nat. Phys.* **8**, 285 (2012).
- [4] S. Lloyd, *Universal Quantum Simulators*, *Science* **273**, 1073 (1996).
- [5] D. S. Abrams and S. Lloyd, *Simulation of Many-Body Fermi Systems on a Quantum Computer*, *Phys. Rev. Lett.* **79**, 2586 (1997).
- [6] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, *Simulated Quantum Computation of Molecular Energies*, *Science* **309**, 1704 (2005).
- [7] I. Kassal, S. P. Jordan, P. J. Love, M. Mohsenia, and A. Aspuru-Guzik, *Polynomial-Time Quantum Algorithm for the Simulation of Chemical Dynamics*, *Proc. Natl. Acad. Sci. U.S.A.* **105**, 18681 (2008).
- [8] A. Friedenauer, H. Schmitz, J. T. Glueckert, D. Porras, and T. Schaetz, *Simulating a Quantum Magnet with Trapped Ions*, *Nature (London)* **484**, 489 (2012).
- [9] B. Lanyon, C. Hempel, D. Nigg, M. Mller, R. Gerritsma, F. Zhringer, P. Schindler, J. T. Barreiro, M. Rambach, G. Kirchmair *et al.*, *Universal Digital Quantum Simulation with Trapped Ions*, *Science* **334**, 57 (2011).
- [10] J. W. Britton, B. C. Sawyer, A. C. Keith, C.-C. J. Wang, J. K. Freericks, H. Uys, M. J. Biercuk, and J. J. Bollinger, *Engineered Two-Dimensional Ising Interactions in a Trapped-Ion Quantum Simulator with Hundreds of Spins*, *Nature (London)* **484**, 489 (2012).
- [11] K. Kim, M.-S. Chang, S. Korenblit, R. Islam, E. E. Edwards, J. K. Freericks, G.-D. Lin, L.-M. Duan, and C. Monroe, *Quantum Simulation of Frustrated Ising Spins with Trapped Ions*, *Nature (London)* **465**, 590 (2010).
- [12] S. Trotzky, P. Cheinet, S. Flling, M. Feld, U. Schnorrberger, A. M. Rey, A. Polkovnikov, E. A. Demler, M. D. Lukin, and I. Bloch, *Time-Resolved Observation and Control of Super Exchange Interactions with Ultracold Atoms in Optical Lattices*, *Science* **319**, 295 (2008).
- [13] J. Simon, W. S. Bakr, R. Ma, M. E. Tai, P. M. Preiss, and M. Greiner, *Quantum Simulation of Antiferromagnetic Spin Chains in an Optical Lattice*, *Nature (London)* **472**, 307 (2011).
- [14] S. Somaroo, C. H. Tseng, T. F. Havel, R. Laflamme, and D. G. Cory, *Quantum Simulations on a Quantum Computer*, *Phys. Rev. Lett.* **82**, 5381 (1999).
- [15] C. Negrevergne, R. Somma, G. Ortiz, E. Knill, and R. Laflamme, *Liquid-State NMR Simulations of Quantum Many-Body Problems*, *Phys. Rev. A* **71**, 032344 (2005).
- [16] K. R. Brown, R. J. Clark, and I. L. Chuang, *Limitations of Quantum Simulation Examined by Simulating a Pairing Hamiltonian Using Nuclear Magnetic Resonance*, *Phys. Rev. Lett.* **97**, 050504 (2006).
- [17] J. Du, N. Xu, X. Peng, P. Wang, S. Wu, and D. Lu, *NMR Implementation of a Molecular Hydrogen Quantum Simulation with Adiabatic State Preparation*, *Phys. Rev. Lett.* **104**, 030502 (2010).
- [18] X. Peng, S. Wu, J. Li, D. Suter, and J. Du, *Observation of the Ground-State Geometric Phase in a Heisenberg XY model*, *Phys. Rev. Lett.* **105**, 240405 (2010).
- [19] C.-Y. Lu, W.-B. Gao, O. Gühne, X.-Q. Zhou, Z.-B. Chen, and J.-W. Pan, *Demonstrating Anyonic Fractional Statistics with a Six-Qubit Quantum Simulator*, *Phys. Rev. Lett.* **102**, 030502 (2009).
- [20] X.-S. Ma, B. Daki, W. Naylor, A. Zeilinger, and P. Walther, *Quantum Simulation of the Wavefunction to Probe Frustrated Heisenberg Spin Systems*, *Nat. Phys.* **7**, 399 (2011).
- [21] A. Orioux, J. Boutari, M. Barbieri, M. Paternostro, and P. Mataloni, *Quantum Simulation of the Wavefunction to Probe Frustrated Heisenberg Spin Systems*, *arXiv:1312.1102*.
- [22] F. Verstraete, J. I. Cirac, and J. I. Latorre, *Quantum Circuits for Strongly Correlated Quantum Systems*, *Phys. Rev. A* **79**, 032316 (2009).
- [23] P. G. Kwiat, E. Waks, A. G. White, I. Appelbaum, and P. H. Eberhard, *Ultrabright Source of Polarization Entangled Photons*, *Phys. Rev. A* **60**, R773 (1999).
- [24] N. Kiesel, C. Schmid, U. Weber, R. Ursin, and H. Weinfurter, *Demonstration of a Simple Entangling Optical Gate and Its Use in Bell-State Analysis*, *Phys. Rev. Lett.* **95**, 210505 (2005).
- [25] N. K. Langford, T. J. Weinhold, R. Prevedel, K. J. Resch, A. Gilchrist, J. L. O'Brien, G. J. Pryde, and A. G. White, *Linear Optics Controlled-Phase Gate Made Simple*, *Phys. Rev. Lett.* **95**, 210504 (2005).
- [26] R. Okamoto, H. F. Hofmann, S. Takeuchi, and K. Sasaki, *Demonstration of an Optical Quantum Controlled-NOT Gate without Path Interference*, *Phys. Rev. Lett.* **95**, 210506 (2005).
- [27] A. G. White, A. Gilchrist, G. Pryde, J. L. O'Brien, M. Bremner, and N. K. Langford, *Linear Optics Controlled-Phase Gate Made Simple*, *J. Opt. Soc. Am. B* **24**, 172 (2007).
- [28] N. Kiesel, C. Schmid, U. Weber, R. Ursin, and H. Weinfurter, *Measuring Two-Qubit Gates*, *Phys. Rev. Lett.* **95**, 210505 (2005).
- [29] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevX.5.021010> for density matrices of all states generated in the experiment.
- [30] H.-D. Chen and Z. Nussinov, *Exact Results of the Kitaev Model on a Hexagonal Lattice: Spin States, String and Brane Correlators, and Anyonic Excitations*, *J. Phys. A* **41**, 075001 (2008).

- [31] M. A. Levin and X.-G. Wen, *String-Net Condensation: A Physical Mechanism for Topological Phases*, *Phys. Rev. B* **71**, 045110 (2005).
- [32] M. Levin and X.-G. Wen, *Detecting Topological Order in a Ground State Wave Function*, *Phys. Rev. Lett.* **96**, 110405 (2006).
- [33] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, *Colloquium: Nonequilibrium Dynamics of Closed Interacting Quantum Systems*, *Rev. Mod. Phys.* **83**, 863 (2011).
- [34] M. A. Cazalilla and M. Rigol, *Focus on Dynamics and Thermalization in Isolated Quantum Many-Body Systems*, *New J. Phys.* **12**, 055006 (2010).
- [35] E. Martín-López, A. Laing, T. Lawson, R. Alvarez, X.-Q. Zhou, and J. L. O’Brien, *Experimental Realization of Shor’s Quantum Factoring Algorithm Using Qubit Recycling*, *Nat. Photonics* **6**, 773 (2012).
- [36] S. Barz, A. A.-G. I. Kassal, M. Ringbauer, Y. O. Lipp, B. Dakic, and P. Walther, “A Two-Qubit Photonic Quantum Processor and Its Application to Solving Systems of Linear Equations”, *Sci. Rep.* **4**, 6115 (2014).
- [37] E. Knill, R. Laflamme, and G. J. Milburn, *A Scheme for Efficient Quantum Computation with Linear Optics*, *Nature (London)* **409**, 46 (2001).
- [38] G. Harder, V. Ansari, B. Brecht, T. Dirmeier, C. Marquardt, and C. Silberhorn, *An Optimized Photon Pair Source for Quantum Circuits*, *Opt. Express* **21**, 13975 (2013).
- [39] J. B. Spring, P. S. Salter, B. J. Metcalf, P. C. Humphreys, M. Moore, N. Thomas-Peter, M. Barbieri, X.-M. Jin, N. K. Langford, W. S. Kolthammer *et al.*, *On-Chip Low Loss Heralded Source of Pure Single Photons*, *Opt. Express* **21**, 13522 (2013).
- [40] J. Silverstone, D. Bonneau, K. Ohira, N. Suzuki, H. Yoshida, N. Iizuka, M. Ezaki, C. Natarajan, M. Tanner, R. Hadfield *et al.*, *On-Chip Quantum Interference Between Silicon Photon Pair Sources*, *Nat. Photonics* **8**, 104 (2014).
- [41] M. A. Broome, A. Fedrizzi, S. Rahimi-Keshari, J. Dove, S. Aaronson, T. C. Ralph, and A. G. White, *Photonic Boson Sampling in a Tunable Circuit*, *Science* **339**, 794 (2013).
- [42] J. B. Spring, B. J. Metcalf, P. C. Humphreys, W. S. Kolthammer, X.-M. Jin, M. Barbieri, A. Datta, N. Thomas-Peter, N. K. Langford, D. Kundys *et al.*, *Boson Sampling on a Photonic Chip*, *Science* **339**, 798 (2013).
- [43] M. Tillmann, B. Dakić, R. Heilmann, S. Nolte, A. Szameit, and P. Walther, *Experimental Boson Sampling*, *Nat. Photonics* **7**, 540 (2013).
- [44] A. Crespi, R. Osellame, R. Ramponi, D. J. Brod, E. F. Galvão, N. Spagnolo, C. Vitelli, E. Maiorino, P. Mataloni, and F. Sciarrino, *Integrated Multimode Interferometers with Arbitrary Designs for Photonic Boson Sampling*, *Nat. Photonics* **7**, 545 (2013).
- [45] T. Gerrits, N. Thomas-Peter, J. C. Gates, A. E. Lita, B. J. Metcalf, B. Calkins, N. A. Tomlin, A. E. Fox, A. L. Linares, J. B. Spring *et al.*, *On-Chip, Photon-Number-Resolving, Telecommunication-Band Detectors for Scalable Photonic Information Processing*, *Phys. Rev. A* **84**, 060301 (2011).
- [46] F. Marsili, V. Verma, J. Stern, S. Harrington, A. Lita, T. Gerrits, I. Vayshenker, B. Baek, M. Shaw, R. Mirin *et al.*, *Detecting Single Infrared Photons with 93% System Efficiency*, *Nat. Photonics* **7**, 210 (2013).
- [47] B. J. Metcalf, J. B. Spring, P. C. Humphreys, N. Thomas-Peter, M. Barbieri, W. S. Kolthammer, X.-M. Jin, N. K. Langford, D. Kundys, J. C. Gates *et al.*, *Quantum Teleportation on a Photonic Chip*, *Nat. Photonics* **8**, 770 (2014).