

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/334662546>

# First report of *Mesocriconema sphaerocephalum* (Taylor, 1936) Loof, 1989 associated with carrot (*Daucus carota* subsp. *Stativus*) in Vietnam

Article in *Journal of nematology* · January 2019

DOI: 10.21307/jofnem-2019-048

CITATIONS

0

READS

7

6 authors, including:



**Duyen Nguyen**

Vietnam National University, Hanoi - University of Languages and International St...

3 PUBLICATIONS 4 CITATIONS

[SEE PROFILE](#)



**Tien Nguyen**

The Institute Of Ecology and Biological Resources

17 PUBLICATIONS 5 CITATIONS

[SEE PROFILE](#)



**Le Mai Linh**

The Institute Of Ecology and Biological Resources

24 PUBLICATIONS 9 CITATIONS

[SEE PROFILE](#)



**Phap Trinh**

The Institute Of Ecology and Biological Resources

35 PUBLICATIONS 141 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Entomopathogenic nematodes in Vietnam [View project](#)



Root-knot nematodes in Vietnam [View project](#)

# First report of *Mesocriconema sphaerocephalum* (Taylor, 1936) Loof, 1989 associated with carrot (*Daucus carota* subsp. *Stativus*) in Vietnam

Thi Duyen Nguyen<sup>1,2</sup>, Huu Tien Nguyen<sup>1,2</sup>, Thi Mai Linh Le<sup>1,2</sup>, Thi Tuyet Thu Tran<sup>3</sup>, Neriza Nobleza<sup>4</sup> and Quang Phap Trinh<sup>1,2,\*</sup>

<sup>1</sup>Institute of Ecology and Biological Resources, Vietnam Academy of Sciences and Technology, 18 Hoang Quoc Viet, Cau Giay, 100000 Hanoi, Vietnam.

<sup>2</sup>Graduate University of Science and Technology, Vietnam Academy of Sciences and Technology, 18 Hoang Quoc Viet, Cau Giay, 100000 Hanoi, Vietnam.

<sup>3</sup>Faculty of Environmental Sciences, VNU University of Science, 334 Nguyen Trai, Hanoi, Vietnam.

<sup>4</sup>College of Agriculture, Mindanao State University, Main Campus, Marawi City, 9700, Lanao del Sur, The Philippines.

\*E-mail: tqphap@gmail.com

This paper was edited by Zafar Ahmad Handoo.

Received for publication May 31, 2019.

## Abstract

Our study recorded the presence of *Mesocriconema sphaerocephalum* on carrot in Hanoi city and Hai Duong province in Vietnam. This species was identified by morphometric, morphological characterizations, and molecular characterization of D2D3 of 28S rDNA sequence. To our knowledge, this is the first report of *M. sphaerocephalum* on carrot in Vietnam.

## Key words

Carrot, Criconematidae, *Daucus carota* subsp. *Stativus*, *Mesocriconema sphaerocephalum*, Ring nematode, Vietnam.

*Mesocriconema* spp. belongs to the family Criconematidae Taylor, 1936 (1914) (Thorne, 1949). Species in this group are known as the ring nematodes, and they are one of important ectoparasitic nematodes that can be a potential threat at high soil population density. Some species of the ring nematodes caused yield loss up to 50% such as *Macroposthonia ornatum* (Raski, 1958) Loof and De Grisse, 1989 in pod field (Siddiqi, 2000).

Our study recorded *Mesocriconema sphaerocephalum* (Taylor, 1936) Loof, 1989 on carrot in Hanoi city and Hai Duong province in Vietnam. *Mesocriconema sphaerocephalum* was found on 10% of 130 soil samples. Density of this nematode was up to 45 individuals/250 g soil. To our knowledge, this is the first report of *M. sphaerocephalum* on carrot.

## Materials and methods

Nematodes were extracted from soil samples using modified Baermann tray method (Whitehead and Hemming, 1965). For morphological characterizations, permanent slides of nematodes were observed through the Carl Zeiss Axio Lab.A1 light microscope. Measurements and pictures were taken using a ZEN lite software on ZEISS AxioCam ERc5s digital camera (Nguyen et al., 2017).

For molecular studies, Primers D2A (5'-ACAAGTACCGTGGGAAA GTTG-3') and D3B (5'-TCGGAAGGAACCAGCTAC TA-3') were used to amplify D2D3 of 28S rDNA region (Nguyen et al., 2017). Obtained sequence was used for a Blast search in GenBank (Altschul et al., 1997). The data set was analyzed using maximum likelihood (ML) method in

**Table 1. Measurements of females of *M. sphaerocephalum* on carrot in Vietnam. All measurements are in  $\mu\text{m}$  and in form: mean  $\pm$  SD (range), except for ratio.**

Measurement	<i>M. sphaerocephalum</i>	
	Population in Hanoi city	Population in Hai Duong province
n	10	10
L	298 $\pm$ 13.8 (274–316)	330.7 $\pm$ 14.4 (309.4–360)
a	8.6 $\pm$ 0.5 (7.5–9.2)	9.7 $\pm$ 0.6 (8.8–10.6)
b	2.8 $\pm$ 0.1 (2.5–3)	3.2 $\pm$ 0.2 (3–3.5)
c	44 $\pm$ 9.8 (31–65)	52 $\pm$ 9.2 (39–69)
c'	0.4 $\pm$ 0.1 (0.3–0.5)	0.4 $\pm$ 0.1 (0.3–0.5)
VL/VB	0.8 $\pm$ 0.1 (0.7–1)	0.7 $\pm$ 0.1 (0.6–1)
V	94 $\pm$ 1 (93–96)	95 $\pm$ 0.4 (94–95)
Stylet length	55 $\pm$ 1.6 (53–58)	51 $\pm$ 1.5 (48–54)
Anterior end to nerve ring	82 $\pm$ 2.4 (78–85)	76 $\pm$ 2.6 (72–82)
Anterior end to end of pharynx	107 $\pm$ 2.7 (104–110)	103.3 $\pm$ 3 (98–109)
Anterior end to secretory-excretory pore	110 $\pm$ 4 (105–117)	108 $\pm$ 2.3 (104–113)
Max. body diam.	34.8 $\pm$ 1 (33–37)	34 $\pm$ 1.4 (31–37)
Body diam. at vulva	23.2 $\pm$ 1.4 (21–25)	24.3 $\pm$ 1.7 (22.2–28)
Body diam. at anus	17.7 $\pm$ 1.8 (14.3–19.4)	18 $\pm$ 2 (14.2–22)
Tail length	7 $\pm$ 1.3 (4.4–8.8)	6.6 $\pm$ 1.2 (4.8–9)
Rst	12.4 $\pm$ 0.5 (12–13)	12 $\pm$ 0.3 (12–13)
Roes	20.7 $\pm$ 0.8 (20–22)	20.4 $\pm$ 0.6 (20–22)
Rex	21.5 $\pm$ 0.7 (20–22)	21.6 $\pm$ 0.5 (21–22)
Rv	4.8 $\pm$ 0.4 (4–5)	4.3 $\pm$ 0.4 (4–5)
Ran	2.2 $\pm$ 0.4 (2–3)	2 $\pm$ 0.3 (2–3)
Rvan	3 $\pm$ 0.3 (2–3)	2.2 $\pm$ 0.4 (2–3)
R	65 $\pm$ 1.6 (63–68)	64 $\pm$ 1.4 (62–66)




Figure 1: *Mesocriconeema sphaerocephalum* on carrot in Vietnam. (A) entire body; (B) head region; (C) stylet; (D) anastomoses at mid-body; (E) pharyngeal region; (F) tail region; (G) anastomoses at tail region (Scale bar: 10µm).

MEGA 6 program with 1,000 bootstrap replications. The best fit model of DNA evolution was obtained using the Model test in MEGA 6 (Nguyen et al., 2017).

## Results and discussion

### Morphological characterization

Measurements of *M. sphaerocephalum* in this study are in agreement with measurement of *M. sphaerocephalum* in Geraert (2010) (Table 1). Females of *M. sphaerocephalum* on carrots are characterized by the following traits: body curved ventrally (Fig. 1A); lip region bearing two annuli with flattened labial disc (Fig. 1B); first body annulus much smaller than second one with smooth edge, sloping posteriorly (Fig. 1B); cuticle annuli at mid-body 4 to 5µm wide;

anastomoses numerous at lateral field, forming zigzag lines (Fig. 1D,G); stylet robust, knobs 9 to 10µm wide (Fig. 1C); vulva located near posterior end; tail rounded (Fig. 1F).

### Molecular characterization

D2D3 of 28S rDNA sequence of *M. sphaerocephalum* in this study was 730bp, submitted to GenBank under accession number: MK026628. It is 99% similar to *M. sphaerocephalum* (AB933464) in GenBank. This sequence varied 0 to 2% compared to other sequences of *M. sphaerocephalum* in GenBank. The maximum likelihood phylogenetic tree placed sequence of *M. sphaerocephalum* on carrots together with other sequences of *M. sphaerocephalum* from GenBank (Fig. 2).




Figure 2: Phylogenetic tree generated from D2D3 of 28S rDNA sequences based on ML method (TN93 + G model) with 1,000 replications. Sequence of *Mesocriconema sphaerocephalum* on carrot in Vietnam is in red.

## Acknowledgments

This research was supported by the project of Vietnam National University, Hanoi (code: QG.16.19); this project provided Criconematidae specimens on citrus in Hoa Binh to compare to the specimen in this study.

## References

Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D. J. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25:3389–402.

Geraert, E. 2010. The Criconematidae of the world: identification of the family Criconematidae (Nematoda). Academia Press, Ghent.

Nguyen, T. D., Le, T. M. L., Nguyen, H. T., Nguyen, T. A. D., Liébanas, G. and Trinh, Q. P. 2017. Morphological and molecular characteristics of *Pratylenchus haiduongensis* sp. n., a new species of root-lesion nematode associated with carrot in Vietnam. *Journal of Nematology*. 49:276–85.

Siddiqi, M. R. 2000. *Tylenchida: parasites of plants and insects*, CABI, London.

Whitehead, A. G. and Hemming, J. R. 1965. A comparison of some quantitative methods of extracting small vermiform nematodes from soil. *Annals of Applied Biology* 55:25–38.