

## **A Machine Learning Approach to Assess Injury Risk in Elite Youth Football Players**

Nikki Rommers<sup>1,2,3</sup>, Roland Rössler<sup>4,5</sup>, Evert Verhagen<sup>5,6,7</sup>, Florian Vandecasteele<sup>8</sup>, Steven Verstockt<sup>8</sup>, Roel Vaeyens<sup>2</sup>, Matthieu Lenoir<sup>2</sup>, Eva D'Hondt<sup>1</sup>, Erik Witvrouw<sup>9</sup>

<sup>1</sup>Department of Movement and Sports Sciences, Vrije Universiteit Brussel, Brussels, Belgium;

<sup>2</sup>Department of Movement and Sports Sciences, Ghent University, Ghent, Belgium; <sup>3</sup>Research

Foundation Flanders (FWO), Brussels, Belgium; <sup>4</sup>Department of Sport, Exercise, and Health,

University of Basel, Basel, Switzerland; <sup>5</sup>Amsterdam Collaboration on Health and Safety in

Sports & Department of Public and Occupational Health, Amsterdam Movement Science, VU

University Medical Center, Amsterdam, the Netherlands; <sup>6</sup>UCT/MRC Research Unit for

Exercise Science and Sports Medicine (ESSM), Department of Human Biology, Faculty of

Health Sciences, University of Capetown, Capetown, South Africa; <sup>7</sup>School of Physical

Education, Faculty of Physical Therapy & Occupational Therapy, Universidade Federal de

Minas Gerais, Belo Horizonte, Brazil; <sup>8</sup>imec, ELIS - IDLab, Ghent University, Ghent, Belgium;

<sup>9</sup>Department of Physical Therapy and Motor Rehabilitation, Ghent University, Ghent, Belgium

Accepted for Publication: 29 January 2020

**Medicine & Science in Sports & Exercise**®. Published ahead of Print contains articles in unedited manuscript form that have been peer reviewed and accepted for publication. This manuscript will undergo copyediting, page composition, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered that could affect the content.

# **A Machine Learning Approach to Assess Injury Risk in Elite Youth Football Players**

Nikki Rommers<sup>1,2,3</sup>, Roland Rössler<sup>4,5</sup>, Evert Verhagen<sup>5,6,7</sup>, Florian Vandecasteele<sup>8</sup>,  
Steven Verstockt<sup>8</sup>, Roel Vaeyens<sup>2</sup>, Matthieu Lenoir<sup>2</sup>, Eva D'Hondt<sup>1</sup>, Erik Witvrouw<sup>9</sup>

<sup>1</sup>Department of Movement and Sports Sciences, Vrije Universiteit Brussel, Brussels, Belgium;

<sup>2</sup>Department of Movement and Sports Sciences, Ghent University, Ghent, Belgium; <sup>3</sup>Research Foundation Flanders (FWO), Brussels, Belgium; <sup>4</sup>Department of Sport, Exercise, and Health, University of Basel, Basel, Switzerland; <sup>5</sup>Amsterdam Collaboration on Health and Safety in Sports & Department of Public and Occupational Health, Amsterdam Movement Science, VU University Medical Center, Amsterdam, the Netherlands; <sup>6</sup>UCT/MRC Research Unit for Exercise Science and Sports Medicine (ESSM), Department of Human Biology, Faculty of Health Sciences, University of Capetown, Capetown, South Africa; <sup>7</sup>School of Physical Education, Faculty of Physical Therapy & Occupational Therapy, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil; <sup>8</sup>imec, ELIS - IDLab, Ghent University, Ghent, Belgium; <sup>9</sup>Department of Physical Therapy and Motor Rehabilitation, Ghent University, Ghent, Belgium

## **Corresponding author:**

Roland Rössler

Birsstrasse 320 B, 4052 Basel, Switzerland

Email: Roland.Roessler@unibas.ch

**Conflicts of Interest and Source of Funding:** The Research Foundation – Flanders kindly supported this study through a PhD research grant awarded to Nikki Rommers (grant number 1116517N). The funding source had no involvement in the conduct and reporting of the study. None of the authors declares a conflict of interest.

ACCEPTED

## **Abstract**

Purpose: To assess injury risk in elite-level youth football players based on anthropometric, motor coordination and physical performance measures with a machine learning approach.

Methods: A total of 734 players in the U10 to U15 age categories (mean age: 11.7 +/- 1.7 years) from seven Belgian youth academies were prospectively followed during one season. Football exposure and occurring injuries were monitored continuously by the academies' coaching and medical staff, respectively. Preseason anthropometric measurements (height, weight, and sitting height) were taken and test batteries to assess motor coordination and physical fitness (strength, flexibility, speed, agility, and endurance) were performed. An extreme gradient boosting algorithms (XGBoost) was used to predict injury based on the preseason test results. Subsequently, the same approach was used to classify injuries as either overuse or acute.

Results: During the season, half of the players (n = 368) sustained at least one injury. Of the first occurring injuries, 173 were identified as overuse and 195 as acute injuries. The machine learning algorithm was able to identify the injured players in the hold-out test sample with 85% precision, 85% recall (sensitivity) and 85% accuracy (f1-score). Furthermore, injuries could be classified as overuse or acute with 78% precision, 78% recall and 78% accuracy.

Conclusion: Our machine learning algorithm was able to predict injury and to distinguish overuse from acute injuries with reasonably high accuracy based on preseason measures. Hence, it is a promising approach to assess injury risk among elite-level youth football players. This new

knowledge could be applied in the development and improvement of injury risk management strategies, to identify youth players with the highest injury risk.

**Keywords:** injury prevention, adolescent, child, soccer

ACCEPTED

## **Introduction**

Elite-level youth football is known to entail a high injury risk (1, 2), which is often attributed to early specialisation, high training loads, and high training and game intensities (3, 4). In order to specifically target injury risk mitigation strategies in this group, knowledge of both modifiable and non-modifiable risk factors is crucial. The few existing studies in elite-level youth football, mainly focus on anthropometric factors, landing strategies and a limited number of functional tasks (5-9). In practice, however, it is often not feasible for clubs and coaches to perform thorough player screening for injury risk management purposes due to limited time and financial means. Therefore, there is a strong interest to assess injury risk based on field-specific and relatively easy screening tests, such as motor performance tests already taken by many clubs to monitor player development.

Previous studies did not show an association between motor performance measures and injury risk in different cohorts of youth football players (8, 10). Due to the complex nature of injuries, a single variable is usually insufficient to provide a meaningful estimate the injury risk of a player (11). Hence, considering the multifactorial nature of injuries, it is key to incorporate a multitude of risk factors in any injury risk analysis (11). More advanced data science approaches, including machine learning, often better integrate the complexity of injury events compared to traditional statistics (12) and these approaches can take many variables and interactions between these into account (13). To date, this kind of approach has mainly been used in relatively small samples of elite athletes, mostly considering training load variables, but already demonstrated promising results (14-18).

To extend this knowledge from data science, the aim of this study was to use a machine learning approach to evaluate the risk of injury in youth elite-level football players, based on anthropometric, motor performance, and demographic measures. The first aim was to use preseason test results to assess the accuracy of a machine learning model predicting injury during the season. The second aim was to apply a similar model to correctly classify different types of injuries, namely overuse and acute injuries.

## **Methods**

### **Participants and design**

The present prospective study followed 734 male youth football players during the 2017-2018 competitive season (August 2017 until May 2018). Players of the under 10 (U10) up to the under 15 (U15) age categories were recruited from the youth academies of seven Belgian premier league football clubs. All players being medically cleared to play at the start of the season were eligible to participate. Verbal and written information about the study design and potential risks and benefits upon participation was provided to the players and their parents or legal caretakers. The parents provided written informed consent and the youth player provided his written assent. The study protocol was approved by the medical ethical committee of the Vrije Universiteit Brussel (B.U.N. 143201628616) and all measurements were performed according to the ethical standards of the Helsinki Declaration.

### **Procedures**

At the start of the competitive season (August 2017) a test battery of anthropometric, motor coordination and physical fitness measures was taken by a group of trained assessors,

coordinated by the principal investigator (NR) to guarantee standardization of all tests. At the same time, demographic information regarding the football experience (in years) and date of birth were collected through a questionnaire.

### **Anthropometry, growth and maturity status**

Body height (Seca 213 Portable Stadiometer, Seca, Germany) and sitting height (Harpender sitting height table, Holtain, UK) were measured to the nearest 0.1 cm. We calculated the players' leg length as the difference between their recorded body height and sitting height. Body weight was determined to the nearest 0.1 kg using a digital scale (Tanita BC-420SMA, Tanita, Japan). The years from peak height velocity (PHV), an indicator for maturational timing, was estimated using the prediction equation of Mirwald and colleagues (19).

### **Motor coordination**

We determined players' level of generic motor coordination using the three subtest short version of the Körperkoordinationstest für Kinder (KTK) (20, 21), consisting of (1) jumping sideways (JS), (2) moving sideways (MS), and (3) balancing backwards (BB). These three subtests were conducted as described by Kiphard and Schilling (22). All individual subtest scores as well as the total sum score were included in the analysis.

Football-specific motor coordination was tested using the Ghent University dribbling test (23). The players completed a prescribed circuit as fast as possible in two conditions: first without the ball (agility), and subsequently with the ball (dribbling skills). The time of both attempts was measured to the nearest 0.01 seconds with a handheld stopwatch.

## **Physical performance**

Flexibility of the lower back and posterior chain was measured using the sit and reach test to the nearest 0.5 cm (24). To test the players' endurance, the YoYo Intermittent Recovery Test Level 1 (YYIR1) was conducted on artificial turf in dry weather conditions according to the guidelines of Krstrup and colleagues (25). In addition, three strength measures were included in our test battery: the standing broad jump (SBJ), countermovement jump (CMJ) and curl-ups. The SBJ was executed as described in the Eurofit test battery's manual (26). The CMJ performance was tested in two conditions, using the Optojump system (OptoJump System, Microgate, Italy): first the player performed 3 jumps with the hands on the hips, followed by 3 jumps with an arm swing (27). The best attempt in each of the conditions of the SBJ and CMJ was used for analysis. We tested abdominal strength by the number of correctly executed curl-ups in 30 sec (26). Speed and agility were tested by a repeated sprint test (28) and a T-test, respectively. The players performed four maximal sprints of 30 m, with 25 sec of rest in between. Fastest split times at 5 m, 10 m, 20 m and 30 m as well as maximum decline over 30 m were used for analysis. Agility was assessed by the T-test that was executed twice: first with all turns performed left, and subsequently with all turns performed right (23). Both tests were performed on artificial turf with time registered to the nearest 0.001 seconds (MicroGate Racetime2, Microgate, Italy).

## **Football exposure**

Individual match and training exposure were recorded by the team coaches according to the consensus statement of Fuller and colleagues (29) in order to calculate the injury incidence.

Attendance at training and individual playing time during matches were registered in the player monitoring system, or by the use of Excel sheets designed by the research team.

### **Injury registration**

Injuries were registered by the medical staff of each youth academy using a prespecified injury registration form. On this form, the date of occurrence, type and location of injury were registered. The date of return to full participation was later added. An injury was defined as a medical attention injury, any injury that required an assessment of medical or paramedical staff (29). All participating youth academies had medical staff available at every training and match, and all players experiencing any physical complaint, had to seek assessment from the medical staff. The injuries were classified as either overuse or acute injuries. An overuse injury was defined as an injury without a single identifiable event responsible for the injury, whereas an acute injury resulted from a specific identifiable event (29). The first occurring injury for each player was taken into account in our analysis.

### **Statistical analysis**

Baseline characteristics of the players are presented as means and standard deviations for all variables. We used a machine learning approach to predict injury based on the preseason screening test results. In a second model, we aimed to correctly classify acute and overuse injuries. All analyses were performed in Python using the XGBoost application (version 0.81), an optimised distributed gradient boosting library. XGBoost is a highly flexible and versatile tool that was built for the purpose of model performance and computational speed. It supports fine tuning of the model and addition of regularisation parameters to control overfitting (13).

Therefore, it is a suitable application for the purpose of our study. We built a model using our training data consisting of a random sample of 80% of all collected data. During the boosting process, a set of weak learners is combined to improve prediction accuracy. Gradient descent was used to minimise the cost function parameterised by a specific model parameter. Finally, to optimise the cost function, hinge loss was used. Hinge loss penalises predictions both when they are incorrect and when they are correct but not confident. After the model was built, it was evaluated using cross-validation. Also, grid search was performed for hyperparameter optimisation. Finally, the best performing model was tested on our test data, consisting of the remaining 20% of all collected data.

To interpret and visualise the output of each model, we used the Shapley Additive exPlanations (SHAP) approach (SHAP summary plot) (30). This approach visualises every single player or injury case and gives an overview of the variables in the model by order of importance (vertically listed features), with the top ones having a higher global impact on the model than bottom ones. The SHAP-values represent the impact of a variable in the decision making process. Dots representing the SHAP-values for each feature value of a player in the dataset are plotted horizontally next to the feature. Positive SHAP-values represent a higher probability of a positive prediction (i.e. being injured). Each dot is coloured by the value (i.e. measured value) of the feature for an individual, where blue represents the lower values (e.g. lower body height), and red the higher values (e.g. higher body height). To assess the value of the models, precision (ratio of the number of true positive observations over the total number of predicted positive observations), recall or sensitivity (ratio of the correctly predicted positive observations to all

observations that were actually positive) and f1-score (harmonic mean of precision and recall) were calculated for both the training data set (80%) and the test data set (20%).

## **Results**

Data from 734 players of the U10 to U15 age categories were used in the analysis. During the 2017-2018 season, 368 of them sustained at least one injury during 129.206 hours of football exposure in total (112.745 hours of training and 16.464 hours of match exposure). An overview of the player characteristics for the total sample, the injured and the non-injured players can be found in Table 1. A detailed overview of the injuries by mechanism, type, location and severity is presented in Table 2.

## **Predicting injury**

The extreme gradient boosting model was able to predict injury with a precision of 84% in the training dataset supported by 587 players. This means that only 16% of players indicated by the model as injured actually remained uninjured. This model had a recall of 83%, and a f1-score of 83%. When the model was run on the test data supported by 147 players, the precision, recall and f1-scores were all 85%, showing a reasonable accuracy and sensitivity of the model. The most important variables related to injury risk, are shown using a SHAP approach (Figure 1A). A higher predicted age at PHV, higher body height and leg length, lower fat percentage and average performance on the standing broad jump were identified as the five most important predictors for injury. A better sit-and-reach performance, average t-test and lower 10 meter sprint performance affected the prediction for a smaller amount.

## **Classifying overuse and acute injuries**

Player characteristics by injury mechanism are displayed in Table 1. All 368 injuries were either classified as overuse ( $n = 173$ ) or acute ( $n = 195$ ) injuries. The extreme gradient boosting model supported by 294 injuries in the training data set, reached a precision and recall score of 82% in classifying injuries, and f1-score of 81%. In the test data set, supported by 74 injuries, a precision, recall and f1-score of 78% was reached. This shows that the model is reasonably accurate in classifying injuries correctly by mechanism. The most influential variables in the decision making process are visualized by the SHAP approach (Figure 1B). Most variables only affected the decision making for a limited amount with SHAP-values not exceeding 0.4. A lower predicted age at PHV, higher sitting height, slower T-test performance and lower MS score were related to overuse injuries.

## **Discussion**

The results of this prospective study show that it is possible to predict injury over the season with 85% accuracy in the unseen test data. Only 15% false positives were observed and 15% injured players misclassified as uninjured based on preseason anthropometric, motor performance and demographic measures. Moreover, overuse injuries could be distinguished from acute injuries with a slightly lower accuracy using the same preseason test results.

Two previous studies did not find an association between preseason performance tests and injury risk using traditional statistics (8, 10). In contrast to traditional approaches, boosted regression tree models, that are designed to maximise the precision and accuracy in classification, allow us to include many more variables than multivariate models used before. This different approach

aiming for the highest classification accuracy, could explain the contradictory findings of our study compared to previous ones identifying associations between certain variables and injury risk. The added clinical value of a machine learning model is the interpretation of the classification based on all variables, next to the interpretation of individual variables as risk factors. Based on the SHAP-values, we can indicate which variables and corresponding values of these variables played the largest role in the classification decisions.

### **Predicting injury**

The five most important variables that predict injury are anthropometric measures. Interpreting the SHAP-values, we notice that a higher predicted age at PHV, longer legs, higher body height, and lower body fat percentage increased injury risk. These findings can be explained by the large increase in injury incidence with age (31-33). Our study confirms the findings of previous studies, identifying higher body height and weight (7), and higher predicted age at PHV (34) as risk factors for injuries. Interestingly, we see that a worse performance on the sit and reach test was protective against injuries, whereas a good performance increased the risk of injury classification to a small extent. This is not in line with previous research showing that a lack of muscle flexibility is a risk factor for muscle injuries in adult players (35). In our sample of youth players, however, only few non-contact muscle strain injuries have been reported. The other motor performance measures that appear amongst the 20 most important variables, influence many classifications, but only affected the classification of being injured or not for a small amount.

### **Overuse and acute injuries**

Our model aimed to distinguish injured players whose first injury was an acute one, from players who sustained an overuse injury. To our knowledge, no previous study aimed to distinguish between types of injuries. Our model showed that performance measures appear amongst the five most influential variables. After age at PHV, the most important variable is performance on the moving sideways task, followed by the 20 m sprint time and the t-test performed with left turns. None of these individual variables have previously been associated with overuse or acute injuries specifically (5, 10). Future research should focus on risk factors for specific types of injuries, to get a more conclusive view of risk profiles.

### **Perspectives and practical applications**

With the models built in this study, a practitioner could screen the players preseason and determine the probability of injury for each player with reasonably high precision and accuracy. Subsequently, a prediction of the most likely type of injury (i.e. overuse or acute) could be made for players likely to get injured, by entering their test data in the other model. The outcomes of the models give the practitioner an estimation of the players who are most in need of injury risk management initiatives. The additional knowledge of the machine learning model could help clubs to spend their available time and funds for extensive screening and injury risk management more effectively.

The possibility to reach a high accuracy in classifying injured and non-injured players and types of injuries based on a combination of anthropometric, motor performance and demographic measures, suggests that risk profiles for certain types of injuries exist, in which none of the risk

factors shows a high individual association with injury occurrence. We therefore suggest that future research continues to look for risk profiles. Furthermore, our models could be validated with unknown data from a different cohort to investigate the generalizability in a larger, potentially international population of elite-level youth football players.

### **Strengths and limitations**

This is the largest prospective study to date closely monitoring youth football players over an entire season and collecting standardised test data. The proposed combined test battery is simple to use in the field and also has its value in talent identification and development purposes (36), which increases the practical relevance of our study protocol. A major limitation of our study is that only the first occurring injury of every player was considered in our analyses, since the motor performance measures potentially change after return from an injury. Consequently, as players can sustain multiple injuries over one season, the analysis does not reflect the complete picture. Furthermore, we only tested the players at the start of the competitive seasons and then monitored injuries over the entire season. Anthropometric and motor performance measures change over the course of the season due to training and natural development. We therefore suggest that future studies take the proposed test battery every few months to have assessments closer to the time of injury.

### **Conclusion**

In this unique study, we observed that a machine learning model was reasonably accurate in the prediction of injury in elite-level youth football players based on preseason test results. It is also possible to classify players sustaining overuse or acute injuries with a slightly lower accuracy

based on the same measures. Practitioners could use this information to assess the risk of particular types of injuries before the start of the competitive season. This information would allow academies to focus the available (financial) resources for injury risk management on those players with a higher injury risk.

ACCEPTED

**Acknowledgement:** The authors would like to thank the participating youth academies, coaches, and players for their collaboration and the test team for their contribution to the data collection. The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation, and results of the present study do not constitute endorsement by ACSM.

**Conflicts of Interest and Source of Funding:** The Research Foundation – Flanders kindly supported this study through a PhD research grant awarded to Nikki Rommers (grant number 1116517N). The funding source had no involvement in the conduct and reporting of the study. None of the authors declares a conflict of interest.

## References

1. Pfirrmann D, Herbst M, Ingelfinger P, Simon P, Tug S. Analysis of Injury Incidences in Male Professional Adult and Elite Youth Soccer Players: A Systematic Review. *Journal of athletic training*. 2016;51(5):410-24.
2. Faude O, Rössler R, Junge A. Football injuries in children and adolescent players: are there clues for prevention? *Sports medicine (Auckland, N.Z.)*. 2013;43(9):819-37.
3. Jayanthi NA, LaBella CR, Fischer D, Pasulka J, Dugas LR. Sports-specialized intensive training and the risk of injury in young athletes: a clinical case-control study. *The American journal of sports medicine*. 2015;43(4):794-801.
4. Read PJ, Oliver JL, De Ste Croix MB, Myer GD, Lloyd RS. The scientific foundations and associated injury risks of early soccer specialisation. *Journal of sports sciences*. 2016;34(24):2295-302.
5. Rössler R, Junge A, Chomiak J et al. Risk factors for football injuries in young players aged 7 to 12 years. *Scandinavian journal of medicine & science in sports*. 2018;28(3):1176-82.
6. Emery CA. Risk factors for injury in child and adolescent sport: a systematic review of the literature. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine*. 2003;13(4):256-68.
7. Kemper GL, van der Sluis A, Brink MS, Visscher C, Frencken WG, Elferink-Gemser MT. Anthropometric Injury Risk Factors in Elite-standard Youth Soccer. *International journal of sports medicine*. 2015;36(13):1112-7.

8. Emery CA, Meeuwisse WH, Hartmann SE. Evaluation of risk factors for injury in adolescent soccer: implementation and validation of an injury surveillance system. *The American journal of sports medicine*. 2005;33(12):1882-91.
9. Read PJ, Oliver JL, De Ste Croix MBA, Myer GD, Lloyd RS. A prospective investigation to evaluate risk factors for lower extremity injury risk in male youth soccer players. *Scandinavian journal of medicine & science in sports*. 2018;28(3):1244-51.
10. Frisch A, Urhausen A, Seil R, Croisier JL, Windal T, Theisen D. Association between preseason functional tests and injuries in youth football: a prospective follow-up. *Scandinavian journal of medicine & science in sports*. 2011;21(6):e468-76.
11. Bahr R, Holme I. Risk factors for sports injuries--a methodological approach. *British journal of sports medicine*. 2003;37(5):384-92.
12. Me E, Unold O. Machine learning approach to model sport training. *Computers in human behavior*. 2011;27(5):1499-506.
13. Chen T, Guestrin C. Xgboost: A scalable tree boosting system. In: *Proceedings of the Proceedings of the 22nd acm sigkdd international conference on knowledge discovery and data mining*. 2016: San Francisco, California (USA). p. 785-94.
14. Carey DL, Ong K, Whiteley R, Crossley KM, Crow J, Morris ME. Predictive modelling of training loads and injury in Australian football. *International Journal of Computer Science in Sport*. 2018;17(1):49-66.
15. Lopez-Valenciano A, Ayala F, Puerta JM et al. A Preventive Model for Muscle Injuries: A Novel Approach based on Learning Algorithms. *Medicine and science in sports and exercise*. 2018;50(5):915-27.

16. Rossi A, Pappalardo L, Cintia P, Iaia FM, Fernandez J, Medina D. Effective injury forecasting in soccer with GPS training data and machine learning. *PloS one*. 2018;13(7):e0201264.
17. Ruddy JD, Shield AJ, Maniar N et al. Predictive Modeling of Hamstring Strain Injuries in Elite Australian Footballers. *Medicine and science in sports and exercise*. 2018;50(5):906-14.
18. Thornton HR, Delaney JA, Duthie GM, Dascombe BJ. Importance of Various Training-Load Measures in Injury Incidence of Professional Rugby League Athletes. *International journal of sports physiology and performance*. 2017;12(6):819-24.
19. Mirwald RL, Baxter-Jones AD, Bailey DA, Beunen GP. An assessment of maturity from anthropometric measurements. *Medicine and science in sports and exercise*. 2002;34(4):689-94.
20. Novak AR, Bennett KJ, Beavan A et al. The Applicability of a Short Form of the KörperKoordinationsTest für Kinder for Measuring Motor Competence in Children Aged 6-11 Years. *Journal of Motor Learning and Development*. 2016;0(0):1-20.
21. Vandorpe B, Vandendriessche J, Lefevre J et al. The KörperkoordinationsTest für Kinder: reference values and suitability for 6-12-year-old children in Flanders. *Scandinavian journal of medicine & science in sports*. 2011;21(3):378-88.
22. Kiphard ES, F. *Körperkoordinationstest für Kinder 2. Überarbeitete und ergänzte Auflage. (Körperkoordinationstest für Kinder 2. Revised and supplemented edition)* Beltz: Test GmbH; 2007.
23. Vandendriessche JB, Vaeyens R, Vandorpe B, Lenoir M, Lefevre J, Philippaerts RM. Biological maturation, morphology, fitness, and motor coordination as part of a selection

- strategy in the search for international youth soccer players (age 15-16 years). *Journal of sports sciences*. 2012;30(15):1695-703.
24. Castro-Piñero J, Chillón P, Ortega F, Montesinos J, Sjöström M, Ruiz J. Criterion-related validity of sit-and-reach and modified sit-and-reach test for estimating hamstring flexibility in children and adolescents aged 6–17 years. *International journal of sports medicine*. 2009;30(09):658-62.
  25. Krstrup P, Mohr M, Amstrup T et al. The yo-yo intermittent recovery test: physiological response, reliability, and validity. *Medicine and science in sports and exercise*. 2003;35(4):697-705.
  26. Council of Europe. Eurofit: European test of physical fitness. In: Council of Europe, Committee for the development of Sport Rome; 1988.
  27. Bosco C, Rusko H, Hirvonen J. The effect of extra-load conditioning on muscle performance in athletes. *Medicine and science in sports and exercise*. 1986;18(4):415-9.
  28. Wragg CB, Maxwell NS, Doust JH. Evaluation of the reliability and validity of a soccer-specific field test of repeated sprint ability. *European journal of applied physiology*. 2000;83(1):77-83.
  29. Fuller CW, Ekstrand J, Junge A et al. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine*. 2006;16(2):97-106.
  30. Lundberg SM, Erion GG, Lee S-I. Consistent individualized feature attribution for tree ensembles. *arXiv preprint arXiv:1802.03888*. 2018.

31. Le Gall F, Carling C, Reilly T, Vandewalle H, Church J, Rochcongar P. Incidence of injuries in elite French youth soccer players: a 10-season study. *The American journal of sports medicine*. 2006;34(6):928-38.
32. Read PJ, Oliver JL, De Ste Croix MBA, Myer GD, Lloyd RS. An audit of injuries in six english professional soccer academies. *Journal of sports sciences*. 2018;36(13):1542-8.
33. Price RJ, Hawkins RD, Hulse MA, Hodson A. The Football Association medical research programme: an audit of injuries in academy youth football. *British journal of sports medicine*. 2004;38(4):466-71.
34. van der Sluis A, Elferink-Gemser MT, Brink MS, Visscher C. Importance of peak height velocity timing in terms of injuries in talented soccer players. *International journal of sports medicine*. 2015;36(4):327-32.
35. Witvrouw E, Danneels L, Asselman P, D'Have T, Cambier D. Muscle flexibility as a risk factor for developing muscle injuries in male professional soccer players: a prospective study. *The American journal of sports medicine*. 2003;31(1):41-6.
36. Deprez DN, Fransen J, Lenoir M, Philippaerts RM, Vaeyens R. A retrospective study on anthropometrical, physical fitness, and motor coordination characteristics that influence dropout, contract status, and first-team playing time in high-level soccer players aged eight to eighteen years. *Journal of strength and conditioning research*. 2015;29(6):1692-704.

## CAPTIONS

**Table 1.** Preseason test results for the total sample and for injured and non-injured players separately, as well as for the players who sustained an overuse or an acute injury.

**Table 2.** Injury characteristics.

**Figure 1.** SHAP (SHapley Additive exPlanations). The features in the model are listed from the relatively most (top) to least (bottom) important by their global impact on the model. Dots representing the SHAP-values for each feature value of an individual in the dataset are plotted horizontally next to the feature. Overlapping points are jittered in y-axis direction, so we get a sense of the distribution of the Shapley values per variable. The higher the absolute value (either positive or negative), the higher the importance in the classification decision-making process. Positive SHAP-values represent a higher probability of a positive prediction (i.e. being injured). Each dot is coloured by the value (i.e. measured value) of the feature for an individual, where blue represents the lower values (e.g. worse standing broad jump score), and red the higher values (e.g. better standing broad jump score). A grey dot represents a missing value.

- A. SHAP values of the variables in the model predicting injury. Positive SHAP values represent a higher chance of injury.

B. SHAP values of the variables in the model classifying overuse and acute injuries. Positive SHAP values represent the risk of acute injuries, whereas negative values correspond to overuse injury risk.

PHV: peak height velocity, y: year, cm: centimetre, sec: seconds, BMI: body mass index, kg: kilogram, m: meter, #: number of, KTK3: 3 subtest short version of the Körperkoordinationstest für Kinder, YoYo IR1: YoYo Intermittent Recovery Test Level 1

ACCEPTED

Figure 1A





Figure 1B



**Table 1.** Preseason test results for the total sample and for injured and non-injured players separately, as well as for the players who sustained an overuse injury and the players who sustained an acute injury.

|                              | <b>Total sample<br/>(N = 734)</b> | <b>Injured<br/>(n = 368)</b> | <b>Non-injured<br/>(n = 366)</b> | <b>Overuse injury<br/>(n = 172)</b> | <b>Acute injury<br/>(n = 196)</b> |
|------------------------------|-----------------------------------|------------------------------|----------------------------------|-------------------------------------|-----------------------------------|
|                              | Mean (SD)                         | Mean (SD)                    | Mean (SD)                        | Mean (SD)                           | Mean (SD)                         |
| Age (y)                      | 11.7 (1.7)                        | 12.2 (1.7)                   | 11.2 (1.5)                       | 12.1 (1.7)                          | 12.2 (1.7)                        |
| Height (cm)                  | 149.4 (12.4)                      | 152.8 (13.4)                 | 146.5 (10.7)                     | 153.0 (13.1)                        | 152.5 (13.7)                      |
| Weight (kg)                  | 39.3 (10.1)                       | 41.9 (11.2)                  | 37.0 (8.3)                       | 42.3 (11.2)                         | 41.6 (11.2)                       |
| Fat percentage (%)           | 17.3 (1.9)                        | 12.4 (3.6)                   | 13.5 (3.3)                       | 12.7 (3.7)                          | 12.2 (3.6)                        |
| BMI (kg/m <sup>2</sup> )     | 13.0 (3.5)                        | 17.6 (2.0)                   | 17.0 (1.6)                       | 17.7 (2.1)                          | 17.5 (1.9)                        |
| Sitting height (cm)          | 77.5 (5.8)                        | 78.8 (6.3)                   | 76.4 (5.2)                       | 78.9 (6.3)                          | 78.7 (6.4)                        |
| Leg length (cm)              | 71.8 (7.4)                        | 74.0 (7.9)                   | 70.0 (6.4)                       | 74.0 (7.6)                          | 73.9 (8.2)                        |
| Maturity offset (y)          | -2.0 (1.4)                        | -1.6 (1.5)                   | -2.3 (1.2)                       | -1.6 (1.5)                          | -1.6 (1.5)                        |
| Age at PHV (y)               | 13.7 (0.6)                        | 13.8 (0.6)                   | 13.6 (0.5)                       | 13.7 (0.6)                          | 13.8 (0.6)                        |
| SBJ (cm)                     | 170.2 (21.5)                      | 174.2 (23.3)                 | 166.7 (19.2)                     | 173.6 (23.3)                        | 174.8 (23.3)                      |
| CMJ (cm)                     | 24.9 (5.2)                        | 26.0 (5.6)                   | 23.9 (4.6)                       | 26.0 (5.7)                          | 26.1 (5.5)                        |
| CMJ with arm swing (cm)      | 29.7 (5.9)                        | 31.1 (6.4)                   | 28.5 (5.2)                       | 30.9 (6.4)                          | 31.3 (6.4)                        |
| Curl-ups (#)                 | 33.9 (6.8)                        | 35.4 (6.9)                   | 32.6 (6.4)                       | 35.6 (6.7)                          | 35.2 (7.0)                        |
| Sit and reach (cm)           | 19.4 (5.8)                        | 20.2 (5.5)                   | 18.8 (6.0)                       | 20.2 (5.2)                          | 20.2 (5.7)                        |
| Jumping sideways (#)         | 94.6 (12.7)                       | 96.0 (12.7)                  | 93.5 (12.5)                      | 95.3 (12.5)                         | 96.6 (12.9)                       |
| Moving sideways (#)          | 57.9 (10.2)                       | 59.3 (10.6)                  | 56.7 (9.6)                       | 58.7 (10.3)                         | 59.8 (10.9)                       |
| Balancing backwards (#)      | 55.5 (11.5)                       | 56.7 (11.2)                  | 54.4 (11.7)                      | 56.0 (10.9)                         | 57.4 (11.4)                       |
| KTK3 sum score               | 207.9 (29.3)                      | 211.9 (29.5)                 | 204.3 (28.6)                     | 210.0 (28.4)                        | 213.7 (30.5)                      |
| T-test left (sec)            | 8.7 (0.5)                         | 8.7 (0.5)                    | 8.8 (0.5)                        | 8.7 (0.5)                           | 8.6 (0.5)                         |
| T-test right (sec)           | 8.8 (0.5)                         | 8.8 (0.5)                    | 8.9 (0.5)                        | 8.8 (0.5)                           | 8.7 (0.5)                         |
| Dribbling without ball (sec) | 12.3 (1.2)                        | 12.1 (1.1)                   | 12.4 (1.2)                       | 12.1 (1.1)                          | 12.0 (1.1)                        |
| Dribbling with ball (sec)    | 20.4 (2.3)                        | 20.2 (2.3)                   | 20.5 (2.2)                       | 20.2 (2.3)                          | 20.2 (2.3)                        |
| Sprint 5 m (sec)             | 1.19 (0.09)                       | 1.18 (0.09)                  | 1.21 (0.09)                      | 1.2 (0.1)                           | 1.2 (0.1)                         |
| Sprint 10 m (sec)            | 2.1 (0.1)                         | 2.0 (0.1)                    | 2.1 (0.1)                        | 2.0 (0.1)                           | 2.0 (0.1)                         |

|                                  |               |              |             |               |                |
|----------------------------------|---------------|--------------|-------------|---------------|----------------|
| Sprint 20 m (sec)                | 3.6 (0.3)     | 3.6 0.3      | 3.7 0.2     | 3.6 (0.3)     | 3.6 (0.3)      |
| Sprint 30 m (sec)                | 5.2 (0.4)     | 5.1 0.4      | 5.2 0.4     | 5.1 (0.4)     | 5.0 (0.4)      |
| Decay 30 m (sec)                 | 0.4 (0.2)     | 0.4 0.2      | 0.4 0.2     | 0.4 (0.2)     | 0.4 (0.2)      |
| YoYo IR test (m)                 | 944.4 (534.6) | 1030.5 589.7 | 871.8 472.1 | 973.5 (559.7) | 1082.6 (613.0) |
| Years of football experience (y) | 5.1 (3.8)     | 5.0 (4.0)    | 5.2 (3.6)   | 6.9 (2.0)     | 7.3 (2.0)      |

SD: standard deviation, y: year, cm: centimetre, kg: kilogram, m<sup>2</sup>: square meter, sec: seconds, #: number of, BMI: body mass index, PHV: peak height velocity, SBJ: standing broad jump, CMJ: counter movement jump, KTK3: 3 subtest short version of the Körperkoordinationstest für Kinder, YoYo IR1: YoYo Intermittent Recovery Test Level 1

**Table 2.** Injury characteristics.

|   | <b>N (%)</b> |
|---|--------------|
| <b>Total injuries</b>                     | 368 (100)    |
| <b><i>Mechanism</i></b>                   |              |
| Overuse injuries                          | 173 (47.0)   |
| Acute injuries                            | 195 (53.0)   |
| <b><i>Injury type</i></b>                 |              |
| Fracture                                  | 24 (6.5)     |
| Other bone and joint injuries             | 14 (3.8)     |
| Meniscus/cartilage                        | 1 (0.3)      |
| Sprain/ligament injury                    | 39 (10.6)    |
| Muscle rupture/tear/strain/cramps         | 116 (31.5)   |
| Tendon injury/rupture/tendinosis/bursitis | 100 (27.2)   |
| Haematoma/contusion/bruise                | 65 (17.7)    |
| Wound                                     | 1 (0.3)      |
| Concussion                                | 1 (0.3)      |
| <b><i>Injury location</i></b>             |              |
| Head/face                                 | 1 (0.3)      |
| Shoulder/clavicular                       | 8 (2.2)      |
| Upper arm                                 | 1 (0.3)      |
| Forearm                                   | 3 (0.8)      |
| Wrist                                     | 13 (3.5)     |
| Hand/finger/thumb                         | 12 (3.3)     |
| Sternum/ribs/upper back                   | 2 (0.5)      |
| Lower back/pelvis/sacrum                  | 12 (3.3)     |
| Hip/groin                                 | 72 (19.6)    |
| Thigh                                     | 63 (17.1)    |
| Knee                                      | 81 (22.0)    |
| Lower leg/Achilles tendon                 | 45 (12.2)    |
| Ankle                                     | 36 (9.8)     |
| Foot/toe                                  | 19 (5.2)     |
| <b><i>Injury severity</i></b>             |              |
| 0 days                                    | 7 (1.9)      |
| 1-3 days                                  | 62 (16.8)    |
| 4-7 days                                  | 95 (25.8)    |
| 8-28 days                                 | 136 (37.0)   |
| > 28 days                                 | 68 (18.5)    |