

Research Article

Vishvesh Kumar*

Pseudo-differential operators on homogeneous spaces of compact and Hausdorff groups

<https://doi.org/10.1515/forum-2018-0155>

Received July 1, 2018; revised August 16, 2018

Abstract: Let G be a compact Hausdorff group and let H be a closed subgroup of G . We introduce pseudo-differential operators with symbols on the homogeneous space G/H . We present a necessary and sufficient condition on symbols for which these operators are in the class of Hilbert–Schmidt operators. We also give a characterization of and a trace formula for the trace class pseudo-differential operators on the homogeneous space G/H .

Keywords: Pseudo-differential operators, Hilbert–Schmidt operators, trace class operators, homogeneous spaces of compact groups

MSC 2010: Primary 35S05, 47G30; secondary 43A85, 43A77

Communicated by: Christopher D. Sogge

1 Introduction

The study of pseudo-differential operators is started by Kahn and Nirenberg [17]. Pseudo-differential operators were used by Hörmander to study problems in partial differential operators [11]. Pseudo-differential operators on different classes of groups are extensively studied by several authors [2–4, 19, 22, 23]. Trace class pseudo-differential operators on \mathbb{S}^1 are studied by Delgado and Wong [4] and recently by Ghaemi et al. [10]. Molahajloo and Pirhayati [20] gave a characterization of and trace formula for trace class pseudo-differential operators with L^2 -symbols. Chen and Wong [1] considered trace class pseudo-differential operators on the unit sphere $\mathbb{S}^{n-1} \cong \text{SO}(n)/\text{SO}(n-1)$ centered in the origin in \mathbb{R}^n . In this paper, we try to replace $\text{SO}(n)$ by a compact Hausdorff group G and $\text{SO}(n-1)$ by a closed subgroup H of G . We study the Hilbert–Schmidt and trace class pseudo-differential operators on the homogeneous space G/H . The homogeneous spaces of compact groups play an important role in mathematical physics, geometric analysis, constructive approximation and coherent state transform, see [12–16, 18] and the references therein.

We consider pseudo-differential operators on homogeneous spaces of compact groups with L^2 -symbols. Pseudo-differential operators with L^2 -symbols are studied by many authors [3, 4, 10]. Recently, Ghaemi et al. [9] characterized nuclear pseudo-differential operators with L^2 -symbols on a compact Hausdorff group. By considering the L^p -conditions, $1 \leq p < \infty$, on symbols, we can allow singularities, and thus it becomes ideal for applications in several areas of mathematics, ranging from functional analysis to operator algebras or quantization. Our work can be considered a generalization of the corresponding work on \mathbb{S}^1 [4], compact Hausdorff groups [20], \mathbb{S}^{n-1} [1] and finite Abelian groups [21]. Our main aim in this paper is to give a charac-

*Corresponding author: Vishvesh Kumar, Department of Mathematics, Indian Institute of Technology Delhi, New Delhi 110 016, India, e-mail: vishveshmishra@gmail.com. <http://orcid.org/0000-0003-4799-9029>

terization of trace class and Hilbert–Schmidt pseudo-differential operators on the homogeneous space G/H of the compact group G , where H is a closed subgroup of G . We give a trace formula for trace class pseudo-differential operators. The main technique used here for obtaining the trace formula is to obtain a formula for the symbol of the product of two pseudo-differential operators on homogeneous spaces on compact groups.

In Section 2, we give some results from harmonic analysis on homogeneous spaces on compact groups [5–8] and operator theory to make the article self-contained. In Section 3, we define the pseudo-differential operator on the homogeneous space G/H of the compact and Hausdorff group G with the help of the Peter–Weyl theorem for G/H given by Ghani Farashahi [7]. We give a characterization of Hilbert–Schmidt pseudo-differential operators on G/H . We also give a product formula for pseudo-differential operators, a characterization of trace class pseudo-differential operators and then a trace formula for pseudo-differential operators on homogeneous spaces of compact groups.

2 Preliminaries

In this section, we recall some basic and important concepts of harmonic analysis on homogeneous spaces of compact groups, developed by Ghani Farashahi in a series of papers [5–8], and operator theory.

Let G be a compact Hausdorff group with normalized Haar measure dx , and let H be a closed subgroup of G with probability Haar measure dh .

The left coset space G/H can be seen as a homogeneous space with respect to the action of G on G/H given by left multiplication. The canonical surjection q from G to G/H is given by $q(x) := xH$. Let $\mathcal{C}(\Omega)$ denote the space of continuous functions on a compact Hausdorff space Ω . Define $T_H: \mathcal{C}(G) \rightarrow \mathcal{C}(G/H)$ by

$$T_H(f)(xH) = \int_H f(xh) dh, \quad xH \in G/H.$$

Then T_H is an onto map. The homogeneous space G/H has a unique normalized G -invariant positive Radon measure μ such that the Weil formula

$$\int_{G/H} T_H(f)(xH) d\mu(xH) = \int_G f(x) dx$$

holds.

Let (π, \mathcal{H}_π) be a continuous unitary representation of a compact group G on a Hilbert space \mathcal{H}_π . It is well known that any irreducible representation (π, \mathcal{H}_π) is finite-dimensional with the dimension d_π (say). Let (π, \mathcal{H}_π) be a continuous unitary representation of a compact group G . Consider the operator-valued integral

$$T_H^\pi := \int_H \pi(h) dh$$

defined in the weak sense, i.e.,

$$\langle T_H^\pi u, v \rangle = \int_H \langle \pi(h)u, v \rangle dh \quad \text{for all } u, v \in \mathcal{H}_\pi.$$

Note that the function $h \mapsto \langle \pi(h)u, v \rangle$ is in $L^1(H)$ for all $u, v \in \mathcal{H}_\pi$. Therefore, the integral $\int_H \langle \pi(h)u, v \rangle dh$ is an ordinary integral of an L^1 -function. Hence, T_H^π is a bounded linear operator on \mathcal{H}_π with norm bounded by one. Further, T_H^π is a partial isometric orthogonal projection, and T_H^π is an identity operator if and only if $\pi(h) = I$ for all $h \in H$. Denote the set of all continuous irreducible unitary representation on G by \widehat{G} . Set

$$\mathcal{K}_\pi^H = \{u \in \mathcal{H}_\pi : \pi(h)u = u \text{ for all } h \in H\}.$$

Then \mathcal{K}_π^H is a closed subspace of \mathcal{H}_π . Let $d_{\pi,H}$ be the dimension of \mathcal{K}_π^H . Then it is evident that $d_{\pi,H} \leq d_\pi$. It can be easily seen that $d_{\pi,H} = d_\pi$ if and only if $[\pi] \in H^\perp := \{[\pi] \in \widehat{G} : \pi(h) = I \text{ for all } h \in H\}$.

Definition 2.1. Let H be a closed subgroup of a compact group G . Then the dual object $\widehat{G/H}$ of G/H is a subset of \widehat{G} and is given by

$$\widehat{G/H} := \{[\pi] \in \widehat{G/H} : T_H^\pi \neq 0\} = \left\{ [\pi] \in \widehat{G} : \int_H \pi(h) dh \neq 0 \right\}.$$

Definition 2.2. Let H be a closed subgroup of a compact group G , and let $[\pi] \in \widehat{G/H}$. An ordered orthonormal basis $\mathfrak{B} = \{e_j\}_{j=1}^{d_\pi}$ of the Hilbert space \mathcal{H}_π is called H -admissible if it is an extension of an orthonormal basis of $\{e_j\}_{j=1}^{d_{\pi,H}}$ of the closed subspace \mathcal{K}_π^H .

Now, we state a corollary of the Peter–Weyl theorem for homogeneous spaces of a compact group and some of its consequences.

Theorem 2.3 ([7, Corollary 4.1]). *Let H be a closed subgroup of a compact group G , and let μ be the normalized G -invariant measure on G/H . For each $[\pi] \in \widehat{G/H}$, let $\mathfrak{B}_\pi = \{e_{\ell,\pi} : 1 \leq \ell \leq d_\pi\}$ be an H -admissible basis for the representation space \mathcal{H}_π . Then we have the following statements:*

- (i) *The set $\mathfrak{B}(G/H) = \{\sqrt{d_\pi} \pi_{ij} : \pi \in \widehat{G/H}, 1 \leq i \leq d_\pi, 1 \leq j \leq d_{\pi,H}\}$ constitutes an orthonormal basis for the Hilbert space $L^2(G/H)$.*
- (ii) *Each $f \in L^2(G/H)$ decomposes as the following:*

$$f = \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{j=1}^{d_{\pi,H}} \sum_{i=1}^{d_\pi} \langle f, \pi_{ij} \rangle_{L^2(G/H)} \pi_{ij},$$

where the series converge in $L^2(G/H)$.

Using the above decomposition of $f \in L^2(G/H)$ and the orthogonality relation

$$\langle \pi_{ij}, \pi'_{kl} \rangle_{L^2(G/H)} = d_\pi^{-1} \delta_{\pi\pi'} \delta_{ik} \delta_{jl},$$

we get the following Plancherel's theorem (see [5]).

Theorem 2.4. *For $f \in L^2(G/H)$, we have*

$$\|f\|_{L^2(G/H)}^2 = \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{j=1}^{d_{\pi,H}} \sum_{i=1}^{d_\pi} |\langle f, \pi_{ij} \rangle_{L^2(G/H)}|^2.$$

Let \mathcal{H} be a complex and separable Hilbert space with the inner product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|_{\mathcal{H}}$. Denote by $\|\cdot\|_{\mathcal{B}(\mathcal{H})}$ the norm in the C^* -algebra of all bounded linear operators on \mathcal{H} .

An operator $T \in \mathcal{B}(\mathcal{H})$ is a Hilbert–Schmidt operator if for any (hence all) orthonormal basis $\{e_j\}_{j=1}^\infty$ of \mathcal{H} , we have $\sum_j \|Te_j\|_{\mathcal{H}} < \infty$. The set of Hilbert–Schmidt operators, denoted by S_2 , is a two-sided ideal of $\mathcal{B}(\mathcal{H})$. The Hilbert–Schmidt norm on S_2 is given by

$$\|T\|_{HS} = \left(\sum_{j=1}^\infty \|Te_j\|_{\mathcal{H}}^2 \right)^{\frac{1}{2}}.$$

An operator $T \in \mathcal{B}(\mathcal{H})$ is a trace class operator if for any (hence all) orthonormal basis $\{e_j\}_{j=1}^\infty$ of \mathcal{H} , we have $\text{tr}(T) = \sum_{j=1}^\infty \langle Te_j, e_j \rangle < \infty$. The set of all trace class operators on \mathcal{H} is denoted by S_1 .

The following well-known theorem describes a relation between a trace class operator and Hilbert–Schmidt operators.

Theorem 2.5. *Let $T \in \mathcal{B}(\mathcal{H})$. Then T is a trace class operator if and only if there exist two Hilbert–Schmidt operators U and V on \mathcal{H} such that $T = UV$.*

3 Pseudo-differential operators on homogeneous space of compact groups

Throughout this section, we always assume that G is a compact Hausdorff group, and H is a closed subgroup of G .

In this section, we assume that the homogeneous space G/H has a unique G -invariant positive Radon measure μ , and $\widehat{G/H}$ is the abstract dual of G/H . The inner product on $L^2(G/H)$ will be denoted by $\langle \cdot, \cdot \rangle$. We will freely use the notation and concepts explained in the previous section. Now, we start with the definition of pseudo-differential operators.

Let σ be a measurable function on $G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N}$. Define the *pseudo-differential operator* T_σ corresponding to the symbol σ as follows. For any measurable function f on G/H , define $T_\sigma f$ formally on G/H by

$$(T_\sigma f)(gH) = \sum_{[\pi] \in G/H} d_\pi \sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} \sigma(gH, \pi, i, j) \langle f, \pi_{ij} \rangle \pi_{ij}(gH) \quad \text{for almost all } gH \in G/H,$$

where $\langle \cdot, \cdot \rangle$ denotes the L^2 -inner product.

Let $L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})$ denote the space of all measurable functions σ on $G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N}$ such that

$$\|\sigma\|_{L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})} := \left(\sum_{[\pi] \in G/H} d_\pi \sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} \int_{G/H} |\sigma(gH, \pi, i, j) \pi_{ij}(gH)|^2 d\mu(gH) \right)^{\frac{1}{2}} < \infty.$$

The following theorem gives the boundedness of pseudo-differential operators on G/H with corresponding symbols in L^2 -space.

Theorem 3.1. *Let G be a compact group, and let H be a closed subgroup of G . Let $\sigma \in L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})$. Then the pseudo-differential operator $T_\sigma: L^2(G/H) \rightarrow L^2(G/H)$ is bounded and*

$$\|T_\sigma\|_{\mathcal{B}(L^2(G/H))} \leq \|\sigma\|_{L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})}.$$

Proof. Let $f \in L^2(G/H)$. Then, by Minkowski's inequality, we have

$$\begin{aligned} \|T_\sigma f\|_{L^2(G/H)} &= \left(\int_{G/H} |(T_\sigma f)(gH)|^2 d\mu(gH) \right)^{\frac{1}{2}} \\ &= \left(\int_{G/H} \left| \sum_{[\pi] \in G/H} d_\pi \sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} \sigma(gH, \pi, i, j) \langle f, \pi_{ij} \rangle \pi_{ij}(gH) \right|^2 d\mu(gH) \right)^{\frac{1}{2}} \\ &\leq \sum_{[\pi] \in G/H} \left(\int_{G/H} d_\pi^2 \left| \sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} \sigma(gH, \pi, i, j) \langle f, \pi_{ij} \rangle \pi_{ij}(gH) \right|^2 d\mu(gH) \right)^{\frac{1}{2}}. \end{aligned}$$

Using Cauchy–Schwarz inequality, we get

$$\begin{aligned} \|T_\sigma f\|_{L^2(G/H)} &\leq \sum_{[\pi] \in G/H} \left(\int_{G/H} d_\pi^2 \left(\sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} |\langle f, \pi_{ij} \rangle|^2 \right) \left(\sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} |\sigma(gH, \pi, i, j) \pi_{ij}(gH)|^2 \right) d\mu(gH) \right)^{\frac{1}{2}} \\ &= \sum_{[\pi] \in G/H} \left(d_\pi \sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} |\langle f, \pi_{ij} \rangle|^2 \right)^{\frac{1}{2}} \left(\int_{G/H} d_\pi \sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} |\sigma(gH, \pi, i, j) \pi_{ij}(gH)|^2 d\mu(gH) \right)^{\frac{1}{2}} \\ &\leq \left(\sum_{[\pi] \in G/H} d_\pi \sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} |\langle f, \pi_{ij} \rangle|^2 \right)^{\frac{1}{2}} \left(\sum_{[\pi] \in G/H} d_\pi \int_{G/H} \sum_{i=1}^{d_{\pi,H}} \sum_{j=1}^{d_\pi} |\sigma(gH, \pi, i, j) \pi_{ij}(gH)|^2 d\mu(gH) \right)^{\frac{1}{2}}. \end{aligned}$$

Therefore, Plancherel's theorem gives

$$\|T_\sigma f\|_{L^2(G/H)} \leq \|f\|_{L^2(G/H)} \|\sigma\|_{L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})}.$$

Hence, T_σ is a bounded operator on $L^2(G/H)$ and $\|T_\sigma\|_{\mathcal{B}(L^2(G/H))} \leq \|\sigma\|_{L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})}$. \square

Our next result provides a characterization of Hilbert–Schmidt pseudo-differential operators on G/H .

Theorem 3.2. *Let G be a compact group, and let H be a closed subgroup of G . Let σ be a measurable function on $G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N}$. Then the corresponding pseudo-differential operator $T_\sigma : L^2(G/H) \rightarrow L^2(G/H)$ is a Hilbert–Schmidt operator if and only if $\sigma \in L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})$. Furthermore, we have*

$$\|T_\sigma\|_{HS} = \|\sigma\|_{L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})}.$$

Proof. Let $\pi' \in \widehat{G/H}$ and $1 \leq i_0 \leq d_\pi$, $1 \leq j_0 \leq d_{\pi,H}$. Then, by the relation $\langle \pi'_{lk}, \pi_{l'k'} \rangle_{L^2(G/H)} = d_\pi^{-1} \delta_{ll'} \delta_{kk'} \delta_{\pi\pi'}$, we get

$$\begin{aligned} (T_\sigma \pi'_{i_0 j_0})(gH) &= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \sigma(gH, \pi, i, j) \langle \pi'_{i_0 j_0}, \pi_{i,j} \rangle \pi_{ij}(gH) \\ &= \sigma(gH, \pi', i_0, j_0) \pi_{i_0 j_0}(gH), \quad gH \in G/H. \end{aligned}$$

Since $\{\sqrt{d_\pi} \pi_{ij} : 1 \leq d_\pi, 1 \leq j \leq d_{\pi,H}\}$ forms an orthonormal basis for $L^2(G/H)$, we have

$$\begin{aligned} \|T_\sigma\|_{HS}^2 &= \sum_{[\pi] \in \widehat{G/H}} \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \|T_\sigma(\sqrt{d_\pi} \pi_{ij})\|_{L^2(G/H)}^2 \\ &= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \int_{G/H} |(T_\sigma \pi_{ij})(gH)|^2 d\mu(gH) \\ &= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \int_{G/H} |\sigma(gH, \pi, i, j) \pi_{ij}(gH)|^2 d\mu(gH) \\ &= \|\sigma\|_{L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})}^2. \end{aligned}$$

Therefore, T_σ is Hilbert–Schmidt operator if and only if $\sigma \in L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})$. \square

Let σ and τ be two measurable functions on $\widehat{G/H} \times G/H \times \mathbb{N} \times \mathbb{N}$. Define $\sigma \otimes \tau$ by

$$\begin{aligned} (\sigma \otimes \tau)(gH, \xi, k, l) &= \int_{G/H} \tau(wH, \xi, k, l) \xi_{kl}(wH) \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \sigma(gH, \pi, i, j) \overline{\pi_{ij}(wH)} \pi_{ij}(gH) d\mu(wH) (\xi_{kl}(gH))^{-1} \end{aligned}$$

for all $gH \in G/H$, $[\xi] \in \widehat{G/H}$, $1 \leq k \leq d_\pi$ and $1 \leq l \leq d_{\pi,H}$.

In the following theorem, we prove that the product of two pseudo-differential operators on G/H is again a pseudo-differential operator on G/H .

Theorem 3.3. *Let G be a compact group, and let H be a closed subgroup of G . Let σ and τ be measurable functions on $G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N}$. Then*

$$T_\sigma T_\tau = T_\lambda,$$

where $\lambda = \sigma \otimes \tau$.

Proof. For $f \in L^2(G/H)$ and $gH \in G/H$, we have

$$\begin{aligned} (T_\sigma T_\tau f)(gH) &= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \sigma(gH, \pi, i, j) \langle T_\tau f, \pi_{ij} \rangle \pi_{ij}(gH) \\ &= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \sigma(gH, \pi, i, j) \left(\int_{G/H} (T_\tau f)(wH) \overline{\pi_{ij}(wH)} d\mu(wH) \right) \pi_{ij}(gH) \\ &= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \sigma(gH, \pi, i, j) \\ &\quad \times \left(\int_{G/H} \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\pi,H}} \tau(wH, \xi, k, l) \langle f, \xi_{kl} \rangle \xi_{kl}(wH) \overline{\pi_{ij}(wH)} d\mu(wH) \right) \pi_{ij}(gH) \end{aligned}$$

$$\begin{aligned}
&= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \sigma(gH, \pi, i, j) \\
&\quad \times \left(\int_{G/H} \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\xi,H}} \tau(wH, \xi, k, l) \langle f, \xi_{kl} \rangle \xi_{kl}(wH) \overline{\pi_{ij}(wH)} d\mu(wH) \right) \pi_{ij}(gH) \\
&= \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\xi,H}} \int_{G/H} \tau(wH, \xi, k, l) \xi_{kl}(wH) \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \sigma(gH, \pi, i, j) \\
&\quad \times \overline{\pi_{ij}(wH)} \pi_{ij}(gH) d\mu(wH) (\xi_{kl}(gH))^{-1} \langle f, \xi_{kl} \rangle \xi_{kl}(gH) \\
&= \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\xi,H}} \left\{ \int_{G/H} \tau(wH, \xi, k, l) \xi_{kl}(wH) \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \sigma(gH, \pi, i, j) \right. \\
&\quad \left. \times \overline{\pi_{ij}(wH)} \pi_{ij}(gH) d\mu(wH) (\xi_{kl}(gH))^{-1} \right\} \langle f, \xi_{kl} \rangle \xi_{kl}(gH) \\
&= \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\xi,H}} \lambda(gH, \xi, k, l) \langle f, \xi_{kl} \rangle \xi_{kl}(gH) \\
&= (T_\lambda f)(gH),
\end{aligned}$$

where $\lambda(gH, \xi, k, l) = (\sigma \otimes \tau)(gH, \xi, k, l)$ defined as above.

Hence, $T_\sigma T_\tau = T_\lambda$. □

Finally, we present the trace formula for a pseudo-differential operator on G/H .

Theorem 3.4. *Let G be a compact group, and let H be a closed subgroup of G . Let λ be a measurable function on $G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N}$. Then T_λ is a trace class operator if and only if there exist two measurable functions σ and τ in $L^2(G/H \times \widehat{G/H} \times \mathbb{N} \times \mathbb{N})$ such that $\lambda = \sigma \otimes \tau$. Furthermore,*

$$\begin{aligned}
\text{tr}(T_\lambda) &= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\pi,H}} \int_{G/H} \lambda(gH, \pi, i, j) |\pi_{ij}(gH)|^2 d\mu(gH) \\
&= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\pi,H}} \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\xi,H}} \langle \tau(\cdot, \xi, i, j) \xi_{ij}, \pi_{kl} \rangle \langle \tau(\cdot, \pi, k, l) \pi_{kl}, \xi_{ij} \rangle.
\end{aligned}$$

Proof. The first part of the theorem follows from Theorem 3.2 and the fact that the product of two Hilbert–Schmidt operators is a trace class operator (see Theorem 2.5). Now, the absolute convergence of the series

$$\sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\pi,H}} \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\xi,H}} \langle \tau(\cdot, \xi, i, j) \xi_{ij}, \pi_{kl} \rangle \langle \tau(\cdot, \pi, k, l) \pi_{kl}, \xi_{ij} \rangle$$

follows from Plancherel’s theorem and Cauchy–Schwarz inequality.

Since the set $\{\sqrt{d_\pi} \pi_{ij} : \pi \in \widehat{G/H}, 1 \leq i \leq d_\pi, 1 \leq j \leq d_{\pi,H}\}$ forms an orthonormal basis for $L^2(G/H)$, we have

$$\begin{aligned}
\text{tr}(T_\lambda) &= \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\xi,H}} \langle T_\lambda \xi_{ij}, \xi_{ij} \rangle \\
&= \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\xi,H}} \int_{G/H} \lambda(gH, \xi, i, j) \xi_{ij}(gH) \overline{\xi_{ij}(gH)} d\mu(gH) \\
&= \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{i=1}^{d_\pi} \sum_{j=1}^{d_{\xi,H}} \int_{G/H} \left\{ \int_{G/H} \tau(wH, \xi, i, j) \xi_{ij}(wH) \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\pi,H}} \sigma(gH, \pi, k, l) \right. \\
&\quad \left. \times \overline{\pi_{kl}(wH)} \pi_{kl}(gH) d\mu(wH) \right\} \overline{\xi_{ij}(gH)} d\mu(gH)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\pi,H}} \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{i=1}^{d_\xi} \sum_{j=1}^{d_{\xi,H}} \int_{G/H} \tau(wH, \xi, i, j) \xi_{ij}(wH) \overline{\pi_{jk}(wH)} d\mu(wH) \\
&\quad \times \int_{G/H} \sigma(gH, \pi, k, l) \pi_{kl}(gH) \overline{\xi_{gH}} d\mu(gH) \\
&= \sum_{[\pi] \in \widehat{G/H}} d_\pi \sum_{k=1}^{d_\pi} \sum_{l=1}^{d_{\pi,H}} \sum_{[\xi] \in \widehat{G/H}} d_\xi \sum_{i=1}^{d_\xi} \sum_{j=1}^{d_{\xi,H}} \langle \tau(\cdot, \xi, i, j) \xi_{ij}, \pi_{kl} \rangle \langle \tau(\cdot, \pi, k, l) \pi_{kl}, \xi_{ij} \rangle. \quad \square
\end{aligned}$$

Acknowledgment: The author expresses his deep gratitude to Prof. M. W. Wong for several insightful conversations. He thanks his supervisors Ritumoni Sarma and N. Shraavan Kumar for their support and encouragement. He would also like to thank the referee for his/her valuable suggestions.

Funding: The author wishes to thank the Council of Scientific and Industrial Research, India, for Senior Research Fellowship (09/086(1204)/2014-EMR-I).

References

- [1] Z. Chen and M. W. Wong, Traces of pseudo-differential operators on \mathbb{S}^{n-1} , *J. Pseudo-Differ. Oper. Appl.* **4** (2013), no. 1, 13–24.
- [2] A. Dasgupta and M. W. Wong, Pseudo-differential operators on the affine group, in: *Pseudo-Differential Operators: Groups, Geometry and Applications*, Trends Math., Birkhäuser/Springer, Cham (2017), 1–14.
- [3] J. Delgado and M. Ruzhansky, Schatten classes and traces on compact groups, *Math. Res. Lett.* **24** (2017), no. 4, 979–1003.
- [4] J. Delgado and M. W. Wong, L^p -nuclear pseudo-differential operators on \mathbb{Z} and \mathbb{S}^1 , *Proc. Amer. Math. Soc.* **141** (2013), no. 11, 3935–3942.
- [5] A. Ghaani Farashahi, Abstract operator-valued Fourier transforms over homogeneous spaces of compact groups, *Groups Geom. Dyn.* **11** (2017), no. 4, 1437–1467.
- [6] A. Ghaani Farashahi, Abstract Plancherel (trace) formulas over homogeneous spaces of compact groups, *Canad. Math. Bull.* **60** (2017), no. 1, 111–121.
- [7] A. Ghani Farashahi, Peter–Weyl theorem for homogeneous spaces of compact groups, *Int. J. Anal. Appl.* **13** (2017), no. 1, 22–31.
- [8] A. Ghaani Farashahi, Trigonometric polynomials over homogeneous spaces of compact groups, *Adv. Oper. Theory* **2** (2017), no. 1, 87–97.
- [9] M. B. Ghaemi, M. Jamalpourbirgani and M. W. Wong, Characterizations, adjoints and products of nuclear pseudo-differential operators on compact and Hausdorff groups, *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.* **79** (2017), no. 4, 207–220.
- [10] M. B. Ghaemi, M. Jamalpour Birgani and M. W. Wong, Characterizations of nuclear pseudo-differential operators on \mathbb{S}^1 with applications to adjoints and products, *J. Pseudo-Differ. Oper. Appl.* **8** (2017), no. 2, 191–201.
- [11] L. Hörmander, *The Analysis of Linear Partial Differential Operators. III*, Grundlehren Math. Wiss. 274, Springer, Berlin, 1985.
- [12] A. A. Kirillov and A. D. Gvishiani, *Theorems and problems of functional analysis* (in Russian), 2nd ed., Nauka, Moscow, 1988.
- [13] V. V. Kisil, Relative convolutions. I. Properties and applications, *Adv. Math.* **147** (1999), no. 1, 35–73.
- [14] V. V. Kisil, Erlangen program at large: An overview, in: *Advances in Applied Analysis*, Trends Math., Birkhäuser/Springer, Basel (2012), 1–94.
- [15] V. V. Kisil, *Geometry of Möbius Transformations*, Imperial College Press, London, 2012.
- [16] V. V. Kisil, Calculus of operators: covariant transform and relative convolutions, *Banach J. Math. Anal.* **8** (2014), no. 2, 156–184.
- [17] J. J. Kohn and L. Nirenberg, An algebra of pseudo-differential operators, *Comm. Pure Appl. Math.* **18** (1965), 269–305.
- [18] R. L. Lipsman, Non-Abelian Fourier analysis, *Bull. Sci. Math. (2)* **98** (1974), no. 4, 209–233.
- [19] S. Molahajloo, K. A. Okoudjou and G. E. Pfander, Boundedness of multilinear pseudo-differential operators on modulation spaces, *J. Fourier Anal. Appl.* **22** (2016), no. 6, 1381–1415.
- [20] S. Molahajloo and M. Pirhayati, Traces of pseudo-differential operators on compact and Hausdorff groups, *J. Pseudo-Differ. Oper. Appl.* **4** (2013), no. 3, 361–369.

- [21] S. Molahajloo and K. L. Wong, Pseudo-differential operators on finite Abelian groups, *J. Pseudo-Differ. Oper. Appl.* **6** (2015), no. 1, 1–9.
- [22] M. W. Wong, *Wavelet Transforms and Localization Operators*, Oper. Theory Adv. Appl. 136, Birkhäuser, Basel, 2002.
- [23] M. W. Wong, *An Introduction to Pseudo-differential Operators*, 3rd ed., Ser. Anal. Appl. Comput. 6, World Scientific, Hackensack, 2014.