

Anyone who has never made a mistake has never tried anything new.

-Albert Einstein

Promotors:

- Prof. Dr. ir. Gijs Du Laing

Department of Green Chemistry and Technology, Faculty of Bioscience Engineering,
Ghent University, Belgium

- Prof. Dr. ir. Piet Lens

UNESCO-IHE Institute for Water Education, Delft, The Netherlands

- Prof. Dr. ir. Ivet Ferrer

Department of Civil and Environmental Engineering, Universitat Politècnica de
Catalunya, Barcelona, Spain

Members of the examination committee:

- Prof. Stefaan De Smet

Department of Animal Sciences and Aquatic Ecology, Faculty of Bioscience
Engineering, Ghent University, Belgium

- Prof. Diederik Rousseau

Department of Green Chemistry and Technology, Faculty of Bioscience Engineering,
Ghent University, Belgium

- Prof. Dongli Liang

Faculty of Natural Resources and Environment, Northwest A&F University, Shaanxi,
China

- Dr. Gary Bañuelos

USDA-ARS, Water Management Research Unit, California, United States

- Dr. Rubén Díez-Montero

Department of Civil and Environmental Engineering, Universitat Politècnica de
Catalunya BarcelonaTech, Spain

Dean of the faculty of Bioscience Engineering, Ghent University

Prof. dr. ir. Marc Van Meirvenne

Rector of Ghent University

Prof. dr. ir. Rik Van de Walle



Selenium and zinc enriched bioproducts generated from wastewater as micronutrient feed supplements and biofertilizers

Jun Li

2021

Thesis submitted in fulfillment of the requirements for the degree of Doctor in Bioscience Engineering, Ghent University & Environmental Engineering, Universitat Politècnica de Catalunya

Dutch translation of the title:

Selenium en zink verrijkte bioproducten gegenereerd uit afvalwater als micronutriënt voedersupplementen en biomeststoffen

Cover illustration:

Picture took in Gent by Jun Li and electron diagram of selenium.

ISBN: 9789463574143

Please refer this work to:

Li, J. (2021). Selenium and zinc enriched bioproducts generated from wastewater as micronutrient feed supplements and biofertilizers. Ph.D. dissertation, Faculty of Bioscience Engineering, Ghent University, Belgium & Department of Civil and Environmental Engineering, Universitat Politècnica de Catalunya, Barcelona, Spain.

© 2021 by Jun Li.

The author and the promoters give the authorization to consult and to copy parts of this work for personal use only. Every other use is subject to the copyright laws. Permission to reproduce any material contained in this work should be obtained from the author.

Table of Contents

Table of Contents

Summary	7
Samenvatting.....	11
Resumen	15
List of abbreviations.....	19
Chapter 1 General introduction	23
1.1 Motivation	24
1.2 Objectives	26
1.3 Outline of the dissertation	27
Chapter 2 Literature review	29
2.1 Occurrence of Se.....	30
2.2 Se toxicity and deficiency.....	32
2.3 Approaches for addressing Se deficiency - biofortification	36
2.4 Approaches for addressing Zn deficiency – biofortification.....	40
2.5 Se/Zn-enriched organic materials as biofertilizers for biofortification	41
2.6 Local “green” production of micronutrient-enriched bioproducts by micronutrient removal from wastewater	43
Chapter 3 Production of selenium- and zinc-enriched <i>Lemna</i> and <i>Azolla</i> as potential micronutrient-enriched bioproducts.....	47
Abstract.....	48
Keywords	48
3.1 Introduction	49
3.2 Materials and methods.....	50
3.2.1 Experimental materials	50
3.2.2 Experimental design	51
3.2.3 Analytical methods.....	51
3.2.4 Statistical analysis.....	53
3.3 Results.....	54
3.3.1 Plant growth	54
3.3.2 Se concentration in <i>Lemna</i> and <i>Azolla</i>	57
3.3.3 Se accumulation and removal by <i>Lemna</i> and <i>Azolla</i>	58
3.3.4 BCF _{Se}	59
3.3.5 Zn concentration in <i>Lemna</i> and <i>Azolla</i>	63
3.3.6 Zn accumulation, removal and BCF _{Zn}	64
3.3.7 Protein content.....	64

Table of Contents

3.3.8 Se speciation in the plants.....	65
3.4 Discussion	67
3.4.1 Se uptake and toxic effects of Se and Zn on <i>Lemna</i> and <i>Azolla</i>	67
3.4.2 Simultaneous uptake of Se and Zn	69
3.4.3 Potential of <i>Lemna</i> and <i>Azolla</i> for wastewater treatment and production of micronutrient-enriched bioproducts	71
3.4.4 Study limitations and future perspectives	73
3.5 Conclusions	74
Supplementary Information	75
Chapter 4 Valorization of Se-enriched sludge and duckweed generated from wastewater as micronutrient biofertilizers	83
Abstract.....	84
Keywords.	84
4.1 Introduction	85
4.2 Materials and methods.....	86
4.2.1 Soil collection and characterization.....	86
4.2.2 Crops screening experiment.....	88
4.2.3 Valorization of Se-enriched biomaterials as Se biofertilizers	89
4.2.4 Analytical methods.....	91
4.2.5 Estimated daily intake and health risk assessment of the bean seeds.....	92
4.2.6 Statistical analysis.....	93
4.3 Results	93
4.3.1 Crops screening.....	93
4.3.2 Characterization of the produced Se-enriched biomaterials	96
4.3.3 Se released to non-planted soils amended with the Se-enriched biomaterials	98
4.3.4 Se released to planted soils amended with Se-enriched biomaterials.....	99
4.3.5 pH, TC and TOC contents in the soil pore water	108
4.3.6 Evaluation of the Se-enriched biomaterials produced from wastewater as slow-release fertilizers for Se biofortification.....	109
4.4 Discussion	110
4.4.1 Se toxicity and accumulation in different crops.....	110
4.4.2 Selenium release from Se-enriched duckweed and sludge	111
4.4.3 The potential of Se-enriched duckweed and sludge as Se biofertilizers ..	112
4.4.4 Study limitations and future perspectives	115
4.5 Conclusions	116

Table of Contents

Supplementary Information	118
Chapter 5 Biofortification of green beans with Se and Zn-enriched duckweed and sludge	123
Abstract.....	124
Keywords	124
5.1 Introduction	125
5.2 Materials and methods.....	125
5.2.1 Soil collection and characterization	125
5.2.2 Valorization of SeZn-enriched biomaterials as Se and Zn biofertilizers ...	125
5.2.3 Analytical methods.....	127
5.2.4. Statistical analysis.....	128
5.3 Results and discussion.....	128
5.3.1 Characterization of SeZn-enriched biomaterials.....	128
5.3.2 pH and TOC content of the extracted soil pore water	128
5.3.3 Selenium content in the soil pore water during plant growth.....	130
5.3.4 Zinc content in the soil pore water during beans growth	132
5.3.5 Biomass of beans	134
5.3.6 Selenium content in the beans grown on soils amended with SeZn-enriched biomaterials	136
5.3.7 Zinc content in the beans grown on soils amended with SeZn-enriched biomaterials	138
5.3.8 Se species in the beans.....	140
5.4 Conclusions	143
Chapter 6 Production of selenium-enriched microalgae as potential feed supplement in high rate algal ponds treating domestic wastewater	145
Abstract.....	146
Keywords:	146
6.1 Introduction	147
6.2 Materials and methods.....	148
6.2.1 Source of biomass and wastewater	148
6.2.2 Se removal by microalgae in batch experiments	149
6.2.3 Se removal in HRAPs and production of Se-enriched biomass.....	150
6.2.4 Wastewater characterization in the HRAP systems	151
6.2.5 Nutritional parameters of the microalgae	151
6.2.6 Pathogenic bacterial content	153
6.2.7 Statistical analysis.....	153

Table of Contents

6.3 Results and discussion.....	154
6.3.1 Microalgae growth, Se accumulation and Se removal in batch experiments	154
6.3.2 Wastewater treatment efficiency and Se removal in HRAPs	157
6.3.3 Nutritional value of microalgae grown in HRAPs	161
6.3.4 Pathogenic bacterial content	170
6.4 Conclusions	171
Supplementary Information	173
Chapter 7 Production of Se-enriched microalgae in raceway ponds treating domestic wastewater as biostimulant and biofertilizer	177
Abstract.....	178
Keywords	178
7.1 Introduction	179
7.2 Materials and methods.....	180
7.2.1 Se-enriched microalgae production in a pilot-scale raceway pond.....	180
7.2.2 Preparation of liquid Se-enriched microalgae extracts	181
7.2.3 Germination test with Se-enriched microalgae extract.....	181
7.2.4 Beans production with foliar spray or soil drench of Se-enriched microalgae extract.....	183
7.2.5 Beans production on soil amendment with Se-enriched microalgae biomass	184
7.2.6 Analytical methods.....	184
7.2.7 Statistical analysis.....	185
7.3 Results and discussion.....	185
7.3.1 Effect of Se-enriched microalgae extract on seed germination and growth of bean seedlings	185
7.3.2 Effect of foliar and soil drench application of Se-enriched microalgae extract on the bean plant growth and selenium concentration in the bean plants	190
7.3.3 Effect of Se-enriched microalgae biomass on the bean plant growth and Se concentration in the plants	193
7.3.4 Practical implications of using Se-enriched microalgae biomass and its extract as biostimulants and Se sources.....	199
7.4 Conclusions	200
Supplementary Information	201
Chapter 8 General discussion, conclusions and future perspectives	209
8.1 Production of micronutrient-enriched bioproducts as potential feed/food supplements.....	210

Table of Contents

8.1.1 Production and valorization of Se/Zn-enriched duckweed as potential feed supplements	210
8.1.2 Production and valorization of Se-enriched microalgae as potential feed supplements	211
8.2 Valorisation of the produced micronutrient-enriched bioproducts as Se/Zn biofertilizers.....	212
8.2.1 Agronomic biofortification of <i>Phaseolus vulgaris</i> with Se/Zn-enriched duckweed and sludge	212
8.2.2 Valorization of Se-enriched microalgae and their extract as potential Se biofertilizers and biostimulants.....	213
8.3 Comparison of Se utilization rates in duckweed, microalgae and sludge based production systems.....	214
8.4 Risks related to the use of Se-enriched biomaterials generated from wastewater as micronutrient biofertilizer and feed supplement	215
8.5 Future perspectives.....	217
8.5.1 Production of micronutrient-rich bioproducts	217
8.5.2 Biofortification.....	219
8.5.3 Feasibility analysis	220
References	221
Acknowledgement	247
Curriculum Vitae	249

Summary

Summary

Selenium (Se) is an essential micronutrient for humans and animals with a narrow window between deficiency and toxicity levels. Dietary Se intake of humans and animals is not adequate in some regions. On the other hand, Se toxicity also frequently occurs worldwide, due to water or soil contamination, as Se is widely applied in or released from industrial and agricultural activities. The trace element zinc (Zn) is also often present in too low concentrations in agricultural soils, but is also toxic at elevated concentrations.

Improvement of the dietary Se and Zn intake through enrichment of food and feed crops (named biofortification) is currently being explored as a possible solution for Se and Zn deficiency. Supplementation of feed and food products with Se and Zn is another solution. In biofortification, the application of conventional chemical Se/Zn fertilizers to increase the Se/Zn content in crops could result in secondary soil and water contamination due to the low utilization efficiency of Se/Zn and fast leaching. Slow-release Se/Zn-enriched fertilizers may therefore be beneficial. Moreover, the use of Se/Zn originating from primary mining for the production of Se/Zn enriched-feed/food supplements is not considered economically and environmental-friendly, taking into account that external Se/Zn is being used and the excess chemicals are currently being discharged as waste. It may thus be beneficial from an economic and environmental point of view to produce slow release Se/Zn-enriched biofertilizers or Se/Zn-enriched feed supplements locally from Se/Zn-bearing water while partially cleaning the water. This may contribute to the worldwide drive for resource recovery and circular economy. Therefore, this thesis aimed to explore the potential of Se/Zn-enriched bioproducts produced from wastewater treatment processes by eco-technologies (phytoextraction, bioreduction and microalgae-based systems) as Se/Zn feed supplements and biofertilizers.

Chapter 1 and **Chapter 2** present the motivation, objectives and background information on the occurrence of Se and Zn in human and animal diets and their deficiency and toxicity for humans and animals. Current studies regarding micronutrient biofortification and the production of Se and Zn supplements to tackle micronutrient deficiency are discussed. This is followed by a discussion on the

Summary

paradigm shift from waste treatment to resource recovery, highlighting the potential of biobased technologies for micronutrient recovery from wastewater, while producing micronutrient-enriched feed/food supplements and biofertilizers.

Since Se can replace sulfur in amino acids and Zn can also be complexed by functional groups in proteins, protein-rich plants have an immense potential for the bioaccumulation/biofortification of these micronutrients. Thus, two aquatic plants (*Lemna* and *Azolla*) with substantial protein content were applied in **Chapter 3** to evaluate the possibility of Se and Zn bioaccumulation/removal from wastewater while producing micronutrient-enriched dietary proteins (for feed/food supplements) and biofertilizers. Nutrient-medium spiked with different concentrations of Se and Zn was used to mimic wastewater. Results of **Chapter 3** demonstrated that both *Lemna* and *Azolla* can accumulate high levels of Se and Zn, while they take up around 10 times more Se(IV) than Se(VI) from the medium. Besides, high transformation to organic Se forms and accumulation in plants after taking up Se(IV), together with the high protein content and fast growth rate, makes *Lemna* (also named duckweed later on) and *Azolla* good candidates for the production of Se- and Zn-enriched biomass, which can be used as crop fertilizers or protein-rich food/feed supplements or ingredients. Considering that a synergetic effect between Se and Zn in *Lemna*, but an antagonistic effect in *Azolla* was observed in **Chapter 3**, *Lemna* loaded with Se/Zn was selected for the subsequent experiments in **Chapter 4** and **Chapter 5**.

Subsequently, **Chapter 4** and **Chapter 5**, respectively, evaluated the valorization potential of the produced micronutrient-enriched duckweed as well as sludge generated in wastewater treatment processes containing single Se or Se combined with Zn as micronutrient biofertilizers. This was conducted in pot experiments using green beans (*Phaseolus vulgaris*). Micronutrient-enriched sludge dominated by the presence of Se in zero oxidation state (Se(0)) was found to be the preferred slow-release Se biofertilizer and an effective Se source to produce Se-enriched beans for Se-deficient populations. This was motivated by the higher Se bioavailability and lower organic carbon content released into the soil from micronutrient-enriched sludge in comparison with micronutrient-enriched duckweed, enabling a higher soil Se supply. The remarkably higher organic carbon content in the soil could result in Se immobilization. On the contrary, the Zn content in the seeds of beans was not successfully improved through the application of micronutrient-enriched biofertilizers in

Summary

comparison with the control. This could be attributed to the lower Zn translocation rate from plant roots to seeds, and the lower Zn amount applied into soils as Se/Zn-enriched biomaterials.

Additionally, microalgae have a great capacity to assimilate/remove excess nutrients from the corresponding growth medium (or wastewater) and metabolize them into valuable compounds such as protein, fatty acids, vitamins and carbohydrates. Microalgae are thus a potential protein source to substitute common animal and plant proteins (e.g. soybean). **Chapter 6** explored the potential of Se removal in high rate algal ponds (HRAPs) treating domestic wastewater, while producing high-value Se-enriched biomass that may be used as feed supplement (dietary protein) or biofertilizer. Results indicated that the wastewater treatment performance of the HRAPs was effective. The produced Se-enriched microalgae in HRAPs fed with domestic wastewater contained a high content of crude protein (48% of volatile suspended solids) and the selenoamino acid selenomethionine (SeMet) (91% of total Se). Besides, the essential amino acid content of the microalgae was comparable to that of soybean, an animal feed protein. This Chapter also highlighted that Se may potentially induce the production of the polyunsaturated fatty acids omega-3 (ω 3) and omega-6 (ω 6), and eicosapentaenoic (EPA) in microalgae, although further research is still needed to confirm this. Therefore, the production of Se-enriched microalgae in HRAPs may offer a promising alternative for upgrading low-value recovered resources into high-value feed supplements.

Chapter 7 aimed to evaluate the Se-enriched microalgae generated in Chapter 6 as a potential biostimulant and/or biofertilizer to enhance crop growth and improve the Se content of the crops. Raw Se-enriched microalgal biomass and an extract thereof were applied in the production of green beans (*Phaseolus vulgaris*) through soil and foliar application. This study demonstrated that the application of raw Se-enriched microalgae biomass to soil (1-10%, soil application) and its extract to leaves (1%, foliar spray) enhanced plant growth, which confirmed that Se-enriched microalgae act as a biostimulant. Besides, a higher Se content in the plant and soil (for soil application) was achieved after the application of Se-enriched microalgae or its extract. This indicated that Se-enriched microalgae cultivated during wastewater treatment can be valorized as a biostimulant and biofertilizer to improve both the seed yields and Se content of beans, leading to a higher market value of the beans.

Summary

Chapter 8 concluded and discussed the key findings of this thesis, taking also economic aspects into account. It also highlighted the limitations of the study. The whole thesis contributes to offering an environmentally friendly and sustainable way for micronutrient biofortification/supplementation in Se/Zn-deficient areas, while recovering nutrients from wastewater.

Samenvatting

Samenvatting

Selenium (Se) is een essentieel micronutriënt voor mens en dier met een nauw bereik tussen deficiëntie en toxiciteit. De inname van Se via de voeding van mens en dier is in sommige regio's niet voldoende. Anderzijds komt Se-toxiciteit ook wereldwijd veel voor als gevolg van water- of bodemverontreiniging, aangezien Se veel wordt toegepast in of vrijkomt bij industriële en agrarische activiteiten. Het sporenelement zink (Zn) komt ook vaak in te lage concentraties voor in landbouwbodems, maar is bij verhoogde concentraties giftig.

Verbetering van de dieetopname van Se en Zn door verrijking van voedsel- en voedergewassen, biofortificatie, wordt momenteel onderzocht als een mogelijke oplossing voor Se- en Zn-deficiëntie. Suppletie van voeder en voedingsmiddelen met Se en Zn is een andere oplossing. Bij biofortificatie kan de toepassing van conventionele chemische Se/Zn-meststoffen om het Se/Zn-gehalte in gewassen te verhogen, leiden tot secundaire bodem- en waterverontreiniging vanwege de lage benuttingsgraad van Se/Zn en de snelle uitspoeling. Se/Zn-verrijkte meststoffen met vertraagde afgifte kunnen daarom nuttig zijn. Bovendien wordt het gebruik van Se/Zn afkomstig uit de primaire mijnbouw voor de productie van met Se/Zn verrijkte diervoeders en voedingssupplementen niet als economisch en milieuvriendelijk beschouwd, rekening houdend met het feit dat extern Se/Zn wordt gebruikt en de overvloedige chemicaliën momenteel als afval worden geloosd. Het kan dus vanuit economisch en ecologisch oogpunt voordelig zijn om Se/Zn-verrijkte biomeststoffen met vertraagde afgifte of Se/Zn-verrijkte voedingssupplementen lokaal te produceren uit Se/Zn-houdende afvalwaters, terwijl het afvalwater gereinigd wordt. Dit kan bijdragen aan het wereldwijde streven naar terugwinning van hulpbronnen en circulaire economie. Daarom was dit proefschrift gericht op het onderzoeken van het potentieel van Se/Zn-verrijkte bioproducten, geproduceerd uit afvalwaterbehandelingsprocessen door middel van eco-technologieën (fyto-extractie, bioreductie en op microalgen gebaseerde methoden), als Se/Zn-voedingssupplementen en biomeststoffen.

Hoofdstuk 1 en **Hoofdstuk 2** geven de motivatie, doelstellingen en achtergrondinformatie over het voorkomen van Se en Zn in de voeding van mens en dier en hun deficiëntie en toxiciteit voor mens en dier. Huidige studies met betrekking tot de biofortificatie van micronutriënten en de productie van Se- en Zn-supplementen

Samenvatting

om het tekort aan micronutriënten aan te pakken, worden besproken. Dit wordt gevolgd door een discussie over de paradigmaverschuiving van afvalverwerking naar terugwinning van hulpbronnen, waarbij het potentieel van biogebaseerde technologieën voor het terugwinnen van micronutriënten uit afvalwater wordt belicht, terwijl met micronutriënten verrijkte voeder-/voedingssupplementen en biomeststoffen worden geproduceerd.

Omdat Se het element zwavel in aminozuren kan vervangen en Zn ook kan worden gecomplexeerd door functionele groepen in eiwitten, hebben eiwitrijke planten een enorm potentieel voor de bioaccumulatie/biofortificatie van deze micronutriënten. Zo werden in **Hoofdstuk 3** twee waterplanten, *Lemna* en *Azolla*, met een substantieel eiwitgehalte, gebruikt om de mogelijkheid van bioaccumulatie/verwijdering van Se en Zn uit afvalwater te evalueren tijdens de productie van met micronutriënten verrijkte voedingseiwitten (voor voeder-/voedingssupplementen) en biomeststoffen. Nutriënten-medium verrijkt met verschillende concentraties Se en Zn werd gebruikt om afvalwater na te bootsen. De resultaten van **Hoofdstuk 3** toonden aan dat zowel *Lemna* als *Azolla* hoge niveaus van Se en Zn kunnen accumuleren, terwijl ze ongeveer 10 keer meer Se(IV) dan Se(VI) uit het medium opnemen. Bovendien maakt de hoge transformatie naar organische Se-vormen en hoge Se accumulatie in planten na het opnemen van Se(IV), samen met het hoge eiwitgehalte en de snelle groeisnelheid, *Lemna* (later eendenkroos genoemd) en *Azolla* goede kandidaten voor de productie van Se- en Zn-verrijkte biomassa, die kan worden gebruikt als gewasbemesting of als eiwitrijk voeder-/voedingssupplementen. Gezien een synergetisch effect tussen Se en Zn in *Lemna*, maar een antagonistisch effect in *Azolla* werd waargenomen in **Hoofdstuk 3**, werd *Lemna* verrijkt met Se/Zn geselecteerd voor de daaropvolgende experimenten in **Hoofdstuk 4** en **Hoofdstuk 5**.

Vervolgens hebben respectievelijk **Hoofdstuk 4** en **Hoofdstuk 5** het valorisatiepotentieel geëvalueerd van het geproduceerde eendenkroos en slib dat wordt gegenereerd in afvalwaterzuiveringsprocessen die enkelvoudig Se of Se gecombineerd met Zn bevatten als micronutriënten biomeststoffen. Dit werd uitgevoerd in potproeven met sperziebonen (*Phaseolus vulgaris*). Met micronutriënten verrijkt slib, gedomineerd door de aanwezigheid van Se in oxidatietoestand nul (Se (0)), bleek de preferentiële Se-biomeststof met langzame afgifte en een effectieve Se-bron te zijn om Se-verrijkte gewassen te produceren voor Se-deficiënte populaties. Dit werd

Samenvatting

gemotiveerd door de hogere biologische beschikbaarheid van Se en het lagere organische koolstofgehalte dat in de bodem vrijkomt uit het met micronutriënten verrijkte slib, waardoor een hogere Se-aanvoer in de bodem mogelijk is in vergelijking met het met micronutriënten verrijkte eendenkroos. Het opmerkelijk hogere gehalte aan organische koolstof in de bodem kan leiden tot immobilisatie van Se. Daarentegen werd het Zn-gehalte in de zaden van bonen niet succesvol verbeterd door de toepassing van met micronutriënten verrijkte biomaterialen in vergelijking met de controle. Dit kan worden toegeschreven aan de lagere translocatie van Zn van plantenwortels naar zaden, en de lagere toegepaste hoeveelheid Zn in bodems als met Se/Zn verrijkte biomaterialen.

Microalgen hebben een groot vermogen om overtollige voedingsstoffen uit het overeenkomstige groeimedium (of afvalwater) te assimileren/verwijderen en deze om te zetten in waardevolle componenten zoals eiwitten, vetzuren, vitamines en koolhydraten. Microalgen zijn dus een potentiële eiwitbron ter vervanging van gewone dierlijke en plantaardige eiwitten (bijv. soja). **Hoofdstuk 6** onderzocht de mogelijkheid van Se-verwijdering door middel van intensieve algenvijvers (HRAPs) die huishoudelijk afvalwater behandelen en tegelijkertijd waardevolle Se-verrijkte biomassa produceren die kan worden gebruikt als een mogelijk voedersupplement (voedingseiwit) of biomeststof. De resultaten gaven aan dat de performantie van de HRAPs op het gebied van afvalwaterbehandeling effectief was. De geproduceerde met Se verrijkte microalgen in HRAPs gevoed met huishoudelijk afvalwater bevatten een hoog gehalte aan ruw eiwit (48% vluchtige gesuspendeerde vaste stof) en het selenoaminozuur selenomethionine (SeMet) (91% van totaal Se). Bovendien was het essentiële aminozuurgehalte van de microalgen vergelijkbaar met dat van sojabonen, een diervoederproteïne. Dit hoofdstuk benadrukte ook dat Se mogelijks de productie van de meervoudig onverzadigde vetzuren omega-3 (ω 3) en omega-6 (ω 6) en eicosapentaeenzuur (EPA) in microalgen kan induceren, hoewel verder onderzoek nodig is om dit te bevestigen. Daarom kan de productie van met Se verrijkte microalgen in HRAPs een veelbelovend alternatief bieden voor het upgraden van laagwaardige afvalstromen tot hoogwaardige voedersupplementen.

Hoofdstuk 7 had als doel de met Se verrijkte microalgen die in Hoofdstuk 6 werden gegenereerd te evalueren als een potentieel biostimulant en/of biomeststof om de gewasgroei te bevorderen en het Se-gehalte van de gewassen te verbeteren. Ruwe

Samenvatting

Se-verrijkte microalgenbiomassa en extracten daarvan werden toegepast bij de productie van sperziebonen (*Phaseolus vulgaris*) via bodem- en bladtoepassing. Deze studie toonde aan dat de toepassing van ruwe Se-verrijkte microalgenbiomassa op de bodem (1-10%, bodemtoepassing) en een extract ervan op bladeren (1%, bladspray) de plantengroei verbeterde, wat bevestigde dat met Se verrijkte microalgen werken als een biostimulant. Bovendien werd een hoger Se-gehalte in de plant en bodem (voor bodemtoepassing) bereikt na het aanbrengen van met Se verrijkte microalgen of een extract daarvan. Dit gaf aan dat Se-verrijkte microalgen die tijdens de afvalwaterzuivering worden gekweekt, kunnen worden gevaloriseerd als biostimulant en biomeststof om de opbrengst van zaden en het Se-gehalte van bonen samen te verbeteren, wat leidt tot een hogere marktwaarde van de bonen.

Hoofdstuk 8 concludeert en bediscussieert de belangrijkste bevindingen van dit proefschrift, tevens economische aspecten in overweging nemend. Het benadrukt ook de beperkingen van het onderzoek. Dit proefschrift draagt bij aan de ontwikkeling van een milieuvriendelijke en duurzame manier voor biofortificatie en supplementatie van micronutriënten in Se/Zn-deficiënte gebieden, terwijl nutriënten worden teruggewonnen uit afvalwater.

Resumen

Resumen

El selenio (Se) es un micronutriente esencial para los humanos y animales, con poco margen entre los niveles de deficiencia y toxicidad. Actualmente, la ingesta de Se por parte de humanos y animales no es adecuada en algunas regiones. Por otro lado, la toxicidad por Se también ocurre con frecuencia en todo el mundo, debido a la contaminación del agua o del suelo, ya que el Se se aplica o se libera ampliamente en actividades industriales y agrícolas. El oligoelemento zinc (Zn) también suele estar presente en concentraciones demasiado bajas en suelos agrícolas, pero también es tóxico en concentraciones elevadas.

Actualmente se está estudiando la mejora de la ingesta de Se y Zn mediante el enriquecimiento de cultivos alimenticios y piensos (denominado biofortificación) como una posible solución para la deficiencia de Se y Zn. La suplementación de piensos y productos alimenticios con Se y Zn es otra solución. En cuanto a la biofortificación, la aplicación de fertilizantes químicos convencionales de Se/Zn para aumentar el contenido de Se/Zn en los cultivos podría causar la contaminación secundaria del suelo y del agua debido a la baja tasa de utilización de Se/Zn y su rápida lixiviación. Por lo tanto, la aplicación de fertilizantes enriquecidos con Se/Zn de liberación lenta sería más adecuada. Además, el uso de Se/Zn procedente de la minería para la producción de piensos/complementos alimenticios enriquecidos con Se/Zn no se considera ni económico ni respetuoso con el medio ambiente, teniendo en cuenta que se está utilizando Se/Zn externo y que los productos químicos excedentarios actualmente se desechan como un residuos. Por lo tanto, sería más beneficioso, desde un punto de vista económico y ambiental, producir localmente biofertilizantes enriquecidos con Se/Zn de liberación lenta o suplementos alimenticios enriquecidos con Se/Zn a partir de aguasresidual que contenga Se/Zn, a la vez que se trata el aguasresidual. Esto contribuiría al impulso mundial de la recuperación de recursos y la economía circular. Por lo tanto, el objetivo de esta tesis fue explorar el potencial de los bioproductos enriquecidos con Se/Zn, producidos a partir del tratamiento de aguasresiduales mediante eco-tecnologías (fitoextracción, biorreducción y sistemas basados en microalgas) como complementos alimenticios y biofertilizantes enriquecidos en Se/Zn.

Resumen

En el **Capítulo 1** y el **Capítulo 2** se presenta la motivación, los objetivos y antecedentes en cuanto a la presencia de Se y Zn en las dietas de humanos y animales, y su deficiencia y toxicidad para humanos y animales. Se discuten estudios actuales sobre la biofortificación de micronutrientes y la producción de suplementos de Se y Zn para abordar la deficiencia de micronutrientes. A esto le sigue una discusión sobre el cambio de paradigma del tratamiento de residuos a la recuperación de recursos, destacando el potencial de las biotecnologías para la recuperación de micronutrientes de las aguas residuales, al tiempo que se producen suplementos alimenticios/piensos y biofertilizantes enriquecidos con micronutrientes.

Dado que el Se puede reemplazar el azufre en los aminoácidos y el Zn también puede formar complejos con grupos funcionales de las proteínas, las plantas ricas en proteínas tienen un inmenso potencial para la bioacumulación/biofortificación de estos micronutrientes. Por lo tanto, en el **Capítulo 3** se utilizaron dos plantas acuáticas (*Lemna* y *Azolla*) con un contenido sustancial de proteínas para evaluar la bioacumulación/eliminación de Se y Zn del agua residual, a la vez que se producían proteínas enriquecidas con micronutrientes (para piensos/complementos alimenticios). Se usó un medio de cultivo enriquecido con diferentes concentraciones de Se y Zn para simular el agua residual. Los resultados del Capítulo 3 demostraron que tanto *Lemna* como *Azolla* pueden acumular altos niveles de Se y Zn, mientras que absorben alrededor de 10 veces más Se(IV) que Se(VI) del medio. Además, la alta transformación a formas orgánicas de Se y la acumulación en las plantas después de absorber Se(IV), junto con el alto contenido de proteínas y la rápida tasa de crecimiento, hacen que *Lemna* (también llamada lenteja de agua en adelante) y *Azolla* sean buenas candidatas para la producción de biomasa enriquecida con Se y Zn, que se puede utilizar como biofertilizante para cultivos o como suplemento alimenticio/pienso rico en proteínas. Considerando el efecto sinérgico entre Se y Zn que se observó para *Lemna*, y antagónico para *Azolla*, se seleccionó *Lemna* enriquecida con Se/Zn para los siguientes experimentos del **Capítulo 4** y **Capítulo 5**.

Así, en el **Capítulo 4** y el **Capítulo 5** se evaluó el potencial de valorización de la lenteja de agua, así como de los lodos de depuradora, con Se simple o Se combinado con Zn, como biofertilizantes de micronutrientes. Esto se llevó a cabo con experimentos en macetas utilizando judías verdes (*Phaseolus vulgaris*) como cultivo. El lodo enriquecido con micronutrientes, con Se en estado de oxidación cero (Se (0)),

Resumen

resultó ser el biofertilizante de Se de liberación lenta más adecuado, siendo una fuente de Se eficaz para producir habas enriquecidas en Se para poblaciones deficientes en Se. Esto fue resultado de la mayor biodisponibilidad del Se y el menor contenido de carbono orgánico liberado al suelo en el caso de los lodos, lo que permitió un mayor suministro de Se al suelo, en comparación con la lenteja de agua enriquecida con micronutrientes. Esto se debe a que un contenido de carbono orgánico notablemente más alto en el suelo puede resultar en la inmovilización del Se. Por el contrario, el contenido de Zn en las semillas de judía verde no se mejoró mediante la aplicación de biofertilizantes enriquecidos con micronutrientes en comparación con el control. Esto podría atribuirse a la menor tasa de translocación del Zn de las raíces de las plantas a las semillas y a la menor cantidad de Zn aplicada al suelo como biofertilizante enriquecido en Se/Zn.

Por otro lado, las microalgas tienen una gran capacidad para asimilar/eliminar nutrientes del medio de cultivo (o de las aguas residuales) y metabolizarlos en componentes valiosos como proteínas, ácidos grasos, vitaminas y carbohidratos. Por lo tanto, las microalgas se están considerando actualmente como una fuente potencial de proteínas para sustituir las proteínas animales y vegetales habituales (por ejemplo, la soja). En el Capítulo 6 se investigó el potencial de eliminación de Se en lagunas de alta carga (HRAP por sus siglas en inglés) para el tratamiento de aguas residuales domésticas, y la producción de biomasa microalgal enriquecida con Se, que podría usarse como complemento alimenticio (proteína) o biofertilizante de alto valor añadido. Los resultados indicaron un tratamiento eficaz de las aguas residuales en las HRAP. Las microalgas enriquecidas con Se producidas en las HRAP alimentadas con agua residual doméstica contenían un alto contenido de proteína cruda (48% de sólidos volátiles en suspensión) y el selenoaminoácido SeMet (selenometionina) (91% del Se total). Además, el contenido de aminoácidos esenciales de las microalgas era comparable al de la soja, una proteína típica de la alimentación animal. Este Capítulo también mostró que el Se puede inducir la producción de ácidos grasos poliinsaturados omega-3 ($\omega 3$) y omega-6 ($\omega 6$) y eicosapentaenoico (EPA) en las microalgas, aunque se necesitaría más investigación para confirmarlo. Por lo tanto, la producción de microalgas enriquecidas en Se en HRAP se presenta como una alternativa prometedora para recuperar recursos de bajo valor y convertirlos en suplementos alimenticios de alto valor.

Resumen

El **Capítulo 7** tuvo por objetivo evaluar las microalgas enriquecidas en Se generadas en el Capítulo 6 como bioestimulante para mejorar el crecimiento de las plantas y como biofertilizante de Se para mejorar el contenido de Se de las plantas. Se aplicaron tanto biomasa de microalgas enriquecida con Se (cruda), como extractos de la misma, en judías verdes (*Phaseolus vulgaris*) a nivel de suelo y foliar. Este estudio mostró que la aplicación de biomasa de microalgas enriquecidas con Se al suelo (1-10%) y sus extractos a las hojas (1%, aspersión foliar) mejoró el crecimiento de las plantas, lo que confirmó que las microalgas enriquecidas con Se actúan como un bioestimulante. Además, se logró incrementar el contenido de Se en la planta y el suelo (para la aplicación al suelo) tras la aplicación de microalgas enriquecidas con Se y de extractos de las mismas. Esto indica que las microalgas enriquecidas con Se, cultivadas mediante el tratamiento de aguas residuales, se pueden valorizar como bioestimulante y biofertilizante para mejorar tanto el rendimiento de semillas como el contenido de Se de las habas, incrementando el valor de mercado de las mismas.

En el **Capítulo 8** se muestran las conclusiones y se discuten los principales resultados de esta tesis. También se enumeran las limitaciones del estudio. Esta tesis contribuye al desarrollo sostenible y respetuoso con el medio ambiente de la biofortificación/suplementación de micronutrientes en áreas deficientes en Se/Zn, junto con la recuperación de nutrientes de las aguas residuales.

List of abbreviations

List of abbreviations

BCF	Bioconcentration factor
BSA	Bovine standard albumin
CEC	Cation exchange capacity
COD _{tot}	Total chemical oxygen demand
COD _{sol}	Soluble chemical oxygen demand
DI	Deionized water
DM	Dry matter
DMDSe	Dimethyl diselenide
DMSe	Dimethyl selenide
DO	Dissolved oxygen
DW1	Se enriched duckweed application at 1.0 mg Se/kg soil
DW5	Se enriched duckweed application at 5.0 mg Se/kg soil
EC	Electrical conductivity
EDI	Estimated daily intake
EFSA	European Food Safety Authority
EPA	Eicosapentaenoic
EXAFS	X-ray absorption fine structure
FAO	Food and Agriculture Organization
GI	Germination index
GP	Germination percentage
HPLC	High performance liquid chromatography
HRAPs	High rate algal ponds
HRAP-Se	High rate algal ponds with continuous Na ₂ SeO ₃ spiking
HRAP-C	High rate algal ponds without Se spiking
HRI	Health risk index
HRT	Hydraulic retention time
ICP-MS	Inductively coupled plasma-mass spectrometry
IDA	Iodoacetic acid
Me-SeMet	Methylselenomethionine
MGT	Mean germination time
MTL	Maximum tolerable levels
MUFA	Monounsaturated fatty acids

List of abbreviations

OM	Organic matter
OPA	O-phthaldialdehyde
PUFA	Polyunsaturated fatty acids
RfD	Reference oral dose
ROS	Reactive oxygen species
SBM	Soybean meal
SDS	Sodium dodecyl sulfate
Se	Selenium
Se(IV)	Selenite
Se(VI)	Selenate
SeCys	Selenocysteine
SeCys ₂	Selenocystine
SeMet	Selenomethionine
SeMetSeCys	Se-methyl-selenocysteine
SeZnDW1	Se and Zn enriched duckweed application at 1.0 mg Se/kg soil
SeZnDW5	Se and Zn enriched duckweed application at 5.0 mg Se/kg soil
SeZnSL1	Se and Zn enriched sludge application at 1.0 mg Se/kg soil
SeZnSL5	Se and Zn enriched sludge application at 5.0 mg Se/kg soil
SL1	Se enriched sludge application at 1.0 mg Se/kg soil
SL5	Se enriched sludge application at 5.0 mg Se/kg soil
SFA	Total saturated fatty acids
SP	Soluble phosphorus
SVI	Seedling vigor index
TC	Total carbon
TKN	Total Kjeldahl nitrogen
TN	Total nitrogen
TOC	Total organic carbon
TP	Total phosphorus
TSS	Total suspended solids
VSS	Volatile suspended solids
ω3	Omega-3
ω6	Omega-6
WHO	World Health Organization

List of abbreviations

XANES	X-ray absorption near edge structure
Zn	Zinc

Chapter 1

Chapter 1 General introduction

Chapter 1

1.1 Motivation

Selenium (Se) and zinc (Zn) are essential trace elements, playing a crucial role in the functioning of enzymes in humans and animals and protecting cells from damage by free radicals (Hatfield et al., 2014). Selenoproteins, i.e. proteins containing selenium, are also well known as antioxidants and catalysts for the production of the active thyroid hormone. Although Se is not essential for the growth and survival of plants, it is considered as a beneficial element for plants as well, which can enhance resistance to stress, whereas Zn is an essential element for plants, that can enhance plant growth (Feng et al., 2013; Subramanyam et al., 2019). The recommended daily Se and Zn intake in an adult human diet are 0.04–0.4 mg and 15 mg per person per day, respectively (FAO/WHO, 2001). However, despite the importance of these trace elements, intake of Se and Zn by animals and humans in a wide range of countries, including e.g. Belgium, the United Kingdom and Kenya, is currently still quite low, resulting in Se and Zn deficiency and causing negative health effects, including increased risk of mortality, poor immune function, and cognitive decline (Broadley et al., 2006; Broadley et al., 2007; Rayman, 2012; Roekens et al., 1986). An estimated one billion people around the world are affected by selenium deficiency and more than 30% of the world's population is Zn-deficient because of low Se and Zn intake (Poblaciones & Rengel, 2017; Rayman, 2004). Besides, also farm animals (Dermauw et al., 2013) and pets (van Zelst et al., 2016) can be affected by Se and Zn deficiencies, leading to economic losses. Therefore, the Se and Zn content in the human and animal diet is a topic of interest to public health systems around the world (Lavu et al., 2012).

Food/feed biofortification (e.g. enrichment of food and feed crops with Se and Zn), food/feed supplements or dietary diversification are proposed as possible solutions to remediate these micronutrient deficiencies. For biofortification, inorganic Se forms (e.g., selenite and selenate) and Zn (Zn^{2+}) are generally added into soils to enhance the micronutrients in crops in order to achieve an optimal Se/Zn level in human and animal diets. However, the direct application of inorganic Se/Zn forms into soils could result in secondary soil and water contamination due to the fast leaching (high mobilization) of the fresh applied Se/Zn and the low utilization efficiency of Se/Zn by plants (Wang et al., 2018). It may thus be beneficial and environmental-friendly to produce slow-release Se/Zn-enriched fertilizers through ecotechnologies in order to supply Se/Zn to soil for plant uptake, such as Se/Zn-enriched organic fertilizer. Likewise, many chemicals

Chapter 1

originating from primary mining (not only Se/Zn but also other micro- and macro-nutrients) are used to cultivate microorganisms for the production of Se/Zn-enriched food/feed supplements, such as the production of Se-yeast (Rayman, 2004) and Se-algae (Gómez-Jacinto et al., 2020). This is not cost-effective and could even introduce new contaminations. The production of Se/Zn-enriched food/feed supplements or biofertilizers could be made more sustainable and environment-friendly by producing them using low-value nutrients in wastewaters as feedstock, while cleaning the wastewater.

Excess of Se and Zn in the environment is also frequently observed. Se and Zn are likely to co-occur in waste streams or in the environment as a result of both industrial and agricultural activities, such as in petroleum refinery effluents (Wake, 2005), in the groundwater of uranium mill tailings repositories (ca. 0.2-0.4 mg Se/L and ca.1-2 mg Zn/L) (Morrison et al., 2002), in leachates of Zn mining (3-8 mg Zn/L) (Etteieb et al., 2020; Mishra, 2014) and gold mining (1.7-3.3 mg Se/L) (Tan et al., 2016), and in agricultural runoff or aquaculture wastewater after application of Se (0.14-1.4 mg Se/L) and Zn (ca. 2-3 mg/L) minerals for improving crop yields or nutritional quality of the crops (Han et al., 2020; Mao et al., 2014; Tan et al., 2016). Selenium exists in these waters mainly as Se(IV) and Se(VI) oxyanions, which have a high solubility. These wastewaters may therefore serve as potential Se and Zn sources, and it may be beneficial to recover not only macronutrients, but also Se and Zn from these wastewaters through different biological technologies (e.g., phytoextraction, bioreduction or microalgae-based methods). Accordingly, bioproducts with high Se/Zn contents could be generated by these technologies, and these may be valorized as slow-release organic Se and Zn fertilizers or feed supplements. Besides, environmentally sustainable fertilizers and feed supplements may also be produced from wastewaters with a low Se/Zn content as feedstock, whereafter Se/Zn may be added to produce an added-value product. Such alternative micronutrient-enriched fertilizers or feed supplements will contribute to the sustainability of our food and feed production systems, as they can be produced on wastewater recovering nutrients during its secondary or tertiary treatment.

Chapter 1

1.2 Objectives

The main objective of this thesis is to explore the potential of Se/Zn-enriched bioproducts generated from wastewater as environmental-friendly micronutrient feed supplements and biofertilizers. Specific research questions being addressed are:

(1) what is the efficiency of different bio-based methods, in particular phytoextraction, bioreduction and microalgae-based methods, for Se/Zn removal from wastewater?

(2) can high Se/Zn-containing bioproducts be generated from the bio-based wastewater treatment processes using the wastewater as feedstock?

(3) can Se/Zn-enriched biomaterials produced from wastewater be used as value-added Se/Zn-containing feed supplement? Can they accumulate high contents of beneficial and bioavailable Zn and Se species, as well as other compounds of high nutritional value?

(4) can Se/Zn-enriched biomaterials be applied to soils or leaves as Se/Zn biofertilizer and/or biostimulant to improve crop growth and increase Se/Zn concentrations in the crops? What are the impacts of soil or foliar application of these materials on crops growth and on total and bioavailable Se and Zn in the crops?

Therefore, specifically, the thesis is aimed to:

- Develop sustainable and biobased methods which are able to remove and recover Se/Zn from aqueous solutions, meanwhile producing micronutrient-enriched biomaterials. Phytoextraction by aquatic plants, bioreduction by sludge and microalgae-based wastewater treatment are investigated in this thesis.
- Evaluate the produced Se/Zn-enriched duckweed and Se-enriched microalgae for potential use as micronutrient feed supplements by evaluating Se/Zn accumulation, Se speciation, Se bioaccessibility and other nutritional compounds in these biomaterials (e.g., amino acids and fatty acids).
- Assess the applicability of the produced Se/Zn-enriched duckweed, sludge and microalgae as potential slow-release organic Se/Zn fertilizers or biostimulants for enhancement of Se/Zn accumulation by plants and stimulating plant growth in pot experiments. The kinetics of the micronutrient release from these biomaterials into soils and the uptake of the trace elements by the crops are

Chapter 1

investigated. The effects of their application on biomass growth and accumulation and speciation of Zn and Se in seeds are also explored.

1.3 Outline of the dissertation

The remainder of this thesis is composed of seven chapters and mainly focuses on Se because Se was less studied so far compared to Zn. Zn was included only in two chapters, studying interaction with Se, considering that Se and Zn are likely to co-exist in the environment and simultaneous occurrence of nutritional deficiencies of more than one micronutrient is more common. Multi-mineral enriched supplements and agronomic biofortification of crops are thus being explored as a possible way to alleviate multi-micronutrient deficiency (Mao et al., 2014).

Chapter 2 presents the background of the doctoral research and a literature review on the occurrence of Se and Zn in human and animal diets and their deficiency and toxicity for humans and animals. The current status on micronutrient biofortification and production of micronutrient feed/food supplements is discussed. This is followed by a discussion on the paradigm shift from waste treatment to resource recovery, highlighting the potential of biobased technologies for micronutrient recovery from wastewater, while producing micronutrient-enriched feed/food supplements and biofertilizers.

Chapter 3 evaluates the possibility to produce Se and Zn-enriched aquatic plants, *Lemna* (duckweed) and *Azolla*, as potential micronutrient-enriched feed supplements or biofertilizers while cleaning water. In Chapters 4 and 5, the potential to use single Se (Chapters 4) or simultaneous Se and Zn enriched (Chapters 5) duckweed (generated by phytoextraction from Chapter 3) and sludge (produced by a bioreduction process) as Se/Zn biofertilizers in the agronomic biofortification of green beans (*Phaseolus vulgaris*) was assessed, respectively. Other researchers have previously reported the ability of sludge to precipitate Se from wastewater under anaerobic reducing conditions and thus the possibility to produce Se-enriched sludge (Staicu et al., 2015a; Staicu et al., 2015b). Therefore, we did not anymore focus on the production of the Se/Zn enriched sludge in this thesis, but only evaluated its potential use as biofertilizer in Chapters 4 and 5.

Chapter 1

Chapter 6 investigates the possibility to produce Se-enriched microalgae in high-rate algal ponds treating domestic wastewater as high-value Se-enriched feed supplements, meanwhile cleaning the wastewater. Chapter 7 evaluates the Se-enriched microalgae obtained from Chapter 6 as potential Se biofertilizers and biostimulants for the enhancement of Se uptake by beans and growth of the beans. A scheme of the doctoral research is presented in Fig. 1.1. A general discussion, integrating also economic aspects, and conclusions drawn from this study are presented in Chapter 8, together with ideas and perspectives for future work.




Figure 1.1. Schematic representation of the doctoral thesis research

Chapter 2

Chapter 2 Literature review

Occurrence, toxicity and deficiency of selenium and zinc and novel biofortification strategies

This chapter was redrafted after:

Li, J., Du Laing, G., Ferrer, I., Lens, P.N.L. 2021. Selenium biofortification for human and animal nutrition, In: Lens, P.N.L., Pakshirajan, K., (Eds.) *Environmental Technologies to Treat Selenium Pollution - Principles and Engineering*. IWA Publishing (In press).

Chapter 2

2.1 Occurrence of Se

Selenium (Se) was identified as a new substance in 1817 by Jöns Jacob Berzelius when studying a method to produce sulfuric acid from sulfur-bearing rocks (Fernández-Martínez & Charlet, 2009). It is a metalloid element belonging to the oxygen group (Group 16 of the periodic table) and closely allied in chemical and physical properties with the elements sulfur and tellurium (Lenz & Lens, 2009). It exists in inorganic forms as selenate (SeO_4^{2-}), selenite (SeO_3^{2-}), selenide (Se^{2-}) and elemental Se (Se^0), and in organic forms such as selenocysteine (SeCys) and selenomethionine (SeMet). Due to this complex chemical behavior, Se is found in all natural materials on earth: soil, rocks, waters, air, plants and animals (Fordyce, 2007). During a long period, Se has been identified as a dangerous substance because of its toxicity (Fordyce et al., 2000). Afterwards, it has also been recognized as an essential trace element for human and animal health due to its crucial role in the functioning of enzymes for humans and animals (Fordyce, 2013; Rayman, 2000). Both Se deficiency and excess in the environment can cause adverse effects on organisms, Se is therefore considered as a “double-edged sword” (Fordyce, 2013). Moreover, the range between its deficiency and toxicity is very narrow. For example, to avoid deficiency, Se in human dietary consumption should exceed 40 $\mu\text{g}/\text{day}$ and concentrations in soil should exceed 0.2 mg/kg to result in sufficiently high Se concentrations in food crops and animal products. To not exceed toxic levels, human dietary consumption cannot exceed 400 $\mu\text{g}/\text{day}$ and concentrations in the soil cannot exceed 3 mg/kg Se to avoid excess Se concentrations in food and feed crops (Dinh et al., 2018). This results in the occurrence of global “hotspots” wherein both Se deficiency and toxicity regularly occur, depending on the Se concentration present in the soil. For instance, Se intoxication events have been regularly reported in the Enshi region of China, where a maximum human consumption of 4990 μg Se/day was estimated, while Keshan disease, the disease caused by Se deficiency, is regularly observed in the Northeast of China, where dietary Se consumption is in the range of only 7-11 $\mu\text{g}/\text{day}$ Se (Huang et al., 2013).

Selenium is released into the environment from both natural and anthropogenic sources. It is naturally found in the Earth's crust in an estimated concentration of 0.05–0.5 mg/kg (Tan et al., 2016). Rocks are the primary source of Se, which comprise approximately 40% of the total Se form in the Earth's crust (Wang & Gao, 2001). A

Chapter 2

greater Se concentration (0.06 mg/kg) is usually found in shales compared to that in limestones and sandstones (Fordyce, 2007). Coals and other organic-rich deposits can be rich in Se (typically from 1 to 20 mg/kg). Very high concentrations of Se have also been reported in some phosphatic rocks (≤ 300 mg/kg) and some black shales (300 mg/kg). Selenium is also often found as a minor component of sulfide mineral deposits (Fernández-Martínez & Charlet, 2009; Fordyce, 2007; Fordyce, 2013).

Human activities contribute to the introduction, mobilization and accumulation of Se in the environment (Winkel et al., 2015). Fordyce (2013) estimated that around 88000 Se tons/year are released globally to the environment from anthropogenic activities, which accounts for 50–65% of the total Se emissions. Industrial processes, in particular coal and petroleum combustion, are thought to be the main processes releasing Se into the atmosphere. Selenium is also mainly a by-product of the extraction of various metal elements (i.e. copper, zinc, uranium and lead) and of processing plants (i.e. sulfuric acid production) (Tan et al., 2016). Other anthropogenic activities, such as glass and electronics production, utilization of rock phosphates as fertilizer and application of sewage sludge or Se-containing fertilizers to agricultural land for the enhancement of the Se concentration in food, also increase the Se content in the environment (Lemly, 1997).

The Se released by anthropogenic (e.g., coal combustion, mining and agricultural) and natural (e.g., biomethylation and volcanic eruptions) activities enters into nature and starts a cycle involving different Se species transformations (Fig. 2.1). Briefly, the released Se enters into water and soil in the form of Se(IV) and Se(VI) by leaching/discharging and weathering/precipitation processes, respectively. Se(IV) and Se(VI) in water or soil can be taken up by organisms (e.g., plants, animals and bacteria) and converted into organic Se. The organic Se in the organisms can be further methylated into volatile Se (e.g. Dimethyl diselenide (DMDSe) or Dimethyl selenide (DMSe)) and released into the atmosphere as H_2Se and SeO_2 . Selenium enters into the atmosphere in the form of SeO_2 and H_2Se . The subsequent solubilization of SeO_2 in the atmosphere results in Se(IV) or Se(VI) precipitating with the rain and entering a new natural cycle.

Alternatively, Se(IV) and Se(VI) in the water and soil can also be reduced by microorganisms into Se(0) or even Se(-II), which are deposited in soils and sediments.

Chapter 2

These biogeochemical processes (e.g. weathering, leaching, rock–water connections and biological activities) mainly control the transport of Se from rocks (the main Se source) to other compartments in the environment, unevenly distributing it over the Earth (Fernández-Martínez & Charlet, 2009). This leads to widely varying Se concentrations in different geo-ecosystems, forming seleniferous or Se-deficient regions (Fernández-Martínez & Charlet, 2009). This uneven distribution is likely to affect the health of both humans and animals throughout the food chain (Tan et al., 2002).




Figure 2.1. Schematic overview of Se transformations in natural systems (Winkel et al., 2015).

2.2 Se toxicity and deficiency

The range between beneficial and harmful Se concentrations is relatively narrow for animals and humans (Li et al., 2015a). Thus, both toxicity and deficiency have been reported over the world (Li et al., 2015a). In humans, chronic Se toxicity is observed

Chapter 2

above intake levels of 400 µg/day and Se deficiency occurs when the dietary intake of Se is below 40 µg/day (Winkel et al., 2012). More specifically, the tolerable upper intake levels are 90 µg/day for children of 1–3 years, 150 µg/day for children of 4–8 years, 280 µg/day for children of 9–13 years, and 400 µg/day for children >14 years and adults (Ngigi, 2019; Sciences, 2000). Besides, for livestock, the toxic Se concentration in animal feed is 2–5 mg/kg dry forage. On the other hand, the minimum requirement is defined as 0.05–0.10 mg/kg (Gupta & Gupta, 2017). The National Research Council (NRC, 2005) has published the following maximum tolerable levels (MTL) for animals: 5 mg Se/kg feed dry matter (DM) for cattle and sheep, 4 mg Se/kg feed DM for pigs, and 3 mg Se/kg feed DM for poultry. The MTL for horses and fish were derived from interspecies extrapolation and amount to 5 and 2 mg Se/kg DM feed, respectively (NRC, 2005).

Se intoxication events for animals and humans, such as selenosis in America, Canada, China, and Mexico, have occurred occasionally because Se has entered the food chain (Li et al., 2015a). These events were caused by excessive Se concentrations in soil and water. For instance, the discovered Se toxicity for humans and animals in the Enshi District, Hubei Province and in Ziyang County, Shaanxi Province in China was related to the exceedingly high Se concentrations in the local food and environment (Fordyce et al., 2000). For humans, Se toxicity (selenosis) could result in garlic breath, hair and nail loss, nervous system disorders, poor dental health and paralysis (Rayman, 2012). For animals, Se can cause alkali disease and blind staggers in livestock, and hooves loss in hooved animals (Fordyce, 2007; Tan et al., 2002). The alkali disease is characterized by dullness, lack of vitality, emaciation, rough coat, sloughing of the hooves, erosion of the joints and bones, anaemia, lameness, liver cirrhosis, and reduced reproductive performance (Reilly, 2006). Blind staggers result in impaired vision and blindness, anorexia, weakened legs, paralyzed tongue, labored respiration, abdominal pain, emaciation, and death (Fordyce, 2007). Hair loss and other abnormalities of farm animals have been observed in areas of Columbia as a result of Se toxicity (Johnson et al., 2009).

On the contrary, Se deficiency is also observed frequently worldwide and is even more widespread than Se toxicity. It is estimated that 0.5–1 billion people are directly affected by Se deficiency on a global scale, due to low Se dietary intake (Haug et al., 2007; Stonehouse et al., 2020). It has been demonstrated that Se deficiency can cause

Chapter 2

the Keshan disease and Kashin-Beck disease (endemic disease) with exceedingly low Se supplies in the food system, i.e. weakening of the heart and also atrophy and necrosis of cartilage tissue in the joints, which has been observed in the middle of China (Stone, 2009), Saudi Arabia, Czech Republic, Burundi, New Guinea, Nepal, Croatia, and Egypt (Wu et al., 2015). Low Se status has also been associated with a significantly increased risk of cancer incidence and mortality, cardiovascular risk, poor immune function, male infertility and lower reproduction (Fordyce, 2007; Haug et al., 2007). In addition, Se deficiency may also be a factor in some other diseases. For instance, studies have found that the prevalences of iodine deficiency diseases were greater among populations with lower Se status than among those with higher Se status in Africa (Combs, 2001). This should probably be attributed to the fact that Se is essential for the metabolic production of thyroid hormone.

Se deficiency is known to adversely affect livestock health around the globe, which has been identified since the 1950s in several countries including South and North America, Africa, Australia, UK and New Zealand (Reilly, 2006). Selenium deficiency causes reproductive and immune response impairment of animals, growth depression (ill-thrift), and white-muscle disease, a myopathy of heart and skeletal muscle principally affecting cattle, sheep, poultry and horse (Rayman, 2000). Wolf et al. (1963) estimated that around 10–15 million sheep or 20 to 30% of the total stock were at risk of developing white muscle disease in New Zealand via an extensive international survey conducted in the 1960s.

Generally, Se deficiency in humans and animals is attributed to a low Se daily dietary intake, with this dietary intake varying considerably between countries/regions. As aforementioned, Se deficiency has mainly been identified in parts of the world which have notably a low content of Se in soil and water, as Se enters the food chain from the environment through crops and plants uptake (mainly local water or soil) (Haug et al., 2007). Therefore, the Se concentration in foods is determined by geological and geographical factors. Globally, the range of total Se concentration in soils is from 0.01 to 2.0 mg/kg (with a mean of 0.4 mg/kg) (He et al., 2010; Rayman, 2008). Some parts of the world (e.g. Denmark, Finland, New Zealand, eastern and central Siberia and a long belt extending from Northeast to Southwest China including parts of Heilongjiang, Jilin, Liaoning, Hebei, Shanxi, Shaanxi, Sichuan and Zhejiang Provinces and Inner

Chapter 2

Mongolia) have relatively low Se contents in their soils and, therefore, resulting in low amounts of Se in their food chains (Combs, 2001).

Table 2.1 summarizes the recommended daily Se intake and the status of daily Se intake in some countries. The table shows that the recommended daily Se intake in some countries is not achieved yet, such as in some European countries (including Belgium) and parts of China. This demonstrates that the food systems of these countries do not provide sufficient Se for consumption. It may, thus, be assumed that many individuals have a potential risk of Se deficiency, which can increase their risks to various diseases, including those of the heart and lungs, as well as cancer, and make them more vulnerable to infectious diseases due to poor functioning of their immune system. There is a clear need to enhance Se in food systems of these countries to remediate Se deficiency.

Table 2.1. Recommended daily Se intake and estimated Se intake statues of adults in some countries ($\mu\text{g}/\text{person}\cdot\text{d}$)

Countries	Males	Females	Se intake
Australia	85	70	57-87
Belgium	70	70	28*-61 7-11* Keshan disease area (e.g. wide belt from Northeast China to Southwest China)
China	60	60	40-120 Moderate Se area (e.g. Guangzhou) 750-4990 Selenosis area (e.g. Hubei and Shaanxi provinces)
Netherlands	50- 150	50-150	-
Germany			35* (Germany)
Austria	30-70	30-70	48 (Austria)
Switzerland			70 (Switzerland)
France	60	50	29*-43
Italy	55	55	35*-42

Chapter 2

Table 2.1 Continued

Ireland	55	55	44-59
Japan	55-60	45	104-199
Nordic countries	60	50	38*-47 (Denmark) 38* (Sweden) 25* (Finland before 1984) 67-110 (Finland after 1984, i.e. initiation of Se biofortification)
USA and Canada	55	55	60-220 (USA) 98-224 (Canada)
United Kingdom	75	60	29-39*
Scientific Committee	55	55	-
Food			
FAO/WHO	40	40	-

Table adapted from: Scientific Committee on Food (2003); Thomson (2004); Rayman (2004); EFSA (2014) and Combs (2001).

* indicates that this level does not meet the recommended requirement according to WHO (2001)

2.3 Approaches for addressing Se deficiency - biofortification

Addressing micronutrient deficiencies to reduce health-related issues can be achieved through various types of interventions, such as through food supplements, dietary diversification, biofortification, or increase of the digestibility of trace elements in products (Lavu et al., 2012; Li et al., 2020). For instance, sodium selenite has been supplemented to feeds in some areas with selenium deficiency in livestock in order to achieve optimal Se intake (EFSA, 2016).

Biofortification is one of the most promising, widespread and accepted strategies, aimed at improving the lacking of a mineral (e.g. Se) content of the diet through its enrichment in food/feed crops, in particular the edible parts of plants, through soil or foliar application of mineral fertilizers (Sánchez et al., 2017). The agronomic approach of applying a fertilizer on the soil or leaves of the crop can improve the nutritional quality

Chapter 2

of the crop without genetic modifications (Storksdieck and Hurrell, 2009). It has been developed as a food-based method to help decreasing widespread deficiencies of minerals (e.g. Se). Although Se is not an essential trace element for plants, it presents chemical similarity to S, and both elements have the same carrier membranes and biochemical pathways of assimilation in plant uptake (Prado et al., 2017). Biofortification through application of Se fertilizers can therefore ensure its sufficient concentration in the edible parts of plants (Sarwar et al., 2020). Se biofortification of food crops is already successfully practiced in some countries (Se-deficient regions) to increase the Se concentration in staple grains and subsequent dietary Se intake, such as in Finland, by adding inorganic Se fertilizer to soils (Bañuelos et al., 2016). For instance, in Finland, a 3-fold increase of mean Se intake within 2 years was observed after Se biofortification of crops through the application of selenate fertilizer, and the concomitant human serum Se concentration increased by 70% (Aro et al., 1995).

Since low concentrations of plant Se can decrease the dietary intake of Se, it is vital to increase Se uptake by plants and to produce plants with higher Se concentrations and bioavailability in their edible tissues (Bañuelos et al., 2017). This is the key issue for effectively developing a biofortification strategy. The Se biofortification efficiency depends on a number of factors associated with the Se concentration in plants during biofortification, such as plant species, Se species and source (chemical Se fertilizer, natural source of Se or organic Se), soil pH and redox conditions, soil texture and organic matter, and the presence of competitive ions (Fordyce, 2007).

Plant species: Table 2.3 summarizes Se concentrations in crops after Se fortification. Plants have been classified as hyperaccumulators (>1000 mg/kg, such as *Stanleya*), secondary accumulators (100–1000 mg/kg, such as *Brassica* species: broccoli), and non-accumulators depending upon Se accumulation inside their cells (Gupta & Gupta, 2017). Vegetables (e.g. *brassica* species: pak choi and cabbage) normally accumulate more Se than legumes (beans), followed by cereals (wheat and rice). The Se concentration accumulated in fruits is generally low, whereas high concentrations (ranging from 0.03–512 mg/kg) have been reported in Brazil nuts as a result of natural biofortification (Prado et al., 2017).

Chapter 2

Table 2.3. Se concentration of some selected Se-enriched plants (crops, vegetable, and fruits) after Se fortification (Gupta & Gupta, 2017)

	Se-enriched parts	Accumulated Se (mg/kg)	Se dose for biofortification
Broccoli	Sprouts	467	60 mg/L
Kale	Sprouts	155	60 mg/L
Pak choi	Shoots	20–4000	2.5–40 mg/kg
Lettuce	Shoots	43	< 2.8 mg/L
Soybean	Seeds	75	130 mg/kg
Rice	Grains	1.3–3.3	2.850 mg/kg
Pear	Fruit	0.199	1 mg/L

Se application methods: Different application methods of Se-based fertilizer affect Se accumulation and transformation in plants. Foliar application is generally more efficient in enhancing the Se concentration in plants in comparison with soil application. Studies showed that foliar Se applications are on average 8 times more efficient than soil applied fertilizers (Ros et al., 2016). Besides, application of Se fertilizers at different plant growth stages can also result in a different biofortification efficiency. Wang et al. (2020) demonstrated that foliar application of selenate or selenite at the pre-filling stage was superior in improving the Se concentration of wheat grains than that at the pre-flowering stage. Zhang et al. (2019) found that the foliar application of selenite during the potato tuber bulking stage resulted in the greatest Se accumulation in tubers, compared to the application during the tuber initiation and maturation stages.

Se species and source: The uptake rates and mechanisms of selenite, selenate and organic Se are different. Some studies showed that selenite is absorbed and taken up in a faster passive way and readily reduced to organic compounds in plants, while selenate is taken up in an active way and easily distributed from roots to shoots (Arvy, 1993; Gupta & Gupta, 2017). Selenate reduction occurs via substitution for sulfate in the ATP sulfurylase reductase system, which is an ATP-consuming process and rate-limiting step, resulting in lower selenate accumulation in plants compared to selenite (Van Hoewyk, 2013). However, Ros et al. (2016) showed that biofortification using selenate-based fertilizers has a high potential to increase Se uptake by crops and subsequently Se intake by animals and humans. This is attributed to the fact that

Chapter 2

selenate is not easily adsorbed into the soil matrix in comparison with selenite, resulting in higher bioavailable Se concentration in the soil, while selenite is readily adsorbed in the soil environment.

Soil pH and redox condition: soil pH and redox conditions have an important effect on Se availability since a combination of these factors determines the Se species present in a given soil environment. For instance, selenate is the predominant Se species in near-neutral pH environments under aerobic conditions, whereas selenite predominates at lower pH and redox potential. Selenate is much more mobile, and thus plant-available, in soils than selenite which is tightly bound to positively charged binding sites in soil (Eich-Greathorex et al., 2007). Besides, soil pH negatively correlates with the amount of Se adsorbed to soil (Li et al., 2015b). Most studies have demonstrated that relatively high pH values in soil solutions lead to a higher Se accumulation by plants in comparison with low pH soil (Li et al., 2016; Li et al., 2017). This is attributed to the fact that soil with low pH contains a high amount of H^+ in the soil matrix, which does not compete with selenite/selenate occurring as oxyanion (e.g., SeO_4^{2-} and SeO_3^{2-}) for positively charged binding sites, thus leading to a relatively high bioavailability of Se in the soil solution.

Soil organic matter: Organic matter (OM) influences Se availability in different ways. On the one hand, OM has a significant capacity to remove Se from the soil solution, and immobilize Se by both biotic and abiotic mechanisms, thus reducing Se bioavailability. On the other hand, OM can improve the soil structure and stimulate oxidizing conditions, thus enhancing Se bioavailability (Li et al., 2017). The release of OM-immobilized Se through mineralization will increase the bioavailable Se concentration in soil.

Competitive ions: The Se accumulation in plants can also be influenced by the presence of other ions, especially phosphate (PO_4^{3-}) and sulfate (SO_4^{2-}). Interactions between Se and other ions may occur in the soil or in the plant (Bingham, 1989). Li et al. (2008) studied the Se uptake in wheat under P and S-starved conditions and demonstrated that selenite uptake is an active process mediated partly by P transporters. Likewise, the Se uptake can be negatively influenced by the addition of sulfur (S) due to the chemical similarity between these two elements. Studies have demonstrated that selenate is taken up by sulfate transporters, thus the competition of

Chapter 2

the same transporters could inhibit Se uptake by plants when S is applied (Li et al., 2008). For instance, a decrease in Se concentration in the shoots and roots of corn (*Zea mays*) was observed when the S concentration in solution increased (Huang et al., 2008). Supplementation of S in the calcareous alluvial and yellow-brown soil reduces the Se contents in soybean (*Glycine max* L.) seeds (Deng et al., 2021).

2.4 Approaches for addressing Zn deficiency – biofortification

Similarly, Zn is also an important micronutrient because it plays an important role in crop production and human nutrition (Broadley et al., 2007; Sánchez et al., 2017). Approximately 10 % of human proteins require Zn to maintain their catalytic activity. Zinc is involved in the biosynthesis of proteins and scavenging of reactive oxygen species (Li et al., 2020b). It is deficient in 30 % of the soils used for agriculture in the world (Poblaciones & Rengel, 2017; Sánchez et al., 2017), and the WHO reports that about 33 % of the population is affected by Zn deficiency, in particular for developing countries. Zn deficiency affects organ functions such as epidermal, gastrointestinal, central nervous, immune, skeletal, and reproductive systems (Roohani et al., 2013). Zinc deficiency will also impair children's physical growth and development. It can result in a syndrome of anemia and increase the risk of pathogenic infections and diseases (Gibson, 2006). Children, pregnant and lactating women require more Zn, which thus have a higher risk of Zn deficiency (Roohani et al., 2013). It is estimated that more than half of the pregnant women and children in developing countries are suffering from Zn deficiencies (Maqbool et al., 2019).

Zn deficiency is also due to the inadequate intake of total dietary Zn in most situations. Besides, a high amount of phytic acid in diets is another important factor for causing Zn deficiency in humans and animals, as phytate has inhibitory effects on Zn absorption by animals or humans, leading to low Zn bioavailability (Lönnerdal, 2002). Increasing Zn levels in crops and/or reducing the phytate content in the diet may lead to more Zn in humans. Zinc biofortification of crops as rice, maize and wheat through foliar or soil application has been approved as an effective and direct strategy to increase the Zn concentration in crops and the diet (Sánchez et al., 2017). Some countries (e.g. China, Mexico, Indonesia and South Africa) have implemented the biofortification of maize or wheat flour with Zn (Gibson, 2006). Zinc biofortification through use of mineral fertilizers has been extensively studied. Many comprehensive

Chapter 2

overviews on the current status, challenges and solutions of Zn biofortification, dealing with the use of mineral of Zn fertilizers for combating Zn deficiency have been written (Maqbool et al., 2019; Palmgren et al., 2008; Zaman et al., 2018).

However, Se and Zn deficiency in humans and animals often co-occur (Darago et al., 2016; Ruz et al., 1999). For instance, in Kenya, Zn deficiency stands at 83.3%, 68.3% and 82.3% for children under 5 years of age, pregnant women, and women of reproductive ages respectively (Health, 2018), and 87% of children and 97% of women were found to be at risk of dietary Se deficiency based on the estimated average requirements (Ngigi, 2019). Joy et al. (2014) estimated the dietary mineral supplies in 46 African countries and found that the mean estimated risks of Zn and Se are 40% and 28%. Selenium and Zn deficiency risks are highest in the Eastern region of Africa. They also demonstrated that many countries have high multiple dietary micronutrient deficiencies, for instance, the estimated deficiency risks for I, Se and Zn in Ethiopia are 64, 38 and 81%, and in Malawi are 27%, 64% and 33% (Joy et al., 2014).

Both the Se and Zn content in the human and animal diet is, therefore, a topic of interest to public health systems around the world. Multi-mineral agronomic biofortification of crops is thus being explored as a simple and effective way to simultaneously alleviate several micronutrient deficiencies (Mao et al., 2014; Poblaciones & Rengel, 2017), e.g. White and Broadley (2009) reviewed the biofortification of crops with essential multi-mineral elements often lacking in human diets (e.g., Zn, Se, Cu, I and Fe) (White & Broadley, 2009; White & Broadley, 2005). However, most previous studies have only investigated single Se or Zn biofortification (or feed supplementation). Combined Se and Zn agronomic biofortification for simultaneously improving concentrations of these two micronutrients in crops was not yet studied thoroughly, particularly not when dealing with slow-release organic biofertilizers. Hence, this thesis explored the case of simultaneous Se and Zn biofortification via the application of Se and Zn-enriched organic fertilizers.

2.5 Se/Zn-enriched organic materials as biofertilizers for biofortification

Application of mineral Se/Zn fertilizers originating from primary mining would cause a waste of Se/Zn due to the low utilization of plants and fast leaching of the freshly added minerals. Biomaterials (e.g. plant residues, sludge, and manures) that come from

Chapter 2

seleniferous and zinciferous areas potentially contain high levels of Se and/or Zn. These micronutrient-enriched materials may serve as potential micronutrient sources and can thus be re-utilized for Se or Zn biofortification of agricultural crops. If Se/Zn-enriched organic biomaterials are used to amend agricultural soils, the decomposition of organic biomaterials will gradually lead to the micronutrients released into the soil solutions, which would be bioavailable for crops uptake (Bañuelos et al., 2015). In this context, biofortification with these micronutrient-enriched biomaterials can thus be achieved, which may be particularly beneficial for crops grown on micronutrient deficient soils (Bañuelos et al., 2016; Li et al., 2017). Some studies have investigated the possibility of using Se-enriched biomaterials as feedstock to improve the Se concentration in crops for biofortification purposes. For instance, the accumulation of Se in canola, grown on soil amended with 1.5 mg/kg seleniferous *Astragalus praelongus* E. and *Medicago saliva* L. tissues, was increased as the amount of application of these materials increased (Ajwa et al., 1998). Moreover, Se-enriched wheat and raya plants straw were used to biofortify sorghum, maize and berseem (Dhillon et al., 2007), and results showed that the Se concentrations in the plant were consistent with the trend of soluble Se in soil.

The supplementation with Se-enriched organic materials in soils as biofertilizer may not only improve the Se concentration in the plants, but also results in value-added plant-based products, as plants can transform the Se taken up during growth into valuable organic Se species (e.g. SeMet, SeCys and MeSeCys), which have important assets in the nutrition of animals and humans. Bañuelos et al. (2015) reported that the Se concentration in the edible parts of broccoli and carrots was increased and that MeSeCys was the main accumulating Se species when the shoots of Se-enriched *Stanleya pinnata* were added to the soil as biofertilizer.

One of the main advantages of micronutrient-enriched organic materials is that they provide a long-lasting micronutrient source, slowly releasing the micronutrient along with the decomposition of the organic materials in the soil (Ajwa et al., 1998). However, the disadvantage is that the application of these materials can introduce additional organic matter into the soil, which can lead to the immobilization of other elements/nutrients in the soil, eventually decreasing the bioavailability for plant uptake (Stavridou et al., 2011).

Chapter 2

It should be noted that Se/Zn biofortification via the application of Se/Zn-enriched organic materials may not be feasible in all Se/Zn-deficient areas. For instance, the Se-deficient region in Northeastern China, characterized by a high content of OM, is not suitable for supplementation with micronutrient-enriched organic materials, as the presence of too much organic matter in the soil will increase the retention of the released Se and Zn, reducing the bioavailability of Se/Zn in the soil. In contrast, some regional soil with strong leaching potential (i.e. high precipitation (rainfall) and humid climates) and low Se/Zn content can benefit from the addition of micronutrient-enriched organic materials since the added organic matter can act as a micronutrient reservoir to avoid the leaching of nutrients and their mobilization to the deeper soil layers (Wang & Gao, 2001).

Although almost all Se-enriched biomaterials studied for use as biofertilizers to improve Se levels in crops are produced on seleniferous soils, wastewater also contains Se, for example flue gas desulfurization process water (1.0-10000 µg Se/L) and agricultural drainage (140-1400 µg Se/L) (Tan et al., 2016), and may be used to enrich crops with Se. Therefore, it can be questioned whether the Se contained in wastewaters can act as a nutrient source for the production of Se/Zn-enriched biomaterials while cleaning the water. For some wastewaters barely containing Se/Zn but sufficient other nutrients, it would be possible to use them as feedstock for the production Se/Zn-enriched biomaterials, wherein Se/Zn enrichment is achieved via external spiking of Se/Zn. However, no research has been done on this before.

2.6 Local “green” production of micronutrient-enriched bioproducts by micronutrient removal from wastewater

In order to achieve optimal Se and Zn levels in the human diet, chemicals containing Se and Zn are commonly added to crops in deficient regions. Similarly, the production of micronutrient-enriched food/feed supplements is being explored as another solution for micronutrient deficiencies. However, for the biofortification process, plants usually can take up only a small amount of those applied elements (Tan et al., 2002). The residual trace elements are leached with rainwater or fixed and accumulated in the soil, thus posing a potential threat to the environment (Broadley et al., 2006; Wang et al., 2018). Micronutrient slow-release fertilizers, such as organic fertilizers (Bañuelos and Hanson, 2010) and chemical nano-Se (Wang et al., 2017), are currently being

Chapter 2

developed to overcome these limitations and risks. Moreover, chemical production processes currently used to produce inorganic fertilizers and food/feed supplements are usually not environmentally and economically sustainable. Given this, it may be beneficial to produce micronutrient-enriched fertilizers and food/feed supplements locally from micronutrient-containing wastewater using environmental-friendly techniques.

Bioremediation is an environmentally friendly method to recover micronutrients from wastewater. Of all treatment technologies, bioremediation approaches may have the lowest construction and operation costs for contaminant removal. Moreover, the macrophytes *Lemna* and *Azolla* have fast growth rates, high tolerance/accumulation to extreme conditions, and can easily be harvested (Miranda et al., 2016; Sasmaz et al., 2015). Ohlbaum et al. (2018) found that the duckweed *Lemna minor* can efficiently remove 76% of Se from seleniferous soil leachates (with 74 µg Se/L). *Azolla filiculoides* grown in synthetic wastewater reduced the Se content up to 40 % after 5 days of treatment (Miranda et al., 2016). *Lemna* and *Azolla* have also a high potential to remove Zn (Sasmaz et al, 2015). Moreover, these two aquatic plants are rich in proteins, so they may also be considered as an alternative (micronutrient-enriched) protein source replacing animal proteins in food and feed systems, contributing to the sustainability of our food and feed production systems (Terry et al., 2000). Based on the above information, we hypothesize that Se/Zn-enriched aquatic plants with high protein content such as *Lemna* and *Azolla* can be produced from wastewater as value-added feed supplement or biofertilizer. Research on the ability of these plants to simultaneously remove Se and Zn, including possibly interaction effects, is needed to confirm this hypothesis..

Similarly, microalgae have fast growth rates and a high protein content (50–60%, depending on nutrients availability). Many studies have proposed that microalgae could be applied in wastewater treatment systems to efficiently clean water (Liu et al., 2016). They are low-cost technologies and can be successfully and easily implemented in locations where weather conditions are favorable for microalgae growth, e.g. high solar radiation and temperature (Arashiro et al., 2019). Besides, the installation and maintenance of algae ponds are also cheaper compared to conventional wastewater treatment systems, e.g. activated sludge systems (Arashiro et al., 2019). Most importantly, microalgae could upgrade low-value products from wastewater and

Chapter 2

synthesize them into high-value compounds, such as protein, lipid and carbohydrate in their cells. Batch tests have already been done to study the Se removal by microalgae. For instance, Liu et al. (2016) developed a batch algae treatment system for Se removal and evidenced that the Se absorption efficiency by *Chlorella vulgaris* was 88%.

Nowadays, many bioproducts from microalgae biomass grown in wastewater have been explored. For instance, high-value phycobiliproteins have been obtained from microalgae cultivated in food-industry wastewater (Arashiro et al., 2020a), municipal wastewater and anaerobic digestion concentrate (Arashiro et al., 2020b); high-quality fatty acids, such as oleic acid for biodiesel production, have been extracted from microalgae biomass grown in sewage (Khan et al., 2019), piggery wastewater (Chen et al., 2020) and starch processing wastewater (Tan et al., 2019); and plant growth-promoting substances (e.g., phytohormones and polysaccharides) have been recovered together with high N and P contents from microalgae biomass cultivated in wastewater, enabling its use as biofertilizer or biostimulant (Supraja et al., 2020). Because of these results, we hypothesize that the production of value-added (Se-enriched) microalgae biomass during wastewater treatment processes is feasible. Currently, there are no studies yet investigating the Se removal efficiency by microalgae in continuous or pilot-scale systems, meanwhile assessing the possibility to use the produced Se-enriched microalgae as Se-rich biofertilizer or feed supplement.

Bioprecipitation or bioreduction, i.e. Se reduction by microorganisms, can also be used to remove Se from water, meanwhile producing more sustainable Se biofertilizers (Staicu et al., 2015a; Staicu et al., 2015b). In wastewater, Se is mainly present as Se oxyanions, namely selenite and selenate, which are soluble, bioavailable and toxic for the environment. However, elemental Se(0) is solid and less toxic in the water phase. Reduction of selenite and selenate in the water phase into solid-phase elemental Se(0) could thus be another sustainable and efficient method to remove Se from wastewater. Staicu et al. (2015a) described the reduction of selenite and selenate by anaerobic granular sludge, and Se nanoparticles (SeNPs) were obtained by the anaerobic reduction process. As these SeNPs were stable in the solid phase, reusing the Se-containing sludge dominated by SeNP as a slow-release micronutrient biofertilizer is therefore theoretically proposed. However, from a practical point of view, it is unknown whether the Se-enriched sludge can slowly release Se, whether the released Se is

Chapter 2

bioavailable for plant growth, and if the Se-enriched sludge produced from wastewater could negatively affect plant growth.

To the best of our knowledge, this thesis is the first study to valorize the application of Se-enriched biomaterials produced from wastewater as Se biofertilizer and feed supplement. Such approach would be beneficial to save mineral Se resources via waste reuse, while avoiding the introduction of new Se contamination into the environment through the use of inorganic Se as fertilizer, the current practice in biofortification. However, potential risks (e.g. heavy metals, pathogens and organic pollutants) arising from the use of products from wastewater should always also be considered.

Chapter 3 Production of selenium- and zinc-enriched *Lemna* and *Azolla* as potential micronutrient-enriched bioproducts

This chapter was redrafted after:

Li, J., Loi, G., Otero-Gonzalez, L., Du Laing, G., Ferrer, I., Lens, P.N.L. 2020. Selenate and selenite uptake, accumulation and toxicity in *Lemna minuta*. *Water Science and Technology*. 81(9), 1852-1862.

Li, J., Lens, P.N.L., Otero-Gonzalez, L., Du Laing, G. 2020. Production of selenium- and zinc-enriched *Lemna* and *Azolla* as potential micronutrient-enriched bioproducts. *Water Research*. 172, 115522.

Chapter 3

Abstract

Selenium (Se) and zinc (Zn) are essential micronutrients that are often lacking in the diet of humans and animals, leading to deficiency diseases. *Lemna* and *Azolla* are two aquatic plants with a substantial protein content, which offer the possibility of utilizing them to remove Se and Zn from wastewater while producing micronutrient-enriched dietary proteins and fertilizers. In this study, we explored interaction effects occurring between Se and Zn when these micronutrients are taken up by *Azolla* and *Lemna*. The two aquatic plants were grown on hydroponic cultures containing 0–5.0 mg/L of Se (Se(IV) or Se(VI)) and Zn. The Se and Zn content of the plants, growth indicators, bioconcentration factor (BCF) and Se/Zn removal efficiency from the water phase were evaluated. The results demonstrated that Se(IV) is more toxic than Se(VI) for both plant species, as evidenced by the remarkable decrease of biomass content and root length when exposed to Se(IV). Both aquatic plants took up around 10 times more Se(IV) than Se(VI) from the medium. Moreover, the Se accumulation and removal efficiency increased by 66–99% for Se(IV) and by 34–59% for Se(VI) in *Lemna* when increasing Zn dosage from 0 to 5.0 mg/L in the medium, whereas it declined by 13–26% for Se(IV) and 21–35% for Se(VI) in *Azolla*, suggesting a synergetic effect in *Lemna*, but an antagonistic effect in *Azolla*. The maximum BCF of Se in *Lemna* and *Azolla* were 507 and 667, respectively. The protein content in freeze-dried *Lemna* and *Azolla* was approximately 17%. The high tolerance and accumulation of Se and Zn in *Lemna* and *Azolla*, combined with their rapid growth, high protein content and transformation of inorganic to organic Se species upon Se(IV) exposure make *Lemna* and *Azolla* potential candidates for the production of Se(IV)- and Zn-enriched biomass that can be used as crop fertilizers or protein-rich feed supplements or ingredients. Accordingly, by growing the *Azolla* and *Lemna* on wastewater, a high-value product can be produced from wastewater while recovering resources.

Keywords: Aquatic plants; dietary protein; micronutrient fertilizer; nutrient accumulation; Se; Zn

Chapter 3

3.1 Introduction

Se and Zn are essential for humans and animals. Certain human proteins require Se or Zn to maintain their catalytic activity. Se and Zn are involved in the biosynthesis of proteins and in scavenging of reactive oxygen species (Sánchez et al., 2017). For plants, Zn is also an essential element that can enhance plant growth, whereas Se is considered as a beneficial element that can enhance resistance to stress (Feng et al., 2013; Subramanyam et al., 2019). Despite the importance of these trace nutrients, Se and Zn intakes are still low in a wide range of countries (Broadley et al., 2006; Thomson, 2004), resulting in Se and Zn deficiencies. In order to achieve optimal Se and Zn levels in the human diet, Se and Zn inorganic fertilizers are commonly added to crops in deficient regions for enrichment of crops with Se and Zn. However, plants cannot take up all of Se applied through conventional inorganic fertilizers, because of fast leaching and high mobilization of the applied inorganic Se and Zn (Broadley et al., 2006; Wang et al., 2018). Micronutrient slow-release fertilizers, such as organic fertilizers (Bañuelos et al., 2016; Bañuelos & Hanson, 2010), are therefore currently being explored.

On the contrary, Se and Zn excess in wastewaters also frequently occurs, as Se and Zn are widely applied in or released from industrial and agricultural activities (Lim & Goh, 2005). Selenium compounds exist in nature in five redox states: Se(-II), Se(0), Se(II), Se(IV) and Se(VI) (Terry et al., 2000). Selenium exists in wastewater mainly as Se(IV) and Se(VI) oxyanions due to their high solubility, with Se(VI) being more abundant than Se(IV). Wastewaters loaded with both Se and Zn can serve as potential sources from which the nutrients may be valorized to produce slow-release organic fertilizers. Accordingly, it could be beneficial to produce micronutrient-enriched fertilizers locally from micronutrient-containing waters using environmentally friendly techniques.

Phytoextraction is an environmentally friendly method to recover micronutrients from wastewater. The floating aquatic plants *Lemna* and *Azolla* have fast growth rates, high tolerance to extreme conditions, and can be easily harvested, making them as potential plant species for natural wastewater/water treatment systems (Miranda et al., 2016; Sasmaz et al., 2015). Most previous studies have investigated the Se or Zn removal by aquatic plants individually, while little studies consider the case of Se and Zn co-occurrence in the environment. Hence, this chapter explored the case of simultaneous

Chapter 3

Se and Zn removal from water by these floating aquatic plants and investigated the simultaneous effects of Se and Zn on these plants. In addition, earlier studies only focused on the removal of contaminants (Tan et al., 2016). In this study, we therefore will not only consider the removal of Se and Zn by *Lemna* and *Azolla*, but also propose the potential use of the Se- and Zn-enriched *Lemna* and *Azolla* as micronutrient-enriched feed supplements and biofertilizers. Especially, these two aquatic plants are rich in proteins, so they may also be considered as alternative (Se- and Zn-enriched) protein sources replacing other plant and animal proteins in feed systems. This may be particularly useful for Se, as Se is preferably accumulated as an organic form, i.e., as selenoaminoacids (Eiche et al., 2015). Such alternative proteins would contribute to the sustainability of our food and feed production systems, as they can be produced on wastewater as nutrient source during secondary or tertiary treatment.

Accordingly, in this study two aquatic plants, *Lemna* and *Azolla*, were planted in media with different dosages of Se and Zn to investigate: 1) the effect of two Se forms (selenite and selenate), supplied together with Zn, on the growth of the two plants; 2) the potential of the two plants to remove Se and Zn together, towards the potential use of the Se- and Zn-enriched *Lemna* and *Azolla* as micronutrient-enriched fertilizers and/or feed supplements; and 3) the potential interactions of Se and Zn on the uptake of these elements by those two plant species.

3.2 Materials and methods

3.2.1 Experimental materials

The aquatic plants *Lemna minuta* and *Azolla cristata* were collected from a natural freshwater canal in Delft (The Netherlands) and cultivated in modified Hoagland solution at pH 6 to acclimatize for seven days in a greenhouse (Hassan & Mostafa, 2016; Ohlbaum et al., 2018). The modified Hoagland solution contained: 472 mg/L $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$, 202 mg/L KNO_3 , 197 mg/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 9 mg/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.62 mg/L $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$, 1.14 mg/L H_3BO_3 , 32 $\mu\text{g/L}$ $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 12.8 $\mu\text{g/L}$ $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$, 1.79 $\mu\text{g/L}$ $\text{NaWO}_4 \cdot 2\text{H}_2\text{O}$, and 4 $\mu\text{g/L}$ $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$. Afterwards, 1 g (wet weight) of each plant was transplanted into 150 mL of modified Hoagland solution with varied concentrations of sodium selenite (Na_2SeO_3), sodium selenate (Na_2SeO_4) and zinc chloride (ZnCl_2).

Chapter 3

3.2.2 Experimental design

The concentrations of Se and Zn ranged from 0 to 5.0 mg/L, including 0, 0.5, 2.5 and 5.0 mg/L, and also additionally 1.0 mg/L for Zn. The selected range of Se and Zn concentrations is chosen based on their concentrations in wastewater streams and literature review on the ability of aquatic plants to tolerate them. Medium without Se and Zn served as control. All experiments were performed in triplicate, with a total of 210 pots for the 70 treatments (Table 3.1). The temperature in the greenhouse varied between 25 and 30 °C and light was provided with a minimum light intensity of 100 $\mu\text{mol}/\text{m}^2/\text{s}$ photons (16 h of light and 8 h of darkness). The whole plants were harvested after seven days of cultivation, washed with deionized (DI) water and analyzed for dry weight, root length, total Se and Zn content, and protein content.

3.2.3 Analytical methods

3.2.3.1 Determination of Se and Zn concentration

All harvested whole plants from the same pot were oven-dried at 60 °C until constant weight, homogenized and then digested using a microwave oven (CEM Mars 5, Matthews, NC, USA). Dry samples were weighed into a digestion vessel followed by the addition of 10 mL concentrated HNO_3 . The digestion temperature was raised to 165 °C in 10 min and kept for 1 min, then raised to 175 °C in 2 min and maintained for 5 min. The digested solution was diluted with DI water and analyzed for total Se and Zn. Total Se was determined using an atomic absorption spectrophotometer coupled to a graphite furnace (GF-AAS, Thermo Elemental Solaar MQZ, GF95, Thermo Fisher Scientific, Waltham, MA, USA) and total Zn with flame atomic absorption spectroscopy (F-AAS, AAnalyst 200, Perkin Elmer, Waltham, MA, USA) as described by Ohlbaum et al. (2018) and Mal et al. (2016), respectively.

Chapter 3

Table 3.1. Se(IV) or Se(VI) concentration and Zn concentration applied in each treatment with *Lemna* or *Azolla*. Each plant received 35 different treatments including 5 treatments of different Zn concentrations in the absence of Se (column 1), 15 treatments of different Se(IV) and Zn concentrations and 15 treatments of different Se(VI) and Zn concentrations (columns 2, 3 and 4).

Treatment	Content		Treatment	Content		Treatment	Content		Treatment	Content	
	(mg/L)			(mg/L)			(mg/L)			(mg/L)	
	Se	Zn		Se	Zn		Se	Zn		Se	Zn
Control	0	0	Se0.5Zn0	0.5	0	Se2.5Zn0	2.5	0	Se5.0Zn0	5.0	0
Se0Zn0.5	0	0.5	Se0.5Zn0.5	0.5	0.5	Se2.5Zn0.5	2.5	0.5	Se5.0Zn0.5	5.0	0.5
Se0Zn1.0	0	1.0	Se0.5Zn1.0	0.5	1.0	Se2.5Zn1.0	2.5	1.0	Se5.0Zn1.0	5.0	1.0
Se0Zn2.5	0	2.5	Se0.5Zn2.5	0.5	2.5	Se2.5Zn2.5	2.5	2.5	Se5.0Zn2.5	5.0	2.5
Se0Zn5.0	0	5.0	Se0.5Zn5.0	0.5	5.0	Se2.5Zn5.0	2.5	5.0	Se5.0Zn5.0	5.0	5.0

Chapter 3

3.2.3.2 Determination of protein content

The protein content was calculated by multiplying the total nitrogen (TN) concentration by 5.0. The conversion factor 5.0 was selected based on the literature (Brouwer et al., 2019; Brouwer et al., 2018; Zhao et al., 2014) and the analyzed amino acid profile of the two plants (Table 3.S1). TN was determined according to Van Ranst et al. (1999). Dry whole plant samples (0.100 g) were weighed and digested with 0.2 g Se catalyst and 10 mL concentrated sulfuric acid at 380 °C until the digestion solution turned clear. After acid digestion, the sample solution was cooled and ammonia was distilled and collected in boric acid. Then, a back titration was performed to measure the N concentration.

3.2.3.3 Determination of Se speciation

Se speciation analysis was determined according to Lavu et al. (2012 and 2013). Specifically, 0.1 g of whole plant samples and 40 mg of the enzyme protease XIV (Sigma Aldrich, St. Louis, MO, USA) were dispersed in 5 mL water in a 10-mL centrifuge tube. The mixture was shaken for 24 h at 37 °C and centrifuged for 30 min at 10000g. The supernatant was filtered through a 0.45- μ m syringe PVDF membrane filter. The filtrate was analysed for total Se and Se speciation by inductively coupled plasma-mass spectrometry (ICP-MS, PerkinElmer DRC-e, Sunnyvale, CA, USA) and ICP-MS coupled to high performance liquid chromatography (Series 200 HPLC, Perkin Elmer, Sunnyvale, CA, USA), respectively. A Hamilton PRP-X100 anion exchange column (250 mm $\times$ 4.6 mm, 5 μ m) was used as stationary phase in the HPLC instrument. The mobile phase was 10 mM citric acid with 5% (v/v) methanol, adjusted to pH 5.0. The standard solutions of the different Se species were prepared with sodium selenite (Na_2SeO_3), sodium selenate (Na_2SeO_4), Se-methionine (SeMet), Se-cystine (SeCys_2), Se-methyl-selenocysteine (SeMetSeCys), γ -glutamyl-methylselenocysteine and γ -glutamyl-selenomethionine.

3.2.4 Statistical analysis

Data were analyzed by ANOVA and Duncan's multiple comparison tests in SPSS 20.0. Graphs and tables were plotted by Excel 2016 and R-3.4.1. The following parameters were calculated:

Chapter 3

$$\text{Bioconcentration factor (BCF)} = \frac{C_{\text{plant}}}{C_{\text{medium}}} \quad (1)$$

$$\text{Plant removal efficiency (\%)} = \frac{C_{\text{plant}} \times M_{\text{plant}}}{C_{\text{medium}} \times V_{\text{medium}}} \times 100\% \quad (2)$$

Where C_{plant} (mg/kg dry weight) is the average Se or Zn concentration in the *Azolla* or *Lemna* biomass, C_{medium} (mg/L or mg/kg) is the total Se or Zn concentration in the corresponding medium, M_{plant} is the average biomass weight and V_{medium} is the volume of the medium (150 mL).

3.3 Results

3.3.1 Plant growth

The effect of Se and Zn on the growth of *Lemna* and *Azolla* was assessed based on the biomass production (i.e., dry weight, Fig. 3.1) and the root length (Fig. 3.2). A decrease of biomass production of both plant species was significantly associated with Se(IV) and Zn application (Fig. 3.1a and 3.1c) ($P < 0.01$). Increasing concentrations of Se(IV) from 0 to 5.0 mg/L reduced the dry weight of *Lemna* stepwise from 0.12 g to 0.05 g (Fig. 3.1a) ($P < 0.01$), while 0.5 mg/L of Se(IV) reduced the dry weight of *Azolla* from 0.10 g to 0.08 g ($P < 0.01$), and no further decrease was observed at higher Se(IV) concentrations (Fig. 3.1c). On the other hand, the exposure to up to 5.0 mg/L of single Se(VI) did not cause growth inhibition on *Lemna* (Fig. 3.1b) ($P = 0.24$). For *Azolla*, the effect of Se(VI) was similar to that of Se(IV) (Fig. 1d). In addition, the application of Se(VI) seemed to slightly counteract the inhibiting effect of Zn on the growth of *Lemna* (Fig. 3.1b). For example, the exposure to 1.0 mg/L of Zn in the absence of Se(VI) caused a remarkable decrease in the dry weight of *Lemna* from 0.12 g to 0.05 g ($P < 0.01$), while in the presence of 0.5–5.0 mg/L of Se(VI) and the same Zn concentration (1.0 mg/L), the dry weight of *Lemna* was approximately 0.11 g ($P = 0.92$).

Chapter 3




Figure 3.1. Dry weight of plants grown at different Se and Zn concentrations: (a) *Lemna* grown on Se(IV) and Zn, (b) *Lemna* grown on Se(VI) and Zn, (c) *Azolla* grown on Se(IV) and Zn and (d) *Azolla* grown on Se(VI) and Zn. Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the same Zn application according to Duncan's multiple comparison tests ($P < 0.05$)

Chapter 3




Figure 3.2. Root length of plants after exposure to different Se and Zn concentrations: (a) *Lemna* grown on Se(IV) and Zn; (b) *Lemna* grown on Se(VI) and Zn; (c) *Azolla* grown on Se(IV) and Zn; and (d) *Azolla* grown on Se(VI) and Zn. Lines are the fitted log-logistic curves. To allow log transformation, a small value (0.01) was added to the zero Se concentration.

The length of the roots of *Lemna* and *Azolla* decreased with increasing Se and Zn concentrations. An increment of the Se(IV) dosage from 0 to 5.0 mg/L significantly reduced the root length of *Lemna* from 1.5 to 0.5 cm ($P < 0.01$) and decreased the root length of *Azolla* from 3.7 to 1.4 cm ($P < 0.01$) (Fig. 3.2a and 3.2c). Additionally, the exposure to 5.0 mg/L of Se(VI) caused a decrease of 0.7 cm in both *Lemna* ($P < 0.01$) and *Azolla* ($P = 0.02$) roots (Fig. 3.2b and 3.2d). Similarly, an increasing Zn application from 0 to 5.0 mg/L in the absence of Se caused a noticeable decrease in the root length of *Lemna* from 1.5 to 0.7 cm ($P < 0.01$), while the growth of the root length of *Azolla* was not associated with the Zn application ($P = 0.08$). The dose-response data could be described satisfactorily by a log-logistic equation for *Lemna* (Fig. 3.2a and 3.2b). From the fitted equation, the effective concentrations of Se that caused a 50% inhibition

Chapter 3

(ED₅₀) of the root length of *Lemna* were estimated at 2.7 and 3.8 mg/L Se for the application of Se(IV) and Se(VI), respectively. These results indicated that *Lemna* was more resistant to Se(VI) than to Se(IV).

3.3.2 Se concentration in *Lemna* and *Azolla*

The Se concentration in the plants differed significantly depending on the chemical form and concentration of the Se amendment (Fig. 3.3) ($P < 0.01$). For both *Azolla* and *Lemna*, the increasing of Se concentration was significantly related to the increasing Se dosage in the medium ($P < 0.01$). Both plants have a higher ability to take up Se(IV) compared with Se(VI), which is reflected by the higher Se content in the plants cultivated on the Se(IV)-enriched medium.

The presence of Zn in the medium affected the Se concentration in *Lemna* and *Azolla* differently (Fig. 3.3). In general, the Se concentration in *Lemna* gradually increased with increasing Zn application dose, while it declined in *Azolla*. Specifically, for the plants exposed to Se(IV), the maximum Se concentration in the plants was found at 5.0 mg/L Se(IV) with 5.0 mg/L Zn in *Lemna* (1665 mg/kg) and 5.0 mg/L Se(IV) without Zn in *Azolla* (1139 mg/kg) (Fig. 3.3a and 3.3c). Similarly, for the Se(VI) application, the highest Se concentration in *Lemna* was 168 mg/kg at 5.0 mg/L Se(VI) coupled with a 5.0 mg/L Zn dose, and the maximum Se concentration in *Azolla* was 196 mg/kg at 5.0 mg/L Se(VI) without Zn (Fig. 3.3b and 3.3d).

Chapter 3




Figure 3.3 Se content in plants grown at different Se and Zn concentrations: (a) *Lemna* grown on Se(IV) and Zn, (b) *Lemna* grown on Se(VI) and Zn, (c) *Azolla* grown on Se(IV) and Zn and (d) *Azolla* grown on Se(VI) and Zn. Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the same Se application according to Duncan's multiple comparison tests ($P < 0.05$).

3.3.3 Se accumulation and removal by *Lemna* and *Azolla*

The total Se accumulation in the plants and the Se removal efficiency are presented in Tables 3.2–3.5. In both plants, the total Se accumulation significantly increased with the increase of Se dose in the medium (Tables 3.2 and 3.4), whereas a decreasing trend was generally observed in the Se removal efficiency (Tables 3.3 and 3.5) ($P < 0.01$).

When both plants were exposed to Se(VI), the Se accumulation and Se removal efficiency were much lower compared to Se(IV) exposure. Specifically, the highest Se accumulation and Se removal efficiency were 15.9 and 17.3 $\mu\text{g}/\text{pot}$, and 3.6 and 3.0% in *Lemna* and *Azolla*, respectively, when exposed to the Se(VI) growth solution. However, when *Lemna* and *Azolla* were exposed to Se(IV) medium, the maximum Se

Chapter 3

accumulation and Se removal efficiency were 89.2 and 90.5 µg/pot, and 30.2 and 38.9%, respectively.

The exposure to Zn increased the Se removal efficiency and Se accumulation in *Lemna* (Tables 3.2 and 3.3), while having an inhibitory effect on the Se removal efficiency and Se accumulation in *Azolla* (Tables 3.4 and 3.5). For instance, for *Lemna* grown on 2.5 mg/L Se(IV), increasing the Zn dose from 0 to 5.0 mg/L remarkably increased the Se accumulation from 44 to 89 µg/pot and the Se removal efficiency from 12 to 24% ($P < 0.01$). In contrast, the addition of 5.0 mg/L of Zn to the growth medium of *Azolla* containing 2.5 mg/L Se(IV) caused a considerable decline of the Se accumulation from 88 to 66 µg/pot and the Se removal efficiency from 24 to 18% ($P < 0.01$).

3.3.4 BCF_{Se}

The BCF_{Se} in *Lemna* ranged from 231 to 552 for the Se(IV) addition and from 20.0 to 55.6 for the Se(VI) treatments. In *Azolla*, the range of the BCF_{Se} values was 182 to 667 for the Se(IV) application and 23.1 to 51.9 for the Se(VI) treatments (Table 3.6).

Chapter 3

Table 3.2. Total Se accumulation in *Lemna* (µg/pot). Values are mean ± standard deviation (n=3). Different letters indicate statistically significant differences among treatments with the same Se application according to Duncan's multiple comparison test ($P < 0.05$).

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
0.0	-	-	-	-	-	-	-	-	-	-
0.5	16.5±0.3 ^c	12.2±0.8 ^d	16.1±0.2 ^c	22.6±1.3 ^a	20.7±1.0 ^b	1.7±0.2 ^c	2.2±0.2 ^b	2.0±0.2 ^{bc}	1.9±0.2 ^{bc}	2.7±0.2 ^a
2.5	44.5±1.0 ^e	50.0±1.3 ^d	56.3±0.2 ^c	65.0±3.0 ^b	88.8±3.7 ^a	7.0±0.1 ^{ab}	7.6±0.5 ^{ab}	8.0±0.3 ^a	6.9±0.6 ^b	5.8±0.4 ^c
5.0	53.6±1.9 ^d	73.9±5.0 ^b	64.3±3.0 ^c	77.6±4.0 ^b	89.2±2.12 ^a	11.9±0.7 ^b	12.6±1.1 ^b	12.5±0.7 ^b	11.9±0.2 ^b	15.9±0.1 ^a

Table 3.3. Se removal efficiency by *Lemna* (%). Values are mean ± standard deviation (n=3). Different letters indicate statistically significant differences among treatments with the same Se application according to Duncan's multiple comparison tests ($P < 0.05$).

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
0.0	-	-	-	-	-	-	-	-	-	-
0.5	22.0±0.4 ^c	16.2±1.0 ^d	21.5±0.3 ^c	30.2±1.8 ^a	27.6±1.4 ^b	2.2±0.2 ^c	2.9±0.3 ^b	2.7±0.3 ^{bc}	2.5±0.3 ^{bc}	3.6±0.2 ^a
2.5	11.8±0.3 ^e	13.3±0.3 ^d	15.0±0.1 ^c	17.3±0.8 ^b	23.7±1.0 ^a	1.9±0.1 ^{ab}	2.0±0.6 ^{ab}	2.1±0.1 ^a	1.8±0.2 ^b	1.5±0.1 ^c
5.0	7.1±0.3 ^d	9.8±0.7 ^b	8.6±0.4 ^c	10.3±0.5 ^b	11.9±0.3 ^a	1.6±0.1 ^b	1.7±0.1 ^b	1.7±0.1 ^b	1.6±0.1 ^b	2.1±0.1 ^a

Chapter 3

Table 3.4. Total Se accumulation in *Azolla* ($\mu\text{g}/\text{pot}$). Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the same Se application according to Duncan's multiple comparison tests ($P < 0.05$).

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
0.0	-	-	-	-	-	-	-	-	-	-
0.5	29.2 $\pm$ 0.1 ^a	23.4 $\pm$ 0.3 ^c	25.9 $\pm$ 0.2 ^b	25.5 $\pm$ 0.8 ^b	23.8 $\pm$ 0.8 ^c	2.2 $\pm$ 0.1 ^a	1.6 $\pm$ 0.1 ^b	1.4 $\pm$ 0.1 ^b	1.4 $\pm$ 0.2 ^b	1.5 $\pm$ 0.1 ^b
2.5	88.4 $\pm$ 3.3 ^a	74.8 $\pm$ 2.3 ^b	74.2 $\pm$ 1.2 ^b	65.9 $\pm$ 3.7 ^c	65.6 $\pm$ 1.8 ^c	11.2 $\pm$ 1.7 ^{ab}	12.4 $\pm$ 0.1 ^a	9.4 $\pm$ 0.9 ^{bc}	7.9 $\pm$ 0.6 ^c	8.9 $\pm$ 1.34 ^c
5.0	90.5 $\pm$ 1.2 ^a	86.6 $\pm$ 5.8 ^a	78.4 $\pm$ 4.3 ^a	79.9 $\pm$ 3.3 ^a	80.7 $\pm$ 2.9 ^a	17.3 $\pm$ 1.3 ^a	15.1 $\pm$ 0.5 ^b	15.3 $\pm$ 1.4 ^b	11.6 $\pm$ 0.8 ^c	11.2 $\pm$ 0.1 ^c

Table 3.5. Se removal efficiency by *Azolla* (%). Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the same Se application according to Duncan's multiple comparison tests ($P < 0.05$).

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
0.0	-	-	-	-	-	-	-	-	-	-
0.5	38.9 $\pm$ 0.1 ^a	31.2 $\pm$ 0.4 ^c	34.6 $\pm$ 0.2 ^b	34.0 $\pm$ 1.0 ^b	31.8 $\pm$ 1.0 ^c	3.0 $\pm$ 0.1 ^a	2.2 $\pm$ 0.1 ^b	1.9 $\pm$ 0.1 ^b	1.9 $\pm$ 0.3 ^b	2.0 $\pm$ 0.1 ^b
2.5	23.6 $\pm$ 0.9 ^a	20.0 $\pm$ 0.6 ^b	20.0 $\pm$ 0.3 ^b	17.6 $\pm$ 1.0 ^c	17.5 $\pm$ 0.5 ^c	3.0 $\pm$ 0.5 ^{ab}	3.3 $\pm$ 0.1 ^a	2.5 $\pm$ 0.2 ^{bc}	2.1 $\pm$ 0.2 ^c	2.4 $\pm$ 0.4 ^{bc}
5.0	12.1 $\pm$ 0.2 ^a	11.5 $\pm$ 0.8 ^a	10.4 $\pm$ 0.5 ^a	10.6 $\pm$ 0.4 ^a	10.8 $\pm$ 0.4 ^a	2.3 $\pm$ 0.2 ^a	2.0 $\pm$ 0.1 ^b	2.0 $\pm$ 0.2 ^b	1.5 $\pm$ 0.1 ^c	1.5 $\pm$ 0.1 ^c

Chapter 3

Table 3.6. Bioconcentration factor of Se (BCF_{Se}) determined for *Lemna* and *Azolla*

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
<i>Lemna</i>										
0.5	343	268	397	552	506	31.0	40.8	34.1	37.6	55.6
2.5	280	275	319	355	508	22.7	26.2	26.5	27.1	30.2
5.0	231	261	260	303	333	20.0	21.5	22.4	24.6	33.7
<i>Azolla</i>										
0.5	667	584	603	562	521	50.9	37.3	40.0	32.6	31.2
2.5	392	366	338	301	294	51.9	44.2	42.9	36.5	36.9
5.0	228	202	188	182	187	39.2	32.3	25.1	23.1	24.6

Chapter 3

3.3.5 Zn concentration in *Lemna* and *Azolla*

An increased Zn concentration in both plants was significantly associated with the increasing Zn dose in the culture solution (Fig. 3.4) ($P < 0.01$). The maximum Zn concentrations in *Lemna* and *Azolla* were 3144 and 1709 mg/kg, respectively, when exposed to the highest Zn amount (5.0 mg/L), but at different Se concentrations.




Figure 3.4. Zn content in plants after incubation on different Se and Zn concentrations: (a) *Lemna* grown on Se(IV) and Zn, (b) *Lemna* grown on Se(VI) and Zn, (c) *Azolla* grown on Se(IV) and Zn and (d) *Azolla* grown on Se(VI) and Zn. Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the same Zn application according to Duncan's multiple comparison tests ($P < 0.05$).

For the same Zn application, the Zn concentration in *Lemna* significantly increased with increasing Se(IV) dose ($P < 0.01$), while it remained almost constant with increasing Se(VI) concentrations (Fig. 3.4a and 3.4b). In contrast, the Zn concentration in *Azolla* generally declined with increasing amounts of Se(IV) and Se(VI) in the medium ($P < 0.01$), except for a slight increase observed at 5 mg/L Se(IV) (Fig. 3.4c and 3.4d). In the presence of 5 mg/L Zn, increasing the dose of Se(IV) from 0 to 5 mg/L raised the

Chapter 3

Zn content in *Lemna* from 1769 to 3144 mg/kg ($P < 0.01$), while it diminished the Zn content in *Azolla* from 1709 to 1530 mg/kg ($P = 0.09$).

3.3.6 Zn accumulation, removal and BCF_{Zn}

The total Zn accumulation, Zn removal efficiency and BCF_{Zn} by *Lemna* and *Azolla* are illustrated in Tables 3.S2–3.S6 (see Supplementary Information). The total Zn accumulation in both plants significantly raised with the increase in Zn application, while the Zn removal efficiency decreased remarkably ($P < 0.01$). The largest Zn accumulation and the highest Zn removal efficiencies in *Lemna* and *Azolla* were up to 196 $\mu\text{g/pot}$ and 90.7%, and 151 $\mu\text{g/pot}$ and 64.7%, respectively. The BCF_{Zn} was much higher than the BCF_{Se} in both plants (Table 3.S6). The maximum BCF_{Zn} reached up to 1310 and 843 in *Lemna* and *Azolla*, respectively.

3.3.7 Protein content

The maximum content of true protein in *Lemna* and *Azolla* was 162 and 170 mg/g when exposed to 2.5 mg/L Se(IV), respectively, whereas a protein content of 131 and 159 mg/g was observed when *Lemna* and *Azolla* were grown in the control medium. The inhibition effect of Se and Zn exposure was not reflected in the protein content, except a decrease of protein content in the presence of 5.0 mg/L Se(IV) for *Lemna* ($P = 0.04$) when increasing the Zn application to 5.0 mg/L (Fig. 3.5).




Figure 3.5. Protein content in *Lemna* and *Azolla* after incubation in the different dosages of Se and Zn medium: (a) *Lemna* and (b) *Azolla*. Values are mean $\pm$ standard deviation ($n=6$). Different letters indicate statistically significant differences among treatments with the same Zn application according to Duncan's multiple comparison tests ($P < 0.05$).

Chapter 3

3.3.8 Se speciation in the plants

The plants grown in the presence of 2.5 mg/L of Se medium were selected for Se speciation determination. The protease extraction efficiency of Se in both plants grown in the Se(IV) containing medium ranged from 13% to 40% compared to the total Se content, while it ranged from 63% to 92% in the presence of Se(VI) medium (Table 3.S7). When plants were grown in the Se(IV) medium, the production rate of Se amino acids was higher than when plants were grown in the medium with Se(VI) (Fig. 3.6, Table 3.S8).

Organic Se species such as SeMet, SeMetSeCys and SeCys₂ were the main Se species of identified Se in *Lemna* grown on Se(IV)-enriched medium (Fig. 3.6a), whereas inorganic Se in the form of selenate (approximately 50%) was the predominant Se species in *Lemna* grown on Se(VI)-containing medium (Fig. 3.6b, Table 3.S8). SeMet accounted for the highest proportion (5.9% of total Se) of identified Se species in *Lemna* grown in Se(IV), while SeMetSeCys predominated (4.6% of total Se) in the Se(IV) treatment with Zn.

In *Azolla*, the main species were organic SeMet and SeMetSeCys and inorganic selenate upon Se(IV) exposure, while selenate was detected in *Azolla* as predominant species (92% and 75% in the absence and presence of Zn, respectively) when exposed to Se(VI). Moreover, excepting selenate species in *Azolla*, SeMetSeCys was the most abundant Se species under Se(IV) exposure with Zn, while SeMet has the highest proportion under only Se(IV) exposure (Figs 6c and 6d, Table S8). Apart from the identified Se species, some unidentified Se species were found in both *Lemna* and *Azolla* grown in the presence of Se(IV) (Figs. 3.6a and 3.6c).

Chapter 3




Figure 3.6. Chromatograms of Se speciation in *Lemna* and *Azolla* compared with that of a Se standard solution of 100 µg/L each species: (1) Se-cystine, (2) Se-methylselenocysteine, (3) selenite, (4) Se-methionine, (5) γ-glutamyl-methylselenocysteine, (6) selenate, and (7) γ-glutamyl-selenomethionine. (a) *Lemna* grown on Se(IV) and Zn, (b) *Lemna* grown on Se(VI) and Zn, (c) *Azolla* grown on Se(IV) and Zn and (d) *Azolla* grown on Se(VI) and Zn.

Chapter 3

3.4 Discussion

3.4.1 Se uptake and toxic effects of Se and Zn on *Lemna* and *Azolla*

Changes in the growth rate of plants, as evidenced by biomass production and root length, are a direct indicator of plant toxicity in contaminated environments (Becker, 2013; Duan et al., 2010). In this study, the decrease of biomass production of both *Lemna* and *Azolla* was significantly associated ($P < 0.001$) with increasing Se(IV) dosage in the medium (Figs. 3.1 and 3.2), while no inhibition effect was found at the highest dose of Se(VI) ($P = 0.24$ and 0.11 for *Lemna* and *Azolla*, respectively). The toxic effect of Se(IV) to aquatic plants has been reported previously. Zhong et al. (2016) reported negative impacts of Se(IV) on the chlorophyll fluorescence, starch content and fatty acid content of the duckweed *Landoltia punctata* after exposure to $> 40 \mu\text{mol/L}$ Se(IV) (equivalent to 3.2 mg/L Se). Carvalho and Martin (2001) also recorded that dry biomass of the duckweed *Lemna obscura* Aust. decreased from 50 to 20 mg when the Se(IV) concentration was increased from 1 to 20 mg/L in the cultivation medium. The higher plant tolerance to Se(VI) compared to Se(IV) has also been observed in other plants such as sunflower and maize when cultivated in a hydroponic system (Garousi et al., 2016). The different toxicity of Se(IV) and Se(VI) on plants growth can be explained by the different mobility, bioavailability and metabolic uptake mechanisms (Li et al., 2008). Specifically, Se(VI) is taken up in an active way and easily distributed from roots to shoots (Arvy, 1993), while Se(IV) is absorbed in a faster passive way and quickly converted into organic forms of Se (Fig. 3.6). The organic Se is then incorporated into proteins by replacing sulfur in plant tissues, resulting in malformed selenoproteins (Arvy, 1993; de Oliveira et al., 2017). Moreover, Se-induced oxidative stress also contributes to plant toxicity (Van Hoewyk, 2013). Accordingly, the reduction of Se(IV) to organic Se forms (e.g., SeMet, SeCys₂) may produce additional organic Se metabolites such as selenodiglutathione, which is more toxic than Se(IV), inducing oxidative stress in plants (Van Hoewyk, 2013; Wallenberg et al., 2010).

The higher toxicity of Se(IV) can also be attributed to the higher Se uptake/accumulation in the plant tissues when the two plants were cultivated in Se(IV) containing medium compared to the Se(VI) treatment. The uptake of Se(IV) in both *Lemna* and *Azolla* was higher than that of Se(VI), resulting in Se concentrations in both plants more than 10-fold higher when they were grown in Se(IV)-supplemented

Chapter 3

medium (Fig. 3.3). Similarly, Broyer et al. (1972) reported more accumulation of Se(IV) than Se(VI) by the hyper-accumulator *Astragalus crotolariae* after 7-12 weeks of hydroponic culture. Zhang et al. (Zhang et al., 2003) observed 4 times higher uptake of Se(IV) than Se(VI) by soybean *Glycine max*, while Arvy (1993) reported a similar uptake rate for the two Se species by bean plants (*Phaseolus vulgaris*) during a 3-h uptake experiment. The higher uptake of Se(IV) can be explained by the different metabolism of Se(IV) and Se(VI) in plants. Se(VI) is taken up by plants, reduced to Se(IV) and then converted into organic Se (Van Hoewyk, 2013). The Se(VI) reduction occurs via substitution for sulfate in the ATP sulfurylase reductase system, which is an ATP-consumption process and rate-limiting step (Salt et al., 2002; Van Hoewyk, 2013). The Se(VI) reduction rate is much slower than the uptake rate, resulting in Se(VI) saturation and lower Se uptake by plant, which is supported by the high proportion of Se(VI) species in both *Lemna* and *Azolla* grown in the presence of Se(VI) (Fig. 3.6).

On the other hand, Versini et al. (2016) reported a lower uptake of Se(IV) compared to Se(VI) in the non-accumulator ryegrass grown in hydroponic cultures. Garousi et al. (2016) also observed that the root-to-shoot Se translocation and total Se uptake was lower in Se(IV)-treated plants (maize and sunflower) than in Se(VI)-treated plants. These discrepancies are likely caused by different external environmental factors, such as temperature, light intensity, Se exposure concentration, and medium composition — especially the content of macronutrient (e.g., sulfate and phosphate) or micronutrient elements — as well as the differences in plant species (Kikkert & Berkelaar, 2013). For instance, *Astragalus crotolariae* is Se hyperaccumulator, whereas ryegrass, maize, and sunflower are non-accumulators. The Se hyperaccumulators differ from the non-accumulators in the capacity of reduction of the intracellular concentration of selenocysteine (SeCys) and SeMet (Terry et al., 2000). Non-accumulators incorporate most SeCys and SeMet into proteins with damaging effects to plant functions; while hyperaccumulators metabolize the SeCys primarily into various non-protein selenoamino acids, such as SeMetSeCys, Se-cystathionine and γ -glutamyl-methylselenocysteine to tolerate high concentrations of Se in their cells (Terry et al., 2000). The high tolerance keeps the cell membranes functional and improves the passive uptake of Se(IV), resulting in more accumulation of Se(IV) than that of Se(VI) by Se hyperaccumulators. Besides, Wang et al. (2019) evidenced that tomato (*Solanum lycopersicum* L.) has a higher uptake ability of Se(VI) than Se(IV) at 0.0175-

Chapter 3

0.2998 mg/L Se exposure, while the opposite was observed when Se exposure was higher than 0.2998 mg/L, which partially confirms that the Se exposure concentration may affect the uptake ability of Se by plants.

Additionally, the results of biomass (Fig. 3.1) indicated that Se could counteract the inhibiting effect of Zn on the growth of plants. This is due to the fact that a suitable amount of Se could act as an antioxidant by inhibiting lipid peroxidation and avoiding the formation of reactive oxygen species (ROS). ROS could harm the plant cells. Besides, Se can up-regulate the amount and activity of antioxidant enzymes (e.g., glutathione peroxidases (GPx) and superoxide dismutase (SOD)), resulting in higher ROS scavenging capacity of cells (Schiavon et al., 2017b; Sun et al., 2014), subsequently counteracting the toxicity of Zn.

3.4.2 Simultaneous uptake of Se and Zn

The Se concentration in *Lemna* increased with increasing Zn dosage in the medium, whereas it declined in *Azolla*. These results indicate that Zn promoted the Se uptake in *Lemna*, but inhibited the Se uptake in *Azolla*. Accordingly, the Zn concentration in *Lemna* also raised with raising Se dose in the culture medium, but it decreased in *Azolla*. These findings demonstrate that Se and Zn have synergetic effects in *Lemna*, but antagonistic effects in *Azolla*. The interactions between Se and other elements during plant absorption have been reported, but ambiguous conclusions have been drawn. In line with our results with *Azolla*, previous studies have demonstrated that Se absorption can decrease the uptake of certain elements. For example, Singh and Singh (Singh & Singh, 1978) found that Se application reduced the Zn and Cu concentration in wheat (*Triticum aestivum*). Similarly, the antagonistic effects between Se and other elements such as Mg, K, P, Fe, Cu, and Zn have been demonstrated (Feng et al., 2009) in the fern *Pteris vittata* L. However, other studies showed that Se could improve Zn uptake, which supports part of our findings in *Lemna* (Fig. 3.4). For instance, Arvy (1992) demonstrated that *Helminthia echinoides* grown under field conditions accumulated Se at concentration ranging from 2.05 to 7.90 mg/kg and that Se accumulation was positively correlated with Mn, Zn, Ni, Co, and Cd uptake. Similarly, the foliar application of Se(IV) in turnip (*Brassica rapa* var. *rapa* Linn.) positively affected the uptake of several elements such as Mg, Fe, Zn, Mn, and Cu (Li et al., 2018).

Chapter 3

The increase of Se concentration in *Lemna* with increasing Zn addition (Fig. 3.3) could be partially explained by bioconcentration because of the decline of biomass (Li et al., 2015a). This is supported by the negative correlation between the Se concentration and biomass of *Lemna* exposed to different Zn dosages within each Se application dose, while this correlation was not observed in *Azolla* (Table 3.S9). Similarly, the increment of the Zn concentration in *Lemna* with increasing Se(IV) application may be related to the decrease of biomass (Table 3.S10), which is also evidenced by the significant decrease of Zn accumulation (multiplying biomass by concentration) and Zn removal efficiency by *Lemna* when the Se application increased ($P < 0.01$) (Tables 3.S2 and 3.S3).

Additionally, Se is detoxified by methylation of SeCys₂ and SeMet to SeMetSeCys and methylselenomethionine (Me-SeMet), which cannot be incorporated into proteins, thereby avoiding toxicity and allowing a safe Se accumulation (Gupta & Gupta, 2017). In this study, SeMet accounted for the highest proportion of the identified Se species in *Lemna* grown in Se(IV), while SeMetSeCys predominated in the Se(IV) treatment with Zn (Fig. 3.6). The formation and accumulation of non-toxic SeMetSeCys species from toxic SeCys₂ could have been stimulated by the addition of Zn (probably by stimulating the expression of selenocysteine methyltransferase (SMT)) (Van Hoewyk, 2013), eventually resulting in the enhancement of Se accumulation and the higher Se concentration in *Lemna* in the treatment with Se and Zn.

Azolla, on the other hand, showed an antagonistic effect between the uptake of Se and Zn, which may be related to the higher metal tolerance ability of *Azolla* and the Se mediated detoxification of heavy metals. For *Lemna*, the supplementation of 5.0 mg/L Se(IV) or Zn decreased the dry weight of the biomass by 61% and 29%, respectively, while for *Azolla*, the reduction was only 20% and 6%, respectively (Fig. 3.1). The higher metal tolerance of *Azolla* suggests a detoxification mechanism was triggered by the exposure to Se and Zn. Accumulated metal ions are normally detoxified by phytochelatins (PCs), which are synthesized from glutathione (GSH) during exposure to heavy metals. PCs form a complex with metal ions and sequester them into the vacuole (Yadav, 2010). Research has evidenced that Zn could induce the PCs synthesis in some plants (Tsuji et al., 2002). However, Hawrylak-Nowak et al. (2014) reported that Se reduced the PCs accumulation in the presence of Cd, due to Se interference with the S metabolism and replacement of S in amino-acids, forming Se-

Chapter 3

amino acids (SeMet and SeCys). Se-amino acids are subsequently incorporated into enzymatic proteins, including phytochelatin synthase as this contains cysteine, affecting their catalytic activity. Thus, the replacement of cysteine by SeCys in the phytochelatin synthase probably affects the biosynthesis and accumulation of PCs in the plant tissues (Malik et al., 2012; Wan et al., 2016). The decrease of PCs induced by Se could explain the decrease of the Zn concentration in *Azolla* when Se was supplied. In addition, increasing Se doses cause saturation of lipids and increases membrane stiffness, resulting in lower membrane permeability and less accumulation of micro- and macronutrients in plants (Filek et al., 2010).

Contrasting interactions between the uptake of metal and metalloids by *Lemna* and *Azolla* have been reported for Cu and As. For example, Cu inhibited As uptake in *Azolla caroliniana*, while it stimulated As uptake in *Lemna minor* (Rofkar et al., 2014). Although the combined interactions of Se and Zn uptake in *Lemna* and *Azolla* were demonstrated in this study, the mechanisms are still unclear. Particularly, the reason why Zn and Se showed opposite effects on Se/Zn uptake between *Azolla* and *Lemna* should be further explored. Therefore, further studies using e.g. ^{77}Se isotope should be conducted to elucidate the mechanisms of micronutrient accumulation and Se species transformations in both *Lemna* and *Azolla* (Di Tullo et al., 2015).

3.4.3 Potential of *Lemna* and *Azolla* for wastewater treatment and production of micronutrient-enriched bioproducts

The species of *Lemna* and *Azolla* used in this study can tolerate and accumulate high Se levels, indicating that the two aquatic plants are optimal Se bioaccumulators. The maximum accumulation of Se in *Lemna* and *Azolla* was 1664 and 1139 mg/kg, respectively (Tables 3.2 and 3.4). These values are much higher than the Se accumulation in the duckweed *Landoltia punctata* (785 mg/kg), which was reported to tolerate up to 80 $\mu\text{mol/L}$ Na_2SeO_3 (~6.3 mg/L Se) (Zhong & Cheng, 2016), and also higher than the Se accumulation in *Azolla caroliniana* (less than 1000 mg/kg) exposed to 1 to 10 mg/L Se(IV) (Hassan & Mostafa, 2016). Moreover, the Se and Zn accumulation in *Lemna* increased with increasing Zn exposure, which confirms that *Lemna* can efficiently remove Se and Zn together (Figs. 3.3 and 3.4).

Chapter 3

The BCF provides information on the ability of a plant to accumulate contaminants from polluted water or soil. In general, a BCF larger than 1 suggests that a plant can be considered as a candidate with good phytoextraction efficiency (Li et al., 2015a). The BCF_{Se} values of *Lemna* and *Azolla* were all larger than 1 (Table 3.6). However, the Se removal efficiency by *Lemna* and *Azolla* was not as high as other indicators (e.g., Se content, BCF). The maximum Se(IV) removal efficiencies by *Lemna* and *Azolla* were 30% and 39%, respectively, and only 3% for Se(VI). This may be due to the small amount of biomass (1.0 g fresh weight) transferred into each treatment and to a limiting water surface area in this study, as floating aquatic plants can only root on the water surface. Regarding the lower removal efficiency of Se(VI), it could be also partially attributed to the competition between Se(VI) and sulfate (0.83 mM) in the medium, resulting in a lower ability of the plant to take up and accumulate Se(VI). It should be noted that the sulfate concentration applied in the medium is a normal concentration in wastewater and suitable for plant growth (Li et al., 2008; Mechora et al., 2015). Therefore, the Se removal efficiencies could improve by increasing the biomass concentration or optimizing the geometry of the treatment tank and the chemical composition of wastewater (e.g., by conducting pretreatments to selectively remove S).

Despite the currently low Se removal efficiencies, the high Se(IV) and Zn accumulation capacity (>1000 mg/kg Se and Zn) of *Lemna* and *Azolla*, together with the high tolerance to Se and Zn, the fast growth rate and the easy harvest make these plant species interesting alternatives for the production of micronutrient-enriched feed supplements or fertilizers. Therefore, the plants could also be grown in treatment ponds treating non-Se-rich wastewater (e.g., domestic wastewater), to which Se(IV) is added to obtain Se-enriched feed supplements or fertilizers. In that way, a high-value product is produced from the wastewater, while recovering resources. Particularly a high protein content is beneficial for feed supplements or ingredients. In this study, the true protein content in *Lemna* and *Azolla* were as high as 162 and 170 mg/g (Fig. 3.5), respectively, which is much higher than other plants, such as turnip (*Brassica rapa* var. *rapa* Linn.) and *Codonopsis lanceolata* (Li et al., 2018; Zhu et al., 2017). The crude protein content in turnip was 59.5-93.5 mg/g when 0-200 mg/L Se(IV) was sprayed on the leaves, and the protein content in *C. lanceolata* was 15.4-17.2 mg/g when 0.5-2.0 mg/kg Se(IV) was applied to the soil (Li et al., 2018; Zhu et al., 2017). Besides, it should be noted that the conversion factor 5 (conversion N to protein) applied in this study

Chapter 3

may potentially underestimate the protein content, due to some missing amino acids were not analyzed (e.g., cysteine and tryptophan) (Table 3.S1). Moreover, most of the identified Se species in both *Lemna* and *Azolla* were organic Se forms (around 88-90% and 54-76% of identified Se species, respectively) when grown on Se(IV) containing medium (Fig. 3.6), which are the preferred forms for Se supplementation in animal feed (Zhan et al., 2007). It has to be noted that the sum of identified Se species ranged from 3.2% to 12.2% when the plants were grown on Se(IV) containing medium (Table 3.S7). The low recoveries of Se indicated that Se(IV) is easily metabolized and incorporated into different Se compounds, complicating their extraction and identification. Moreover, the inorganic Se content also seem to be relatively high when plants are exposed to Se(IV) and Se(VI), particularly for use in food or feed. Further product processing through fractionation of the obtained micronutrient-enriched *Lemna* and *Azolla* and/or applying lower Se dosages may help to remove excess inorganic Se. For use as micronutrient-rich feed supplement, a high Se concentration is preferred, but for use as feed ingredient, which will be consumed in higher amounts, a high protein content combined with a lower Se concentration is sufficient. Furthermore, the produced Se- and Zn-enriched bioproducts could be reused and applied as slow-release organic fertilizers in Se/Zn-deficient fields to improve the Se and Zn levels in the soil and the crops growing on these soils. The potential of Se- and Zn-enriched *Lemna* and *Azolla* for this application also requires further investigation, including pot or field trials with relevant crops.

3.4.4 Study limitations and future perspectives

It should be noticed that the nutrient medium in the current study cannot completely mimic real wastewater, which may affect the results. Real wastewaters generally contain a certain amount of organic carbon, bacteria, solids and possibly harmful elements, e.g. heavy metals or an extremely high content of S, P and N. These compounds will interfere with Se uptake. For instance, the high content of organic carbon in wastewater could stimulate the activities of microorganisms, leading to the change of redox potential. Se(VI) can be reduced into Se(IV) or even Se(0) under reducing conditions, therefore affecting Se uptake by plants. Besides, Se(IV) and Se(VI) can be adsorbed by solids existing in real wastewaters and then precipitated, preventing them from being taken up by plants. Very high concentrations of P, S or N (NO_3^- and NO_2^-) in some real wastewaters would largely reduce the Se uptake by plants,

Chapter 3

as these elements could compete for transporters with Se during the plant uptake phase. Therefore, growing *Lemna* and *Azolla* in real wastewater is necessary for validation of the results obtained in this study.

Safety issues are a main concern when using bioproducts produced from real wastewater as biofertilizer or feed supplement, as these biomaterials may harbor contaminants, such as heavy metals, pathogens, and organic pollutants, and ultimately pose threats to human/animal health and the natural environment. A thorough risk assessment of using these bioproducts will be needed in the future prior to upscaling production on real wastewater. In future cases or applications, wastewater pretreatments may need to be considered to selectively remove any harmful components, such as heavy metals, prior to the production of Se/Zn-containing biomaterials. However, for wastewaters with a low Se/Zn content, such as domestic wastewater, not having serious issues due to absence of heavy metals and only containing suitable concentrations of nutrients, such as N and P, Se/Zn can be externally added to produce an added-value product.

3.5 Conclusions

This study investigated the accumulation and combined effects of Se and Zn in two aquatic plants, *Lemna* and *Azolla*, grown in hydroponic culture. The results demonstrated that Se(IV) is more toxic than Se(VI) for both plant species investigated, as evidenced by the considerable decrease of biomass content and root length when exposed to Se(IV) rich medium. Both aquatic plants took up around 10 times more Se(IV) than Se(VI) from the medium. The Se accumulation and removal efficiency by *Lemna* increased with increasing Zn dosage in the medium, whereas it declined in *Azolla*, suggesting a synergetic effect in *Lemna*, but an antagonistic effect in *Azolla*. Both *Lemna* and *Azolla* can tolerate and accumulate high levels of Se(IV) and Zn, which, combined with the observed transformation to organic species, high protein content and rapid plant growth, makes them good candidates for the production of Se- and Zn-enriched biomass that may be used as crop fertilizers or protein-rich feed supplements or ingredients.

Chapter 3

Supplementary Information

Table 3.S1. amino acid profile of *Lemna* and *Azolla*

Amino Acid	<i>Lemna</i>		<i>Azolla</i>	
	g/100g	± Stdev	g/100g	± Stdev
Aspartic acid	1.85	0.04	1.32	0.11
Glutamic acid	1.82	0.03	2.49	0.23
Asparagine	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>
Serine	0.89	0.02	0.71	0.05
Glutamine	<i>N.D.</i>	<i>N.D.</i>	<LOQ*	<LOQ*
Histidine	0.42	0.00	0.26	0.01
Glycine	0.97	0.01	0.71	0.05
Threonine	0.90	0.02	0.68	0.06
Citrulline	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>
Arginine	1.07	0.01	0.84	0.07
Alanine	1.23	0.02	0.87	0.08
Tyrosine	0.68	0.01	0.51	0.04
Valine	0.89	0.01	0.64	0.06
Methionine	0.30	0.01	0.16	0.02
Phenylalanine	0.98	0.00	0.67	0.06
Isoleucine	0.78	0.02	0.57	0.06
Ornithine	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>
Leucine	1.50	0.01	1.07	0.09
Lysine	1.01	0.02	0.58	0.05
Hydroxyproline	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>
Proline	1.04	0.01	2.23	0.20
Total amino acid	16.51	0.18	14.37	1.04
Total N	3.23	0.08	3.18	0.06
*Protein content	16.16	0.50	15.89	0.32
**Crude protein content	20.20	0.50	19.87	0.38

N.D.= not detected; *LOQ* = limit of detection

*calculated by multiplying the TN by 5.0 (this study)

**calculated by multiplying the TN by 6.25 (typical method)

Chapter 3

Table 3.S2. Total Zn accumulation in *Lemna* for each pot ($\mu\text{g}/\text{pot}$). Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the same Zn application according to Duncan's multiple comparison tests ($P < 0.05$).

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
0.0	-	61.2 $\pm$ 2.8 ^a	79.8 $\pm$ 2.6 ^a	120 $\pm$ 4.3 ^a	156 $\pm$ 8.1 ^c	-	61.2 $\pm$ 2.8 ^a	79.8 $\pm$ 2.6 ^c	120 $\pm$ 4.3 ^b	156 $\pm$ 8.1 ^b
0.5	-	32.5 $\pm$ 3.5 ^c	69.7 $\pm$ 2.0 ^b	83.3 $\pm$ 3.8 ^c	175 $\pm$ 3.0 ^a	-	55.5 $\pm$ 6.4 ^a	101 $\pm$ 3.9 ^b	160 $\pm$ 8.9 ^a	196 $\pm$ 4.8 ^a
2.5	-	38.8 $\pm$ 0.3 ^b	60.3 $\pm$ 2.8 ^c	107 $\pm$ 2.1 ^b	163 $\pm$ 2.7 ^{bc}	-	63.2 $\pm$ 3.3 ^a	107 $\pm$ 1.8 ^{ab}	158 $\pm$ 2.0 ^a	184 $\pm$ 5.7 ^a
5.0	-	31.7 $\pm$ 2.5 ^c	55.8 $\pm$ 2.9 ^c	112 $\pm$ 1.1 ^b	168 $\pm$ 6.4 ^{ab}	-	68.0 $\pm$ 3.5 ^a	111 $\pm$ 4.3 ^a	154 $\pm$ 4.8 ^a	185 $\pm$ 7.0 ^a

Table 3.S3. Zn removal efficiency by *Lemna* for each pot (%). Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the same Zn application according to Duncan's multiple comparison tests ($P < 0.05$).

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
0.0	-	81.6 $\pm$ 3.6 ^a	53.2 $\pm$ 1.7 ^a	32.1 $\pm$ 1.2 ^a	20.8 $\pm$ 1.1 ^c	-	81.6 $\pm$ 3.7 ^a	53.2 $\pm$ 1.7 ^c	32.1 $\pm$ 1.2 ^b	20.8 $\pm$ 1.1 ^b
0.5	-	43.3 $\pm$ 4.7 ^c	46.4 $\pm$ 1.4 ^b	22.2 $\pm$ 1.02 ^c	23.4 $\pm$ 0.4 ^a	-	74.0 $\pm$ 8.5 ^a	67.7 $\pm$ 2.6 ^b	42.6 $\pm$ 2.4 ^a	26.1 $\pm$ 0.6 ^a
2.5	-	51.8 $\pm$ 0.4 ^b	40.2 $\pm$ 1.8 ^c	28.7 $\pm$ 0.6 ^b	21.7 $\pm$ 0.4 ^{bc}	-	84.2 $\pm$ 4.4 ^a	71.7 $\pm$ 1.2 ^{ab}	42.2 $\pm$ 0.5 ^a	24.6 $\pm$ 0.8 ^a
5.0	-	42.3 $\pm$ 3.3 ^c	37.2 $\pm$ 1.9 ^c	29.9 $\pm$ 0.3 ^b	22.5 $\pm$ 0.2 ^{ab}	-	90.7 $\pm$ 4.7 ^a	74.0 $\pm$ 2.9 ^a	41.2 $\pm$ 1.3 ^a	24.7 $\pm$ 0.9 ^a

Chapter 3

Table 3.S4. Zn accumulation in *Azolla* for each pot ($\mu\text{g}/\text{pot}$). Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the same Zn application according to Duncan's multiple comparison tests ($P < 0.05$).

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
0.0	-	48.5 $\pm$ 3.6 ^a	66.1 $\pm$ 3.2 ^a	118 $\pm$ 1.2 ^a	151 $\pm$ 2.5 ^a	-	48.5 $\pm$ 3.6 ^a	66.1 $\pm$ 3.2 ^a	119 $\pm$ 1.2 ^a	151 $\pm$ 2.5 ^a
0.5	-	30.7 $\pm$ 3.2 ^c	46.3 $\pm$ 3.3 ^c	74.7 $\pm$ 6.0 ^c	119 $\pm$ 9.4 ^c	-	29.8 $\pm$ 1.3 ^b	47.2 $\pm$ 2.5 ^b	88.7 $\pm$ 6.7 ^b	125 $\pm$ 10.0 ^b
2.5	-	35.7 $\pm$ 1.3 ^{bc}	49.0 $\pm$ 3.0 ^c	72.2 $\pm$ 4.6 ^c	107 $\pm$ 5.1 ^c	-	26.5 $\pm$ 0.7 ^b	39.7 $\pm$ 1.3 ^c	66.7 $\pm$ 0.4 ^d	128 $\pm$ 5.8 ^b
5.0	-	36.8 $\pm$ 2.4 ^b	59.2 $\pm$ 2.8 ^b	95. 7 $\pm$ 7.7 ^b	133 $\pm$ 1.3 ^b	-	28.5 $\pm$ 1.4 ^b	42.7 $\pm$ 1.7 ^{bc}	79.7 $\pm$ 0.4 ^c	114 $\pm$ 10.5 ^b

Table 3.S5. Zn removal efficiency by *Azolla* for each pot (%). Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the same Zn application according to Duncan's multiple comparison tests ($P < 0.05$).

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
0.0	-	64.7 $\pm$ 4.8 ^a	44.1 $\pm$ 2.2 ^a	31.6 $\pm$ 0.3 ^a	20.2 $\pm$ 0.3 ^a	-	64.7 $\pm$ 4.8 ^a	44.1 $\pm$ 2.2 ^a	31.6 $\pm$ 0.3 ^a	20.2 $\pm$ 0.3 ^a
0.5	-	41.0 $\pm$ 4.2 ^c	30.9 $\pm$ 2.2 ^c	19.9 $\pm$ 1.6 ^c	16.0 $\pm$ 1.3 ^c	-	39.8 $\pm$ 1.7 ^b	31.5 $\pm$ 1.6 ^b	23.7 $\pm$ 1.8 ^b	16.7 $\pm$ 1.3 ^b
2.5	-	47.6 $\pm$ 1.7 ^{bc}	32.7 $\pm$ 2.0 ^c	19.3 $\pm$ 1.2 ^c	14.3 $\pm$ 0.7 ^c	-	35.3 $\pm$ 0.94	26.4 $\pm$ 0.8 ^c	17.8 $\pm$ 0.1 ^d	17.1 $\pm$ 0.8 ^b
5.0	-	49.0 $\pm$ 3.1 ^b	39.4 $\pm$ 1.8 ^b	25.5 $\pm$ 2.0 ^b	17.8 $\pm$ 0.6 ^b	-	38.0 $\pm$ 1.8 ^b	28.5 $\pm$ 0.7 ^{bc}	21.3 $\pm$ 0.1 ^c	15.2 $\pm$ 1.4 ^b

Chapter 3

Table 3.S6. Bioconcentration factor of Zn (BCF_{Zn}) for *Lemna* and *Azolla* cultivated in different concentrations of Se and Zn.

Se treatment (mg/L)	Se(IV)					Se(VI)				
	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0	Zn 0.0	Zn 0.5	Zn 1.0	Zn 2.5	Zn 5.0
<i>Lemna</i>										
0.0	-	1064	905	547	354	-	1064	905	547	354
0.5	-	696	818	428	413	-	1085	897	625	397
2.5	-	1069	853	616	465	-	1086	979	617	416
5.0	-	1310	1128	877	629	-	1106	973	602	387
<i>Azolla</i>										
0.0	-	843	748	540	342	-	843	748	540	342
0.5	-	633	571	340	269	-	662	538	373	276
2.5	-	766	519	335	244	-	577	462	320	266
5.0	-	833	625	446	306	-	609	467	331	232

Chapter 3

Table 3.S7. Extraction efficiency of Se after protease hydrolysis and the identification rate of Se by anion exchange column

Treatments		Extraction efficiency (%)	Column recovery (%)
Se(IV)	<i>Lemna</i>	37	41
	<i>Azolla</i>	13	43
Se(IV)+Zn	<i>Lemna</i>	31	39
	<i>Azolla</i>	26	19
Se(VI)	<i>Lemna</i>	63	122
	<i>Azolla</i>	81	130
Se(VI)+Zn	<i>Lemna</i>	67	113
	<i>Azolla</i>	92	107

Chapter 3

Table 3.S8. Se species concentration found in *Lemna* and *Azolla* grown on 2.5 mg/L Se at the absence of Zn or at the presence of 5.0 mg/L Zn (percentage of identified Se species comparison with total Se in plants)

Treatments		Se species concentration (mg/kg)					
		SeCys ₂	SeMetCys	Se(IV)	SeMet	Se(VI)	Unknown
<i>Duckweed</i>	Se(IV)	5.4 (0.8%)	23.3 (3.3%)	5.6 (0.8%)	41.1 (5.9%)	9.9 (1.4%)	Detected
	Se(IV)+Zn	10.2 (0.8%)	58.3 (4.6%)	15.1 (1.2%)	34.7 (2.7%)	11.6 (0.9%)	Detected
	Se(VI)	1.4 (2.4%)	1.7 (3.1%)	1.8 (3.2%)	12.5 (22.1%)	28.1 (49.7%)	ND
	Se(VI)+Zn	1.0 (1.8%)	3.2 (5.6%)	2.0 (3.5%)	6.2 (11.0%)	30.1(53.5%)	ND
<i>Azolla</i>	Se(IV)	2.1 (0.3%)	6.0 (0.9%)	3.7 (0.5%)	24.5 (3.5%)	11.6 (1.7%)	Detected
	Se(IV)+Zn	2.9 (0.4%)	4.4 (0.6%)	1.8 (0.3%)	3.8 (0.5%)	10.9 (1.5%)	Detected
	Se(VI)	0.8 (0.7%)	ND	1.5 (1.3%)	2.1 (1.9%)	101 (91.7%)	ND
	Se(VI)+Zn	0.2 (0.3%)	ND	ND	ND	68.9 (75.5%)	Detected

Chapter 3

Table 3.S9 Pearson's correlation coefficients between biomass and Se concentration in *Lemna* and *Azolla* within each Se application. Significant differences are indicated with * for *P*-value <0.05 and ** for *P*-value <0.01.

		Se application (mg/L)		
		0.5	2.5	5.0
<i>Lemna</i>	Se(IV)	-0.54*	-0.07	0.51*
	Se(VI)	-0.67*	-0.48*	-0.81**
<i>Azolla</i>	Se(IV)	-0.80*	0.12	-0.35
	Se(VI)	-0.01	0.49	-0.33

Table 3.S10. Pearson's correlation coefficients between biomass and Zn concentration in *Lemna* and *Azolla* within each Zn application. Significant differences are indicated with * for *P*-value <0.05 and ** for *P*-value <0.01.

		Zn application (mg/L)			
		0.5	1.0	2.5	5.0
<i>Lemna</i>	Se(IV)	-0.46	-0.82**	-0.86**	-0.80**
	Se(VI)	-0.49	0.12	0.57*	-0.33
<i>Azolla</i>	Se(IV)	0.55	0.37	0.13	0.50
	Se(VI)	0.62*	-0.002	0.22	-0.26

Chapter 4 Valorization of Se-enriched sludge and duckweed generated from wastewater as micronutrient biofertilizers

This chapter was redrafted after:

Li, J., Otero-Gonzalez, L., Parao, A., Tack, P., Folens, K., Ferrer, I., Lens, P.N.L., Du Laing, G. 2021. Valorization of Se-enriched sludge and duckweed generated from wastewater as micronutrient biofertilizers. *Chemosphere*, 130767.

Chapter 4

Abstract

In this study, the potential of Se-enriched biomaterials (sludge and duckweed) as slow-release Se biofertilizers was evaluated through pot experiments with and without planted green beans (*Phaseolus vulgaris*). The Se concentration in the bean tissues was 1.1–3.1 times higher when soils were amended with Se-enriched sludge as compared to Se-enriched duckweed. The results proved that the Se released from Se-enriched biomaterials was efficiently transformed to health-beneficial selenoamino acids (e.g., Se-methionine, 76–89%) after being taken up by beans. The Se-enriched sludge, containing mainly elemental Se, is considered as the preferred slow-release Se biofertilizer and an effective Se source to produce Se-enriched crops for Se-deficient populations, as shown by the higher Se bioavailability and lower organic carbon content. This study could offer a theoretical reference to choose an environmental-friendly and sustainable alternative to conventional mineral Se fertilizers for biofortification, avoiding the problem of Se losses by leaching from chemical Se fertilizers while recovering resources from wastewater. This could contribute to the driver for a future circular economy.

Keywords: Biofortification, green beans, resource recovery, Se-enriched biomaterials, selenium bioavailability.

Chapter 4

4.1 Introduction

Biofortification is being explored as a possible solution for Se deficiency (Lavu et al., 2012; Li et al., 2010; Thavarajah et al., 2008). Selenium biofortification of food crops is practiced in Se-deficient regions of different countries, such as Finland, by adding inorganic Se fertilizer to soils (Bañuelos et al., 2016). Although the application of inorganic Se fertilizer can be an effective way to produce Se-enriched food and feed products, secondary contamination of soil and water can occur due to the low utilization rate of Se by crops (Wang et al., 2018). Therefore, other sources of Se, such as organic Se, could be useful as an alternative soil additive to produce Se-enriched crops. This approach would not only increase the organic matter content in soils, but also enhance Se uptake by plants due to the slower release of Se to the pore water (Wang et al., 2018). In this context, the synchronization of the release of Se from decomposing organic matter with crop uptake is beneficial to avoid leaching problems and to reduce the amount of Se that should be applied.

For that purpose, some Se-rich organic materials produced in seleniferous areas have already been investigated as potential Se sources to provide a suitable amount of Se for plant uptake, e.g., Se-rich animal manure or plant residues (Ajwa et al., 1998; Bañuelos et al., 2016; Dhillon et al., 2007; Stavridou et al., 2011; Wang et al., 2018). Specifically, the supplementation of 4.2 g Se-enriched wheat straw and pak choi into Eum-Orthic Anthrosol soil significantly increased the Se content in the shoot of pak choi from 0.16 mg/kg (in the untreated control) to 0.40 and 2.23 mg/kg, respectively (Wang et al., 2018). The total Se concentration in the edible parts of broccoli and carrot increased from 0.5 to 3.5 mg/kg, and from 0.3 to 2.3 mg/kg, respectively, after soil application of Se-enriched *Stanleya pinnata* (Bañuelos et al., 2015). All of these materials are so far generated by crop growth on seleniferous soils, while the valorization of organic Se-containing biomaterials produced from wastewater as Se sources for plant uptake has not been reported yet.

However, excess of Se in the aquatic environment is also frequently observed. Some wastewaters have a high Se content as a result of both agricultural and industrial activities (Lim & Goh, 2005). Nutrient-rich wastewater loaded also with Se, such as aquaculture wastewater (Han et al., 2020) and agricultural drainage (CH2MHill, 2010), may thus also serve as a potential Se source from which the Se may be recovered and

Chapter 4

valorized to produce slow-release organic biofertilizers. Products generated from Se-bearing wastewater normally have a high Se content, such as Se-enriched granular sludge (Staicu et al., 2015a) and aquatic plants after phytoremediation and phytoextraction (Li et al., 2020b). Thus, it can also be explored whether the Se contained in these products can also be potentially used for biofortification purposes as an organic nutrient-enriched organic fertilizer to improve Se levels in soils for plant uptake. This approach would be beneficial to save Se resources and mitigate Se contamination, meanwhile also avoiding introduction of new Se contamination into the environment through the use of inorganic Se as fertilizer in biofortification. It would also contribute to the worldwide drive for resource recovery and circular economy.

Therefore, the main objective of this study was to assess whether Se-enriched duckweed and sludge produced on wastewater can be an efficient Se source for elevating Se contents in food crops. Specifically, the study aims to (1) to preliminarily screen five crops for Se accumulation and select one suitable crop; (2) to study the release of plant-available Se from Se-enriched duckweed and sludge produced through ecotechnologies after amending them into two types of soils during a long-term period; (3) to monitor the effect of soil amendment with these two Se-enriched biomaterials on green beans (*Phaseolus vulgaris*), in terms of their growth, production and Se accumulation; and (4) to assess the potential of these two Se-enriched biomaterials as Se biofertilizers to improve the Se content in green beans. *Phaseolus vulgaris*, known as the common bean, is an annual plant, grown worldwide for its edible beans. Its leaf is occasionally used as a vegetable, and straw as fodder for animals (Azhar et al., 2020). The relatively high protein content in beans makes them a good candidate for Se biofortification, as Se is preferably accumulated as an organic form in proteins, i.e., as selenoaminoacids (Li et al., 2020b). Accordingly, Se-biofortified beans could simultaneously improve human and animal Se and protein intake.

4.2 Materials and methods

4.2.1 Soil collection and characterization

Two types of non-contaminated soils, classified as sandy and loamy, were collected at a depth of 0–20 cm from fields in Evergem (51°6′57" N, 3°39′40" E) and Wortegem-Petegem (50°50′20" N, 3°33′22" E), Belgium, respectively. The soils were dried,

Chapter 4

homogenized and passed through a 2-mm sieve mesh. The physicochemical properties of the soils were analyzed according to Van Ranst et al. (1999).

For soil pH-H₂O determination, 10.0 g of the collected soil was added into 50 mL deionized water (DI) and mixed thoroughly, followed by a 16-h equilibrium. The pH in the extract was determined using a pH-meter (Orion Star A211, Thermo fisher scientific, Waltham, MA, USA). Soil pH-KCl was measured after 1M KCl extraction for 10 min with a soil-to-solution ratio of 1:2.5. Electrical conductivity (EC) was determined after extraction with water (soil:water = 1:5) for 0.5 h and analyzed using a conductivity meter (Orion Star A211, Thermo fisher scientific, Waltham, MA, USA). Cation exchange capacity (CEC) was determined through ammonium acetate percolation, followed by 1 M KCl percolation. The extract was distilled and titrated with 0.1 M HCl. Organic matter (OM) was quantified by measuring the weight loss after 400 °C combustion. Water-holding capacity (WHC) was measured by soaking the soil samples in water for 2 h and then draining for 2 h. Soil texture was measured via the hydrometer method. Specifically, 40 g of the soil was added into 250 mL DI water and 100 mL dispersing agent prepared by (NaPO₃)₆ and Na₂CO₃, and shaken thoroughly overnight. The mixture was transferred into a 1-L sedimentation cylinder located in a 20 °C water bath and measured soil texture was determined based on evolution of density, measured using a hydrometer for 24 h. Total elements concentrations of the soils were determined after the aqua regia digestion using an inductively coupled plasma-optical emission spectrometer (ICP-OES; Varian Vista MPX, Palo Alto, USA), while Se was determined using inductively coupled plasma-mass spectrometer (ICP-MS, ELAN DRC-e, PerkinElmer, Waltham, MA, USA). Soil available Se was determined via 0.016 M KH₂PO₄ extraction as described by Ngigi et al. (2019). The results of soil characterization showed that the loamy soil had a higher electrical conductivity (EC), cation exchange capacity (CEC) and organic matter (OM) content than the sandy soil (Table 4.1).

Chapter 4

Table 4.1. Physicochemical properties and (trace) element content of the tested sandy and loamy soil (mean $\pm$ standard deviation; n=3, except for texture).

	Loamy soil	Sandy soil
pH-KCl	6.14 $\pm$ 0.14	6.45 $\pm$ 0.02
pH-H ₂ O	6.92 $\pm$ 0.08	7.07 $\pm$ 0.12
EC (dS/m)	0.175 $\pm$ 0.011	0.035 $\pm$ 0.002
CEC (cmol/kg)	7.38 $\pm$ 0.07	1.79 $\pm$ 0.18
OM (g/kg)	78.8 $\pm$ 0.0	21.4 $\pm$ 0.0
WHC (g H ₂ O/g dry soil)	0.39	0.33
<i>Texture</i>		
Sand (%)	53.7	91.5
Silt (%)	40.2	6.0
Clay (%)	6.1	2.5
<i>Elements (mg/kg)</i>		
Total Se	0.24 $\pm$ 0.03	0.10 $\pm$ 0.01
Available Se	0.013 $\pm$ 0.001	0.030 $\pm$ 0.001
Total P	1464 $\pm$ 52	300 $\pm$ 6
Total S	359 $\pm$ 11	76 $\pm$ 10
Total Cu	29.7 $\pm$ 0.5	2.7 $\pm$ 0.1
Total Zn	119 $\pm$ 3	8.5 $\pm$ 0.5
Total Ca	4285 $\pm$ 154	1103 $\pm$ 30
Total Mg	1866 $\pm$ 52	489 $\pm$ 31
Total Fe	10749 $\pm$ 136	3612 $\pm$ 104
Total Al	8420 $\pm$ 472	3772 $\pm$ 270

4.2.2 Crops screening experiment

Five crops, pak choi (*Brassica chinensis*), lettuce (*Lactuca sativa*), wheat (*Triticum aestivum*), maize (*Zea mays*) and beans (*Phaseolus vulgaris*), were screened for their capacity to bioaccumulate Se under different soil exposures. A quantity of Na₂SeO₄ stock solution equivalent to 0, 1.0, 3.0 and 5.0 mg Se/kg soil containing 100 mg/kg NPK fertilizer were sprayed onto 7.5 kg dried sandy soil for the growth of five crops. The treated soils were homogenized and divided into 0.5 kg in each pot to stabilize for two weeks for planting crops. All experiments were conducted in triplicate.

Chapter 4

Seeds of the five crops were placed in Petri dishes filled with wet mineral vermiculite and incubated at 27 °C to stimulate germination. Four germinated seeds were transplanted into each pot containing the 0.5 kg treated soils and then grew in the indoor conditions with artificial light (2000 lux, 36W) placed 50 cm above of the plants (16 h of light and 8 h of darkness). The soil moisture during the growing period was maintained at approximately 80% of the water holding capacity by weighing the pots every day and refilling with an appropriate amount of deionized (DI) water.

The plants were harvested after 6 weeks of growth, cleaned with DI water, separated into different tissues (root, shoot, leaf and seed) and dried in an oven at 60 °C until constant weight. The dry biomass weight and total Se concentration of each tissue of the plants were determined.

4.2.3 Valorization of Se-enriched biomaterials as Se biofertilizers

4.2.3.1 Preparation of Se-enriched biomaterials (Se-enriched duckweed and sludge)

The duckweed was collected from a natural freshwater canal in Delft (The Netherlands) and cultivated in a Hoagland solution with 5.0 mg/L of Se added as Na_2SeO_4 . The Se-enriched duckweed was harvested after 7 days of cultivation, oven-dried and stored for subsequent fertilization experiments. Further details on the Se-enriched duckweed production and Se removal have been described in Chapter 3. The Se-enriched sludge was generated as described by Staicu et al. (2015b). The sludge collected from a full-scale upflow anaerobic sludge blanket reactor treating pulp and paper wastewater in Eerbeek (The Netherlands) was added to an oxygen-free growth medium with 5.0 mg/L of Se in the form of Na_2SeO_4 and incubated for 14 days at 30 °C. Afterwards, the sludge enriched with Se was settled, separated and oven-dried. The supernatant was filtered and measured for total Se content in order to quantify the Se removal efficiency. The result showed that up to 93% of Se was removed from wastewater (Fig. 4.S1). The generated sludge samples were milled and passed through a 0.45-mm sieve mesh for further fertilization experiments. The generated duckweed was pulverized using an electrical grinder equipped with a 0.50-mm sieve. The Se-enriched duckweed and sludge contained 209 mg Se/kg dry weight and 314 mg Se/kg dry weight, respectively. A more complete chemical composition of the Se-enriched duckweed and sludge is presented in Table 4.S1.

Chapter 4

4.2.3.2 Selenium availability in the non-planted soils through perennial monitoring

The Se availability in the two soil types amended with the biomaterials was studied during a long-term incubation trial (around 15 months, 471 days). The Se-enriched biomaterials were applied to the soils in an amount equivalent to 1.0 and 5.0 mg Se/kg soil. The biomaterials were thoroughly mixed with 0.5 kg sandy or loamy soil, respectively, and placed into plastic bags within plastic pots (10 cm in height and 10 cm in diameter) during the entire incubation. Non-amended soils prepared similarly served as blank (noted as Blank). The following abbreviations are used indicating the amendments in this study: Se-enriched duckweed 1.0 mg Se/kg soil (DW1) and 5.0 mg Se/kg soil (DW5) and Se-enriched sludge 1.0 mg Se/kg soil (SL1) and 5.0 mg Se/kg soil (SL5).

Specifically, the entire incubation consisted of two monitoring periods by collecting soil pore water: 6 weeks at the beginning of the incubation period (first monitoring period) and 4 weeks at the end of the incubation period (second monitoring period, 13.5 months after the first incubation period). In each monitoring period, soil moisture was kept at 80% of the water holding capacity and the soil pots were placed indoor at room temperature and under natural light conditions. After the first monitoring period, due to the limited lab space for storage and long-term incubation, the amended soils were partially air-dried, sealed in plastic bags and stored for 13.5 months at room temperature until the second monitoring period (Egene et al., 2018). It should be noted that the plastic bags were not sealed tightly, and the soils were not completely dry, so microbial activity was still present in the soil. In the first monitoring period, soil pore water was collected twice per week by using Rhizon soil moisture samplers (Eijkelkamp Agrisearch Equipment, Giesbeek, the Netherlands) and analyzed for total Se concentration, pH, total carbon (TC) and total organic carbon (TOC) (Egene et al., 2018). In the second monitoring period, deionized (DI) water was added to the pots to bring the soil moisture again to 80% of the water holding capacity. Soil pore water was extracted weekly using the Rhizon samplers and analyzed for total Se concentration.

4.2.3.3 Selenium bioavailability in planted soils

Similar to the non-planted experiment, an amount equivalent to 1.0 and 5.0 mg Se/kg soil of Se-enriched duckweed (DW1 and DW5) and sludge (SL1 and SL5) was applied

Chapter 4

to 0.5 kg loamy and sandy soils. For the control experiment, the same amount of non-enriched duckweed and sludge as in the Se-enriched biomaterials amendments was applied to the two types of soil (noted as control-DW1, control-DW5, control-SL1 and control-SL5). Soils without any biomaterial amendment served as blank soil (noted as Blank). In each treatment, inorganic fertilizer (NPK) and the Se-enriched/control biomaterials were uniformly mixed with 0.5 kg sandy or loamy soil. Pots were placed indoors (at 24 °C, 53% relative humidity and 100 $\mu\text{mol}/\text{m}^2/\text{s}$ light intensity) with 80% of water holding capacity and pre-incubated for one week before transferring 4 seedlings of green beans (*Phaseolus vulgaris*) previously cultivated in trays with wet vermiculite for one week. The scheme of this experiment is presented in Figure 4.S2. All experiments were conducted in triplicate. The soil pore water was collected every week through Rhizon extraction for total Se measurement. The bean plants were harvested after 6 weeks of growth, washed and separated into different tissues (root, stem, leaf and seed) for biomass and Se concentration analysis. Se speciation analysis was performed on selected bean seeds (at the DW1 and SL1 amendment for sandy soil, and sandy soil blank) after lyophilization.

4.2.4 Analytical methods

4.2.4.1 Analysis of total Se concentration in plants

For the determination of total Se in plants, 0.3 g of dry samples were weighed into a digestion vessel followed by the addition of 10 mL of concentrated ultrapure HNO_3 . The tubes were sonicated for 1 h, then placed in a microwave oven (CEM Mars 6, Matthews, NC, USA) for digestion with the following program: ramp to 180 °C in 25 min and holding for 20 min at 1200 W power. The digested samples were diluted to 50 mL with Milli-Q water for Se measurement using inductively coupled plasma-mass spectrometry (ICP-MS, ELAN DRC-e, PerkinElmer, Waltham, MA, USA). Internal standards (10 $\mu\text{g}/\text{L}$ ^{103}Rh and ^{69}Ga) and an external multi-element standard solution were used during ICP-MS analysis to validate the accuracy of Se measurements. White clover samples (BCR-42, 6.7 ± 0.25 mg/kg) were included as certified reference materials in each analytical batch as quality control with recoveries of 97 (± 7)%.

4.2.4.2 Selenium speciation analysis

Chapter 4

Selenium speciation in the plant samples (Se-enriched duckweed and Se-enriched bean seeds) was analyzed as described in Chapter 3.

Selenium speciation in the solid sludge was determined by extended X-ray absorption fine structure (EXAFS) and X-ray absorption near edge structure (XANES) (Lenz et al., 2008). To prepare the samples for analysis, dried sludge samples and reference materials of different oxidation numbers were ground to fine powders, mixed with cellulose and pressed into pellets to allow a straightforward sample handling during the experiments. The amount of sludge and reference powders used for each sample was determined using the calculated optimum amount of Se per sample. The EXAFS/XANES spectra were collected over the Se-K (12.658 keV) absorption edge on the sludge sample at the DUBBLE beamline (BM26A) at the European Synchrotron and Radiation Facility (ESRF, Grenoble, France). Measurement for samples was carried out in transmission mode at room temperature. EXAFS data were processed using the Viper software package (Klementev, 2001). XANES data was processed by performing a pre- and post-edge normalization, followed by linear combination analysis using the reference compound spectra.

4.2.4.3 pH, TC and TOC analysis of soil extract

The pH of the soil extracts was determined by using a pH-meter (Orion Star A211, Thermo fisher scientific, Waltham, MA, USA). 2.0 mL of Rhizon extract from each pot was diluted with Milli-Q water to obtain 20 ml volume for total carbon (TC) and total organic carbon (TOC) measurement through a TOC-analyzer (TOC-5000, Shimadzu, Tokyo, Japan) as described by Eugene et al. (2018).

4.2.5 Estimated daily intake and health risk assessment of the bean seeds

The estimated daily intake of Se (EDI, $\mu\text{g/kg/day}$) was calculated using equation (1):

$$EDI = \frac{C_{Se} \times C_{factor} \times D_{intake}}{B_{weight}} \quad (1)$$

Where: C_{Se} is the Se concentration in the bean seeds (mg/kg), C_{factor} represents the conversion factor from fresh weight to dry weight (0.075, calculated by dry weight/fresh weight in this study), D_{intake} (g) and B_{weight} (kg) represent the daily intake of beans and the average body weight (BW), respectively. According to the daily food consumption

Chapter 4

issued by the European Food Safety Authority (EFSA), the chronic legume consumption (D_{intake}/B_{weight}) is 0.55 g/kg BW per day for adults and 1.01 g/kg BW per day for children in Belgium (EFSA).

To assess the human health risk of Se-enriched bean seeds obtained in this study, it is necessary to calculate the level of human exposure to those seeds. The health risk index (HRI) for Se was calculated using equation (2):

$$HRI = \frac{EDI}{RfD} \quad (2)$$

Where: EDI is the estimated daily intake and RfD is the reference oral dose. According to the USEPA, the RfD value for Se is 5.0 µg/kg /day (IRIS, 2006). An estimated value of HRI <1 indicates no evident health risk to the exposed population. Otherwise, it may raise health risks (Sihlahla et al., 2019).

4.2.6 Statistical analysis

Descriptive statistics were performed using Sigma plot 13, Excel 2016 and SPSS 20.0. Results are expressed as mean ± SD. The different treatments were compared with a one-way analysis of variance (ANOVA) and Duncan's multiple comparison tests ($P < 0.05$).

4.3 Results

4.3.1 Crops screening

4.3.1.1 Plant biomass

The aboveground and underground biomass of pak choi and lettuce was significantly decreased by the Se(VI) amendment (Figs. 4.1a and 4.1b). Increasing applications of Se(VI) from 0 to 5.0 mg/kg reduced the dry weight of the shoot and root by 36% and 84% for pak choi, and 45 and 71% for lettuce, respectively. Additionally, for wheat, the exposure to high Se(VI) concentrations at 5.0 mg/kg did not cause any significant growth inhibition (Fig. 4.1c). The low dosage of Se(VI) applications even stimulated the wheat growth: the biomass of wheat shoot and root raised by 21% and 27%, respectively, in the 1.0 mg/kg Se(VI) amendment compared to the control. Similarly, for the maize, the root and stem biomass was increased under the low Se(VI) exposure,

Chapter 4

while significantly declined at the 5.0 mg/kg Se(VI) application compared to the control (Fig. 4.1d). Exposure of beans to Se(VI) significantly stimulated the growth of each bean tissue except there was no significant effect on the growth of the root and stem at the 5.0 mg/kg of Se(VI) exposure in comparison with the control (Fig. 4.1e). Specifically, at 1.0, 3.0 and 5.0 mg/kg of Se(VI) treatments, the yield was improved by 14, 26 and 24% for the leaves, and 25, 31 and 16% for the seeds, respectively, compared to the control treatment.

4.3.1.2 Se content in plants

Se concentrations in all plants considerably raised with increasing dosages to the sandy soil (Fig. 4.2). Pak choi had the highest Se concentration among the five crops investigated, with a content of 732 and 335 mg/kg, respectively, for shoot and root at 5.0 mg/kg of Se(VI) treatment, respectively. The Se content in the shoots of pak choi was 2-4 fold higher than that in the corresponding roots (Fig. 4.2a). The Se content in the shoots of lettuce was 1.1-1.3 times higher than in the roots, but was 1.3-3.0 times lower than in the shoots of pak choi at the same Se amendments (Fig. 4.2b). Maize had an almost similar Se content to lettuce in the root and shoot at the 5.0 mg/kg of Se(VI) amendment, where the maximum Se concentration in the maize was 249 mg/kg in the roots, 136 mg/kg in the stems and 267 mg/kg in the leaves (Fig. 4.2d). The Se concentration was evenly distributed in each bean tissue (root, stem, leaf and seed) and was comparable to that of maize (Fig. 4.2e). Wheat presented the lowest Se concentration among all crops with 166 mg/kg for the shoots and 145 mg/kg for the roots under the highest Se dosage (Fig. 4.2c).




Figure 4.1. Dry weight of the five crops grown at different Na_2SeO_4 applications: (a) pak choi, (b) lettuce, (c) wheat, (d) maize and (e) beans. Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences among treatments with the different Se applications according to Duncan's multiple comparison tests ($P < 0.05$).




Figure 4.2. Se content in plants grown at different Na_2SeO_4 concentrations: (a) pak choi, (b) lettuce, (c) wheat, (d) maize and (e) beans. Values are mean $\pm$ standard deviation ($n=6$). Different letters indicate statistically significant differences among treatments with the different Se applications according to Duncan's multiple comparison tests ($P < 0.05$).

4.3.2 Characterization of the produced Se-enriched biomaterials

Understanding Se speciation in the produced Se-enriched biomaterials could aid to predict and explain the evolution of Se release into soils. The Se species in the Se-

Chapter 4

enriched duckweed and sludge are illustrated in Fig. 4.3a and b, respectively. Se in the Se-enriched duckweed consisted of 80% selenate (Se(VI)), 1.3% selenite (Se(IV)), 0.8% SeCys_2 , 0.7% SeMetSeCys , 0.9% SeMet , and a small amount of an unidentified Se compound. In contrast, Se in the Se-enriched sludge was mainly present as the zerovalent form (94.2%) and the remaining Se species was Se(IV) ..




Figure 4.3. Se speciation in the two Se-enriched biomaterials: (a) chromatogram of Se speciation in the Se-enriched duckweed compared with that of a Se standard solution of 100 $\mu\text{g/L}$ of each species, and (b) XANES spectra of the Se species in the Se-enriched sludge (yellow) as compared to standards.

Chapter 4

4.3.3 Se released to non-planted soils amended with the Se-enriched biomaterials

The Se release from the Se-enriched biomaterials into the soil was quantified by measuring the Se concentration in the soil pore water. Quantification of Se concentration in the soil pore water could also aid to estimate the Se bioavailability in the soils. Both Se-enriched biomaterials were effective for improving the availability of Se in the two types of soil. The pore water of soils amended with Se-enriched duckweed and anaerobic sludge contained a significantly higher amount of Se compared to the blank soil (Fig. 4.4). After 3 days of incubation (first monitoring period), the supplement of DW1 and DW5 increased the Se concentration in the pore water from the soil background values to 537 and 4375 $\mu\text{g/L}$ for sandy soil, and to 413 and 1238 $\mu\text{g/L}$ for loamy soil, respectively; whereas the supplement of SL1 and SL5 led to an increment of the Se content in the pore water to 65 and 322 $\mu\text{g/L}$ for sandy soil, and 72 and 387 $\mu\text{g/L}$ for loamy soil, respectively.

Generally, the Se concentration in the pore water of the two soils supplied with Se-enriched duckweed was significantly higher than that of the two soils supplied with Se-enriched sludge, except for the SL5 and DW5 application in the loamy soil after 14 days incubation of the first monitoring period (Fig. 4.4). Moreover, the Se concentration in the pore water of the sandy soil with the Se-enriched duckweed supplement was 1.1–5.0 times higher than that of the pore water of the loamy soil, while no significant difference was observed between the two soil types with Se-enriched sludge amendment before 42 days of incubation (first monitoring period) (Fig. 4.4).

The difference in the Se release pattern between Se-enriched duckweed and sludge continued to exist over time. The Se concentration in the pore water decreased along with the incubation time with the DW5 supplement in both soil types in the first monitoring period. Specifically, increasing the incubation time from 3 to 42 days reduced the Se content in pore water by 92% for the sandy soil and 89% for the loamy soil. Conversely, the Se concentration in the pore water of both soils supplied with Se-enriched sludge was stable. After one year of incubation (second monitoring period), the Se concentration in the pore water of the two soils supplied with Se-enriched duckweed was slightly lower compared to the 42 days incubation of the first monitoring

Chapter 4

period, while it was around 1.5–2.0 times higher for the Se-enriched sludge supplement in both soil types investigated (Fig. 4.4).




Figure 4.4. Evolution of Se concentration in the Rhizon extracts of non-planted soils amended with Se-enriched duckweed and sludge along with incubation time: (a) sandy soil, and (b) loamy soil. Values are mean $\pm$ standard deviation ($n = 3$), DW: duckweed amendment, SL: sludge amendment, $\circ$ DW5, $\bullet$ DW1, $\triangle$ SL5, $\blacktriangle$ SL1 and $\blacksquare$ Blank.

4.3.4 Se released to planted soils amended with Se-enriched biomaterials

4.3.4.1 Se content in the soil pore water

Similarly, as in the non-planted soils, amendment of soils planted with beans with Se-enriched duckweed or sludge significantly increased the Se concentration in the pore water of both soils compared to the blank (Fig. 4.5 and Table 4.S2). The effect of the Se dosage was also evident from the higher Se concentration in the soil pore water for the most Se-enriched biomaterials (DW5 and SL5). After 7 days of incubation, the

Chapter 4

supplement of DW1 and DW5 increased the Se concentration in the pore water from the soil background values (1.3 and 1.0 µg/L for the sandy and loamy soil, respectively) to 920 and 2330 µg/L for the sandy soil, and 595 and 1851 µg/L for the loamy soil, respectively; whereas the application of SL1 and SL5 led to an increment of the Se content in the pore water to 85 and 200 µg/L for the sandy soil, and 58 and 252 µg/L for the loamy soil, respectively.

The Se concentrations in the soil pore water with DW1 and DW5 amendments decreased by 70% and 80% for the sandy soil, and by 98% and 85% for the loamy soil from day 7 to day 42, respectively. Similarly, for the SL1 amendment, the Se concentration in the pore water decreased by 48% and 32% from day 7 to day 42 for the sandy and loamy soil, respectively. The decrease in Se pore water concentration for the SL1 amendment was less pronounced than the decrease observed for the Se-enriched duckweed amendment. However, the amendment of SL5 significantly increased the Se concentration in the pore water by 143% for the sandy soil after 42 days compared to 7 days, whereas it increased the Se concentration in pore water for the loamy soil in the middle of the planting phase (28 days) and remained stable afterwards (Fig. 4.5 and Table 4.S2).

Additionally, the Se concentration in the pore water of the two soil types supplied with Se-enriched duckweed was significantly higher than that of the soils supplied with Se-enriched sludge before 21 days of incubation, while no significant difference was observed afterwards, except for the application of SL1 and DW1 to the loamy soil (Table 4.S2).

Chapter 4




Figure 4.5. Evolution of Se concentration in the Rhizon extracts of soils amended with Se-enriched duckweed and sludge over the growth period of beans in (a) sandy soil, and (b) loamy soil. Values are mean $\pm$ standard deviation ($n=3$), ○ DW5, ● DW1, △ SL5, ▲ SL1 and ■ Blank.

4.3.4.2 Biomass of bean tissues

The effect of biomaterials application on the growth of beans can be assessed based on biomass production of the beans. Table 4.2 shows the biomass of beans grown in soils amended with Se-enriched biomaterials, non-enriched biomaterials (the corresponding controls) and without any biomaterials (Blank). There was no significant

Chapter 4

difference between the blank and the application of Se-enriched biomaterials except for the DW5 application in the sandy soil, which significantly declined the root biomass of the beans. However, when compared with application of non-enriched biomaterials, the biomass of beans grown in soil amended with the Se-enriched biomaterials is generally higher. The amendment with non-enriched duckweed and sludge (control) significantly decreased the root biomass of beans in the sandy soil, and considerably reduced the seed and total biomass of beans in the loamy soil (Table 4.2). However, a higher biomass production was observed with the supplementation of Se-enriched biomaterials to both soils, indicating that Se released from the Se-enriched biomaterials could promote the growth of beans.

Table 4.2. Dry biomass of the roots, seeds and whole plant (sum of root, stem, leaf and seed) of beans grown in soils with different biomaterials application (Mean $\pm$ standard deviation, $n = 3$). Different lowercase letters indicate statistically significant differences between different Se dosages for the same type of Se-enriched biomaterial application and the same type of soil according to Duncan's multiple comparison tests ($P < 0.05$). Different uppercase letters indicate significant differences between different dosages for the same type of non-enriched biomaterials (control) application and the same type of soil ($P < 0.05$).

	Treatments	Root	Seed	Whole plant
<i>Sandy soil</i>	Blank	0.54 $\pm$ 0.01 b ^A	0.72 $\pm$ 0.53	4.21 $\pm$ 0.26 a ^A
	DW1	0.65 $\pm$ 0.04 a	0.79 $\pm$ 0.18	4.03 $\pm$ 0.32 a
	DW5	0.33 $\pm$ 0.02 c	0.57 $\pm$ 0.42	3.19 $\pm$ 0.38 b
	Control-DW1	0.48 $\pm$ 0.32 ^A	0.57 $\pm$ 0.23	4.14 $\pm$ 1.26 ^A
	Control-DW5	0.12 $\pm$ 0.04 ^B	0.06 $\pm$ 0.03	1.46 $\pm$ 0.05 ^B
	Blank	0.54 $\pm$ 0.01 ^A	0.72 $\pm$ 0.53	4.21 $\pm$ 0.26
	SL1	0.65 $\pm$ 0.13	0.53 $\pm$ 0.10	3.65 $\pm$ 0.60
	SL5	0.50 $\pm$ 0.05	0.40 $\pm$ 0.03	3.85 $\pm$ 0.30
	Control-SL1	0.36 $\pm$ 0.21 ^{AB}	0.69 $\pm$ 0.45	3.52 $\pm$ 1.5
	Control-SL5	0.20 $\pm$ 0.03 ^B	0.51 $\pm$ 0.31	2.80 $\pm$ 0.45
<i>Loamy soil</i>	Blank	0.48 $\pm$ 0.03 ^A	0.51 $\pm$ 0.24 ^A	3.70 $\pm$ 0.11 ^A
	DW1	0.37 $\pm$ 0.08	0.86 $\pm$ 0.39	3.79 $\pm$ 0.79
	DW5	0.49 $\pm$ 0.11	0.71 $\pm$ 0.41	4.12 $\pm$ 0.68
	Control-DW1	0.23 $\pm$ 0.09 ^B	0.25 $\pm$ 0.05 ^{AB}	2.72 $\pm$ 0.87 ^B
	Control-DW5	0.22 $\pm$ 0.00 ^B	0.12 $\pm$ 0.08 ^B	2.21 $\pm$ 0.11 ^B
	Blank	0.48 $\pm$ 0.03	0.51 $\pm$ 0.24 ^A	3.70 $\pm$ 0.11 ^A
	SL1	0.65 $\pm$ 0.13	0.55 $\pm$ 0.11	3.67 $\pm$ 0.37
	SL5	0.50 $\pm$ 0.05	0.41 $\pm$ 0.21	3.86 $\pm$ 0.37
	Control-SL1	0.30 $\pm$ 0.11	0.36 $\pm$ 0.12 ^B	2.87 $\pm$ 0.74 ^{AB}
	Control-SL5	0.23 $\pm$ 0.00	0.25 $\pm$ 0.02 ^B	2.61 $\pm$ 0.26 ^B

Chapter 4

4.3.4.3 Se content in the different bean tissues

Selenium concentration in the bean plants may directly reflect the bioavailability of Se released in the soil from the Se-enriched biomaterials and evaluate their potential for use as Se biofertilizers. The amendment of Se-enriched duckweed and sludge significantly increased the Se concentration in the different tissues of beans (seed, leaf, stem and root) growing on both the sandy and loamy soil (Fig. 4.6). Increasing the amount of Se-enriched duckweed and sludge amendment from 1.0 to 5.0 mg Se /kg soil raised the Se content in the roots of beans from 17 to 52 mg/kg and from 27 to 117 mg/kg for the sandy soil, respectively, and increased the Se content in the roots of beans from 14 to 38 mg/kg and from 26 to 118 mg/kg for the loamy soil, respectively. The Se concentration in the beans amended with Se-enriched sludge was mostly 1.1–3.1 times higher than in those amended with Se-enriched duckweed. Besides, the highest Se content was observed in the roots among all tissues for both Se-enriched duckweed and Se-enriched sludge amendment, being 2–12 fold higher than those in the stem, leaf and seed.

4.3.4.4 Correlation between Se in the beans and Se in the soil pore water

The correlation between Se in the plants and Se in the soil pore water could help to identify the source of Se taken up by the plants and to determine whether the concentration of Se released in the soil pore water can be adopted to predict Se bioavailability. Given this, Fig. 4.7 shows the correlation between the Se concentration in the seeds and roots of beans and the Se concentration in the pore water extracted from the sandy and loamy soils amended with Se-enriched duckweed and sludge after 7 days incubation (the first day of transplanting beans). The Se content in the soil pore water positively and significantly correlated with the Se concentration in both the seeds and roots. The high correlation coefficient (between 0.80 and 0.99) between the Se concentration in the tissues of beans and in the soil pore water indicates that the Rhizon extraction can properly assess the bioavailability of Se in soils.




Figure 4.6. Se content in the different tissues of harvested beans (42 days growth) fertilized with Se-enriched duckweed and sludge in (a) sandy soil, and (b) loamy soil. Values are mean $\pm$ standard deviation (n=3).




Figure 4.7. Correlation between the concentration of Se in the soil pore water (7 days incubation) and seeds (○) and roots (●) of beans: (a) sandy soil with Se-enriched duckweed supplement, (b) sandy soil with Se-enriched sludge supplement, (c) loamy soil with Se-enriched duckweed supplement, and (d) loamy soil with Se-enriched sludge supplement.

4.3.4.5 Selenium speciation in the seeds

Selenium speciation in the Se-enriched beans is an important factor to evaluate when exploring the potential of these biofortified beans as beneficial foods for a Se deficient population. The seeds of beans grown in the sandy soil with an amendment of 1.0 mg Se/kg Se-enriched duckweed (DW1) and sludge (SL1) and the blank soil were selected for the determination of Se speciation (Fig. 4.8 and Table 4.3). Organic Se (SeCys2, SeMetCys and SeMet) was the main Se species in all seeds samples. SeMet accounted for the highest percentage of the total Se in the seeds. The percentage of SeMet in the seeds with the amendment of DW1 (72%) was notably lower than that of the amendment of SL1 (85%). Besides, in the blank and DW1 amendment, Se(VI) was

Chapter 4

the dominant inorganic Se species (accounted for 13.7% and 12.2% of total Se, respectively). Compared to the Se-enriched duckweed amendment (13%), the percentage of inorganic Se species of total Se in the bean seeds was much lower for the Se-enriched sludge amendment (1.7%) (Table 3). Apart from the identified Se species, unidentified Se peaks in the beans were observed both for the Se-enriched duckweed and sludge application (Fig. 4.8).

Table 4.3. Percentage of Se species in the seeds of beans grown in sandy soil amended with 1.0 mg Se/kg soil of Se-enriched duckweed (DW1) and sludge (SL1), and in the blank without amendment (Blank) comparison with total Se in bean seeds.

	Total Se	SeCys ₂	SeMetCys	SeMet	Se(IV)	Se(VI)	Se species recovery
	(mg/kg)				(%)		
Blank	0.12	3.2	3.5	77.2	2.4	13.7	99
DW1	4.63	1.8	5.2	71.7	0.4	12.2	91
SL1	6.95	2.7	11.6	84.6	0.3	1.4	101

Chapter 4




Figure 4.8. Se speciation in the harvested seeds of beans after 42 days grown in the sandy soil amended with 1.0 mg Se/kg of Se-enriched duckweed (DW1) and sludge (SL1): (a) Se standard solution of 10 $\mu\text{g/L}$, (b) Blank soil (higher magnification in insert), (c) DW1 amendment and (d) SL1 amendment.

Chapter 4

4.3.5 pH, TC and TOC contents in the soil pore water

The influence of Se-enriched biomaterials application on soil characteristics was assessed via pH, TOC and TC measurement. These parameters help to explain differences in Se bioavailability in soils amended with different Se-enriched biomaterials. The pH, and the TC and TOC concentration significantly increased upon the amendment of Se-enriched biomaterials (Table 4.4). The highest TC content was found for the DW5 amendment for both the sandy and loamy soils, which was 7.2 and 3.2 times higher than that of the blank in the non-planted soil, and 4.8 and 2.9 times higher than that of the blank in the planted soil, respectively. However, no significant difference was observed between the amendment of Se-enriched sludge and the blank in the planted loamy soil. Moreover, the TC and TOC content in the Se-enriched duckweed amendments were 1.2–2.7 times higher than those of the Se-enriched sludge amendments for both planted and non-planted sandy and loamy soils. The values of pH, TC and TOC in the planted soils were slightly lower, in comparison with the non-planted soils.

Table 4.4. Characterization of Rhizon extracts after 42 days of Se-enriched biomaterials application in planted and non-planted soils.

	Treatments	Sandy soil			Loamy soil		
		pH	TC (mg/L)	TOC (mg/L)	pH	TC (mg/L)	TOC (mg/L)
Non-Planted	Blank	6.0 ± 0.2	33 ± 5.2	30 ± 5.9	6.5 ± 0.1	72 ± 5.0	70 ± 2.3
	DW1	6.8 ± 0.0	63 ± 8.4	57 ± 5.5	6.3 ± 0.3	110 ± 13.2	108 ± 15.8
	DW5	7.6 ± 0.2	236 ± 22.1	201 ± 18.2	7.4 ± 0.1	233 ± 113.8	212 ± 102.0
	SL1	7.8 ± 0.1	48 ± 4.3	47 ± 4.6	7.7 ± 0.1	68 ± 2.2	63 ± 10.3
	SL5	7.8 ± 0.0	136 ± 25.7	133 ± 24.8	7.8 ± 0.0	128 ± 39.2	111 ± 38.5
Planted	Blank	4.9 ± 0.5	36 ± 8.4	34 ± 5.6	6.2 ± 0.2	67 ± 16.5	67 ± 15.6
	DW1	5.4 ± 0.2	76 ± 4.4	75 ± 4.3	6.6 ± 0.3	77 ± 27.3	76 ± 24.9
	DW5	6.7 ± 0.2	172 ± 17.0	166 ± 17.2	7.1 ± 0.2	193 ± 65.4	181 ± 62.3
	SL1	7.0 ± 0.1	43 ± 10.3	41 ± 10.4	6.9 ± 0.2	52 ± 13.7	52 ± 12.4
	SL5	7.5 ± 0.1	79 ± 14.6	77 ± 14.8	7.6 ± 0.2	65 ± 15.2	58 ± 16.6

Chapter 4

4.3.6 Evaluation of the Se-enriched biomaterials produced from wastewater as slow-release fertilizers for Se biofortification

The Se utilization efficiency by beans amended with Se-enriched duckweed and sludge is calculated and presented in Table 4.S3. Results showed that approximately 5% of the applied Se in DW1 and SL1 was accumulated in bean plants. Similarly, around 2% and 4% of the applied Se in DW5 and SL5 was accumulated in the beans, respectively.

The Se concentrations ranged from 4-9 mg/kg and 3-23 mg/kg in the seeds of bean upon supplementation of the Se-enriched duckweed and sludge, respectively. If we assume that a daily serving size of the Se-enriched bean seeds is 7.2 g dry weight (96 g fresh correspondingly), consumption of the bean seeds biofortified with Se-enriched duckweed and sludge amendment would result in the ingestion of 28-64.8 µg Se and 21.6-166 µg Se/day, respectively. These values are close to the recommended daily Se requirements (70 µg Se/day in Belgium).

All EDI values for the amendment of Se-enriched duckweed and sludge were also lower than the oral reference dose of Se (5.0 µg/kg·day), while the value for the amendment of chemical Na₂SeO₄ at a dose of 5.0 mg/kg (unpublished data from a previous experiment) was higher than 5.0 µg/kg·day (Table 4.5). Similarly, the estimated HRI in all treatments was below 1, except in the treatment of 5.0 mg/kg of Na₂SeO₄ (1.860). The EDI and HRI values of children were slightly higher than those of adults. These estimations suggest that normal consumption of Se-enriched beans produced through soil amendment of Se-enriched duckweed and sludge would not pose a potential risk of excessive Se intake, whereas beans grown on soil amended with 5.0 mg/kg Na₂SeO₄ might cause a significant health hazard for both adults and children.

Chapter 4

Table 4.5. Estimated daily Se intake ($\mu\text{g/kg/day}$) through seeds of beans grown in sandy and loamy soil fertilized with Se-enriched biomaterials.

Treatments	EDI				HRI			
	Adult		Children		Adult		Children	
	Sandy	Loamy	Sandy	Loamy	Sandy	Loamy	Sandy	Loamy
Blank	0.01	0.01	0.03	0.01	0.003	0.001	0.005	0.002
DW1	0.19	0.21	0.35	0.38	0.038	0.041	0.070	0.076
DW5	0.40	0.36	0.74	0.65	0.080	0.071	0.148	0.130
SL1	0.29	0.15	0.53	0.27	0.057	0.030	0.105	0.054
SL5	0.94	0.79	1.73	1.45	0.188	0.158	0.346	0.291
Na ₂ SeO ₄ 1.0	1.23	-	2.26	0.00	0.246	0.000	0.452	-
Na ₂ SeO ₄ 5.0	5.06	-	9.30	0.00	1.013	0.000	1.860	-

Note: Na₂SeO₄ at a dose of 1.0 and 5.0 mg/kg concerned unpublished data from a previous experiment.

- : no data obtained in the corresponding treatment

4.4 Discussion

4.4.1 Se toxicity and accumulation in different crops

This study showed that pak choi had the highest ability of Se uptake among pak choi, lettuce, maize, wheat and bean, reflecting on the largest Se content in the shoot and root (1.3–3.0 times higher than that in the lettuce), while wheat represented the lowest Se uptake ability (Fig. 4.2). Pak choi is a vegetable species of *Brassicaceae* commonly found in China, and has been shown as Se accumulator (Li et al., 2015a). *Brassicaceae* species such as *Stanleya pinnata* and *Brassica juncea* (indian mustard) can indeed accumulate and tolerate a high Se content and may potentially be a Se hyperaccumulator applied in seleniferous areas (White, 2016; Yawata et al., 2010). In contrast, cereals such as wheat, generally have a lower Se uptake ability, in comparison with legumes, some brassicas and fruits (Díaz-Alarcón et al., 1996; Wu et al., 2015), which corroborates to the findings of this study (Fig. 4.2).

Chapter 4

The biomass yield obtained in this investigation showed that vegetables (pak choi and lettuce) have a lower tolerance to Se exposure among the five crops tested (Figs. 1a and b). The greatest reductions of root biomass under Se exposure among all crops were observed in pak choi and lettuce. In contrast, the application of Se stimulated the growth of wheat and bean in this study, even at the 5.0 mg/kg of Se application for the seeds and leaves of beans. The largest root biomass of these two crops was at the amendment 1.0 mg/kg Se, indicating that 1.0 mg/kg of Se is the optimal Se application for the two plants grown (Fig. 4.1) and these two crops have a higher tolerance for Se. In agreement with the findings in this research, other researchers have also demonstrated the dual effect of Se on crops (Guerrero et al., 2014; Wang et al., 2017). The optimal dose of Se stimulates crop growth, while excessive Se inhibits plant growth (Figs 4.1c-d).

Selenium taken up by plants will partly replace sulfur in amino acids and selenoaminoacids such as SeMet will subsequently be incorporated into proteins, which is beneficial for human and animal nutrition (Eiche et al., 2015; Schrauzer, 2000). Given that beans have the highest protein content in seeds as well as high Se tolerance ability among the five crops in this study, it was selected for the following experiment of Se-enriched biomaterials fertilization.

4.4.2 Selenium release from Se-enriched duckweed and sludge

The results showed that the amendment of soils with Se-enriched duckweed and sludge increased the availability of soluble Se in the soils, as evidenced by the increment of the Se concentration in the pore water with larger doses of enriched biomaterials (Fig. 4.4). Similarly, Bañuelos et al. (2015) found that the increase of available soluble Se positively correlated with the amount of Se-enriched *Stanleya pinnata* applied. Dhillon et al. (2007) indicated that the application of Se-enriched wheat (*Triticum aestivum* L.) and raya (*Brassica juncea* L.) straw from 0% to 1% (ratio of straw weight to soil weight) increased the hot water-soluble Se (bioavailable Se) fraction in the soil from 18 µg/kg to 36 and 79 µg/kg, respectively.

However, the available Se in soils amended with Se-enriched duckweed decreased by 90% over an incubation period of 42 days even in the absence of planted crops (Fig. 4.4). This may be partly attributed to the immobilization of the released Se in the soil

Chapter 4

solid phase (both inorganic and organic) and the adsorption of Se onto the extra organic matter introduced with the supplementation of the Se-enriched biomaterials (Li et al., 2017). Specifically, the organic content of the applied duckweed and sludge introduces organic compounds, including some organic acids, polysaccharides and lignin, which augment the soil organic matter (Wang et al., 2018). These organic compounds can bind soluble Se into stable compounds, resulting in Se immobilization (Ebrahimi et al., 2019; Li et al., 2017). The increase in organic compounds was confirmed by the significant increment of the TC and TOC concentration in the Rhizon extracts of both soils after the biomaterials application (Table 4.4). Research on Se species transformation in straw-amended soil further demonstrated that straw-derived organic carbon could accelerate the conversion of soluble Se (SOL-Se) to organic matter-bound Se (OM-Se) and residual Se (RES-Se), resulting in less soluble Se in soils (Arbestain, 2001). Wang et al. (2018) also found that most SOL-Se was reduced and transformed to RES-Se in a soil fertilized with Se-enriched wheat straw. Similarly, in this study, the decrease of Se in the pore water along the incubation time (Figs. 4.4 and 4.5) may be partly explained by the transformation of SOL-Se to other, unavailable Se fractions in soils, e.g. OM-Se and RES-Se.

On the other hand, the higher quantity of Se in the pore water of the Se-enriched duckweed amended soil (14% of the total applied Se after 3 days of DW1 amendment) compared to that of the Se-enriched sludge amended soil (2% of the total applied Se after 3 days of SL1 amendment) (Fig. 4.4) may be mainly ascribed to different Se species in the Se-enriched biomaterials. Specifically, inorganic Se(VI) represented a large proportion of the Se in duckweed (ca. 80%, Fig. 4.3a), whereas elemental Se(0) dominated in the sludge (94.2%, Fig. 4.3b), especially nano-Se(0) (Staicu et al., 2015b). Se(VI) is highly soluble and readily mobilized in soils and waters (Li et al., 2020a). In contrast, elemental Se(0) is stable and difficult to mobilize or oxidize in the environment (Hu et al., 2018), resulting in less and slower Se released to the soil pore water from the Se-enriched sludge amendment.

4.4.3 The potential of Se-enriched duckweed and sludge as Se biofertilizers

The increase in Se uptake by beans positively correlated with the application rates of Se-enriched duckweed and sludge (Fig. 4.7), implying that Se in the soil pore water released from the Se-enriched duckweed and sludge is the major source of Se in the

Chapter 4

tissues of beans. Interestingly, the higher Se concentration in the soil pore water with the Se-enriched duckweed amendment compared to that with the Se-enriched sludge supplement did not result in a higher Se concentration in the corresponding plant biomass (Figs. 4.5 and 4.6). This is probably due to the different Se species released to the soil solution from the two Se-enriched biomaterials (Fig. 4.1) and the different behavior of Se after being released into soils (Figs. 4.4 and 4.5), resulting in a different metabolism of Se in beans and consequently leading to a different Se uptake/accumulation ability. Specifically, the high amount of Se(VI) present in the duckweed (80%, Fig. 4.3a) is immediately released into the soil solution where it is quickly adsorbed by the soil matrix or extra organic matter (Li et al., 2016), thus leading to a significant decrease of the plant bioavailable Se concentration in the soil pore water along time, even lower than that in the sludge amendment after 28 days of growth (Fig. 4.5 and Table 4.S2). In contrast, the zerovalent form of Se in the Se-enriched sludge (Fig. 4.3b) may have been slowly released or oxidized to Se(IV) or Se(VI) in the soil solution. The slowly released Se in the soil pore water could continuously supply and satisfy the Se uptake requirement of the beans during the growth period, contributing to the higher Se uptake (Fig. 4.6). Furthermore, the Se(0) in the Se-enriched sludge may mostly exist as biogenic Se(0) nanoparticles with an average size of about 166 nm (Staicu et al., 2015b), which has a similar behavior to Se(IV) after being taken up by plants (El-Ramady et al., 2020; Moreno-Martin et al., 2020). This could also explain the higher Se concentration in bean tissues grown in Se-enriched sludge amendment soil. Besides, the speciation of Se accumulated in the seeds of beans fertilized with Se-enriched biomaterials further supports the different uptake patterns and different Se species released from Se-enriched duckweed and sludge (Fig. 4.8 and Table 4.3). Specifically, this study found that a higher fraction of inorganic Se(VI) (13.3%) accumulated in the seeds of beans amended with Se-enriched duckweed, compared to only 1.6% of Se(VI) in the seeds fertilized with Se-enriched sludge (Table 3), which indirectly confirms that the form of Se released from Se-enriched duckweed was mainly Se(VI). Because Se(VI) taken up by plants is reduced into Se(IV) and then converted to selenaminoacids through a reduction process via substitution for sulfate in the ATP sulfurylase reductase system (Gupta & Gupta, 2017). This reduction process is an ATP-consuming process and rate-limiting step, resulting in the accumulation of Se(VI) in the beans with the Se-enriched duckweed amendment (Table 4.3). In contrast, the Se-enriched sludge may release Se(0) into soils and then

Chapter 4

be oxidized to Se(IV) by microorganisms (Sarathchandra & Watkinson, 1981; Winkel et al., 2015). It is also possible that the nano-Se(0) released in the soil solution was directly taken up and is metabolized in a similar way as Se (IV) in plants (El-Ramady et al., 2020; Moreno-Martin et al., 2020). Se(IV) is quickly and easily converted to organic Se forms (e.g., SeMet and SeCys₂) in the beans after being taken up by the roots and translocated to the seeds.

The significantly higher amount of TOC in the soil pore water caused by the Se-enriched duckweed supplement as compared with the Se-enriched sludge amendment (Table 4.4) also partially contributed to the lower Se uptake by the beans (Fig. 4.6), as organic carbon in the soil pore water may bind with Se (Supriatin et al., 2016; Weng et al., 2011), and eventually lower the Se bioavailability (Li et al., 2017; Wang et al., 2018). Besides, soil pH also plays a role in the mineralization of soluble organic Se. Supriatin et al. (2016) demonstrated that soil pH is the primary factor to control the solubility of Se in dissolved organic carbon and confirmed that the solubility of Se-containing organic matter increases with pH. In this study, the higher pH of the pore water of the soils with Se-enriched sludge compared to the Se-enriched duckweed supplement (Table 4) may have resulted in a higher solubility of Se, hence more Se available for plant uptake and thus leading to a higher Se content in the beans (Fig. 4.6). These results indicated that Se-enriched sludge is considered as the preferred slow-release Se biofertilizer and an effective Se source to produce Se-enriched crops for Se-deficient populations.

The application of Se-enriched biomaterials did not show negative effects on the biomass production of beans, while the higher amount of non-enriched biomaterials application may have caused an abiotic stress, resulting in a decreased biomass of beans (Table 4.2). Similarly, this has been demonstrated in other studies where amendment of high amounts of compost, weeds or crops residues (such as the residues of *Chenopodium murale* and *Parthenium hysterophorus*) were associated with the release of a wide range of inhibitors (e.g. tannins and phenols compounds) to the environment that may be toxic or cause stress to plants (Batish et al., 2007; Mushtaq et al., 2020; Singh et al., 2003), which is broadly associated with allelopathic interactions. Therefore, the presence of inhibitory compounds in the non-enriched biomaterials amended soils in this study may partially explain the significant reduction in the bean biomass upon non-enriched biomaterials supplementation. On the other

Chapter 4

hand, no significant reduction of bean biomass was observed when Se-enriched biomaterials were used as biofertilizers. This indicates that the Se released from the biomaterials could protect the plants against various types of external stresses (Handa et al., 2016). Various studies have revealed the direct effect of Se on the antioxidative defense system, which increases the potential of the plants to combat stressful conditions (Handa et al., 2016; Jozwiak & Politycka, 2019; Subramanyam et al., 2019). Besides, P has been proposed to negatively affect Se uptake by plants in many studies. In this study, the large difference of total P in sandy and loamy soil could partially contribute to the slightly lower Se in loamy soil pore water and in beans grown in loamy soil.

Furthermore, the high amount of SeMet (around 80%, Table 4.3) present in the seeds of the beans suggests that these Se-enriched seeds obtained by the amendment of Se-enriched biomaterials may be of interest for animals or human nutrition in Se-deficient regions. SeMet is a selenoamino acid that is highly suitable for nutritional supplementation because it is more bioavailable, less toxic, and can provide higher Se concentrations in organs than inorganic Se (Gómez-Jacinto et al., 2020). SeMet is also one of the precursors of methyl selenol, a potent anticarcinogen that inhibits tumor invasion and angiogenesis (Gómez-Jacinto et al., 2020). Additionally, the HRI and EDI data suggest that Se-enriched beans produced from this study would not pose a potential risk of excessive Se intake for adults and children (Table 4.5). In contrast, the beans produced from the treatment with 5.0 mg/kg Na_2SeO_4 could cause a significant health hazard due to Se over-consumption. The result thus demonstrated that the Se recovered from wastewater through ecotechnologies (phytoremediation and bioreduction) can be reused as biofertilizers to efficiently improve Se level in feeds/foods. Biofertilizers from wastewater can thus be valorized, while recovering resources.

4.4.4 Study limitations and future perspectives

Even if the biofortification of beans through soil amendment with Se-enriched biomaterials produced from wastewater has been achieved in this study, additional studies should particularly address the safety and other risks associated with the application of these Se-enriched biomaterials on a short and long term, such as their potential heavy metal content and pathogen load. Besides, regulations on use of

Chapter 4

biomaterials from wastewater should be complied as well. Furthermore, an economic analysis of Se recovery from wastewater and reusing Se as biofertilizer is also still necessary.

Besides, the results of bean biomass analysis showed that some potential side effects of the application of duckweed and sludge into soils may exist. For instance, a high amount of duckweed or sludge applied into soils may disturb the C/N ratio of the soils and thus affect soil microbial activities, as also reported in other studies where rice and wheat straw were amended into soils (Jin et al., 2020). Additionally, the relatively lower Se bioavailability and the significant decrease of Se content in the soils amended with Se-enriched duckweed indicate that post-treatment of the duckweed harvested from the wastewater should be further considered before application as biofertilizer, such as through composting and extracting. Continuous cultivation under field conditions should be conducted to investigate the evolution of release and bioavailability of Se on longer term and its long-lasting effects on the environment and nutritional value of the crops.

In terms of estimation of daily Se intake through seeds of beans produced in this study, total Se in the raw seed of beans was simply adopted for assessment. This could result in the overestimation of real Se intake, as beans are generally consumed after cooking, e.g. boiling and frying. Boiling or frying may cause Se losses through the cooking water or volatilization. This has been demonstrated for other crops, such as during cooking of Se-biofortified potato and mushroom (Dong et al., 2021; Zhou et al., 2021).

4.5 Conclusions

Soil amendment with Se-enriched biomaterials produced from wastewater was shown to be a promising approach for Se biofortification, which was able to provide sufficient bioavailable Se for crop uptake and subsequent dietary supplementation of Se deficient populations. Both the Se content in the tissues of beans and soil pore water were significantly enhanced by the Se-enriched duckweed and sludge amendment. The Se in the Se-enriched duckweed, mainly contained in proteins, was released quicker to the soil than the Se contained in the Se-enriched sludge, mainly present as Se(0). However, the Se-enriched sludge (nano-Se(0)) was more efficient than the Se-enriched duckweed in increasing the Se content in beans, in particular the SeMet

Chapter 4

concentration of the seeds. The value-added Se-enriched sludge may thus be considered as the preferred slow-release Se biofertilizer for reusing the recovered Se. Nevertheless, further studies are still needed to validate these results under real field conditions with different soil types and growth environments.

Supplementary Information




Figure 4.S1. Selenium content in the wastewater/medium at the beginning (Before) and end (After) of the experiment and Se removal efficiency by sludge.



Figure 4.S2. Schematic representation of the experiment

Chapter 4

Table 4.S1. Elemental composition of the Se-enriched duckweed and sludge

	Se-enriched duckweed	Se-enriched sludge
	mg/kg	
Cu	31 ± 1	36 ± 4
Zn	370 ± 3	117 ± 3
Fe	1658 ± 37	3837 ± 146
Mn	4751 ± 930	2312 ± 23
Al	188 ± 7	1711 ± 25
Ni	5 ± 3	44 ± 1
Cr	<LOQ	96 ± 1.8
Pb	<LOQ	<LOQ
Cd	<LOQ	<LOQ
Co	<LOQ	<LOQ

< LOQ: values lower than the limit of quantification

Chapter 4

Table 4.S2. Selenium concentrations ($\mu\text{g/L}$) in the pore water extracted from planted and non-planted soils amended with Se-enriched duckweed and sludge. Mean $\pm$ standard deviation, $n=3$. Different lowercase letters indicate statistically significant differences between different incubation days according to Duncan's multiple comparison tests ($P < 0.05$). * indicates significant differences between different Se-enriched biomaterials.

			Day 7	Day 21	Day 28	Day 42
Planted	Sandy soil	Blank	1.8 ± 0.4 a	1.5 ± 0.0 a	1.3 ± 0.3 a	1.2 ± 0.1 a
		DW1	920 ± 85.4 a*	314 ± 120.2 b*	264 ± 120.8 b*	278 ± 231.1 b
		SL1	85 ± 9.0 a*	45 ± 6.0 b*	43 ± 5.0 b*	44 ± 15.2 b
		DW5	2330 ± 646.1 a*	549 ± 466.0 b	319 ± 113.2 b	305 ± 143.2 b
		SL5	200 ± 44.0 c*	255 ± 8.3 bc	312 ± 62.1 b	485 ± 108.7 a
	Loamy soil	Blank	1.0 ± 0.2 a	0.7 ± 0.4 a	1.5 ± 0.8 a	1.2 ± 0.4 a
		DW1	595 ± 44.2 a*	107 ± 12.3 b*	42 ± 3.7 c	7.9 ± 1.3 d*
		SL1	58 ± 0.5 a*	50 ± 3.0 a*	48 ± 18.0 a	39 ± 13.3 a*
		DW5	1851 ± 391.1 a*	823 ± 345.4 b*	341 ± 160.3 b	272 ± 220.2 b
		SL5	252 ± 7.0 c*	313 ± 7.0 ab*	355 ± 44.8 a	292 ± 53.9 bc
Non-planted	Sandy soil	Blank	1.4 ± 0.2 a	1.5 ± 0.3 a	1.1 ± 0.0 a	0.9 ± 0.2 a
		DW1	819 ± 96 a*	667 ± 15 b*	510 ± 28 c*	401 ± 33 c*
		SL1	69 ± 7.0 a*	39 ± 5.2 b*	34 ± 3.5 bc*	24 ± 0.1 c*
		DW5	1996 ± 187 a*	1404 ± 104 b*	605 ± 229 b	332 ± 230 b
		SL5	196 ± 24 a*	251 ± 46 a*	260 ± 53 a	196 ± 18 a
	Loamy soil	Blank	0.9 ± 0.2 a	0.6 ± 0.3 a	1.0 ± 0.2 a	0.9 ± 0.3 a
		DW1	586 ± 6.0 a*	432 ± 0.0 a*	436 ± 112 a*	459 ± 105 a*
		SL1	52 ± 3.3 a*	50 ± 9.2 a*	51 ± 0.9 a*	42 ± 3.9 a*
		DW5	1242 ± 132 a*	2406 ± 215 b*	2082 ± 441 b*	996 ± 406 b
		SL5	322 ± 35 a*	452 ± 23 a*	468 ± 95 a*	432 ± 198 a

Chapter 4

Table 4.S3. Selenium utilization efficiency by beans upon amendment of soil with Se-enriched duckweed and sludge

		Amount of Se applied in each pot (µg/pot)	Se accumulation in plants of each pot (µg/pot)	Se utilization rate (%) *
Sandy soil	DW1	500	29	5.8%
	SL1	500	29	5.8%
	DW5	2500	42	1.7%
	SL5	2500	112	4.5%
Loamy soil	DW1	500	23	4.7%
	SL1	500	25	5.1%
	DW5	2500	53	2.1%
	SL5	2500	109	4.4%

*Se utilization efficiency= Se accumulation in plants of each pot (g)/ amount of Se applied in each pot (g)

Chapter 5

Chapter 5 Biofortification of green beans with Se and Zn-enriched duckweed and sludge

Chapter 5

Abstract

The potential of biomaterials (duckweed and sludge) enriched with Se and Zn as micronutrient biofertilizers for simultaneously improving the Se and Zn content in green beans (*Phaseolus vulgaris*) was evaluated in pot experiments. Both the Se and Zn concentrations in the soil pore water increased upon amending both biomaterials. The concentration of Se released from SeZn-enriched duckweed rapidly decreased in the first 21 days and slowly declined afterwards, while it remained stable during the entire growth period upon application of SeZn-enriched sludge. The Zn content in the soil pore water gradually increased over time. In addition, the application of the SeZn-enriched biomaterials significantly increased the Se concentrations (in particular organic Se-methionine) in plant tissues including the seeds, without a negative impact on plant growth, except for a remarkable decrease in biomass production upon the high amount of SeZn-enriched duckweed application (SeZnDW5). This indicates that the SeZn-enriched biomaterials could be used as organic Se biofertilizers for Se-deficient soils, but an appropriate dose should be determined. In contrast, as the Zn content of the beans was not noticeably improved by supplementation of the SeZn-enriched biomaterials, increasing the Zn concentration in the biomaterials may need to be considered.

Keywords: Selenium and Zn bioavailability, Se and Zn-enriched biomaterials, biofortification, biofertilizer, resource recovery

Chapter 5

5.1 Introduction

Selenium and Zn are both essential micronutrients for humans and animals, playing an irreplaceable role in the functioning of enzymes (Hatfield et al., 2014). Multiple micronutrient (e.g. Se and Zn) deficiencies have been found worldwide, particularly in developing countries (Ngigi, 2019; Sazawal et al., 2018), which is mainly associated with the low dietary micronutrient intake in diets or the low diversity of foods (Mao et al., 2014). Improving the micronutrient content in plants, crops and foods is a possible solution for micronutrient deficiency. As plants and crops take up and accumulate micronutrients from the soil where they grow, multi-mineral agronomic biofortification of crops is thus being explored as a simple and effective way to alleviate micronutrient deficiency (Mao et al., 2014; Poblaciones & Rengel, 2017). On the other hand, discharged wastewaters may simultaneously contain excessive Se and Zn due to insufficient treatment (Lim & Goh, 2005). In this context, Se and Zn loaded in those wastewaters may serve as potential nutrient sources from which the nutrients may be recovered and valorized to produce slow-release micronutrient organic fertilizers. Accordingly, the main objective of this study was to explore the possibility of the two biomaterials (duckweed and sludge) generated from Se and Zn-containing water as potential micronutrient (Se and Zn) biofertilizers. The specific objectives include: (1) to monitor the evolution of the released micronutrients (Se and Zn) from the SeZn-enriched biomaterials in the soil pore water, (2) to evaluate the influence of the supplementation of the two biomaterials on the growth of green beans (*Phaseolus vulgaris*) on sandy and loamy soil and its micronutrient accumulation.

5.2 Materials and methods

5.2.1 Soil collection and characterization

Two types of soil (similar to those used in Chapter 4) were collected and classified as sandy and loamy soil. The soil collection and characterization were detailed in the materials and methods section of Chapter 4.

5.2.2 Valorization of SeZn-enriched biomaterials as Se and Zn biofertilizers

5.2.2.1 Preparation of SeZn-enriched biomaterials (duckweed and sludge)

Chapter 5

SeZn-enriched duckweed was produced as described in Chapter 4. Briefly, duckweed collected from a natural freshwater canal in Delft was cultivated in a Hoagland solution together with 5.0 mg/L of Se and 5.0 mg/L of Zn added as Na_2SeO_4 and ZnCl_2 , respectively. The cultivated duckweed was harvested after 7 days of growth, oven-dried and ground for use in the subsequent fertilization experiment. The SeZn-enriched sludge was generated as described in Chapter 4, with minor modification. The sludge collected from a full-scale upflow anaerobic sludge blanket reactor treating pulp and paper wastewater in Eerbeek (The Netherlands) was added to a nutrient medium with 5.0 mg/L of Se and 5.0 mg/L of Zn in the form of Na_2SeO_4 and ZnCl_2 under anaerobic conditions. After 14 days of incubation at 30 °C, the sludge enriched with Se and Zn was separated from the supernatant, oven-dried and ground for further use. The obtained SeZn-enriched duckweed and sludge contained 103 mg Se/kg and 2289 mg Zn/kg dry weight, and 287 mg Se/kg and 563 mg Zn/kg dry weight, respectively.

5.2.2.2 Selenium and Zn bioavailability for planting green beans

An amount equivalent to 1.0 and 5.0 mg Se/kg soil of SeZn-enriched duckweed (4.9 and 24 g) and sludge (1.7 and 8.7 g) was applied to 0.5 kg of the soil. Accordingly, the amendments in this study were 1.0 mg Se/kg soil and 5.0 mg Se/kg soil of SeZn-enriched duckweed (noted as SeZnDW1 and SeZnDW5), and 1.0 mg Se/kg soil and 5.0 mg Se/kg soil of SeZn-enriched sludge (noted as SeZnSL1 and SeZnSL5). The soils without any biomaterials addition served as blank (noted as Blank). For the control experiment, the same amount of non-enriched duckweed and sludge as in the SeZn-enriched biomaterials amendments was applied to the two soil types (noted as control-SeZnDW1, control-SeZnDW5, control-SeZnSL1 and control-SeZnSL5). The non-enriched duckweed and sludge contained 1.9 mg Se/kg and 295 mg Zn/kg dry weight, and 0.5 mg Se/kg and 117 mg Zn/kg dry weight, respectively. The biomaterials were mixed thoroughly with the soils and placed into 10 x 10 cm plastic pots before incubation.

Subsequently, 100 mL of chemical fertilizer solution including 600 mg/L of N, P_2O_5 and K_2O were added to each pot. Pots were incubated indoors at 24°C, 53% relative humidity and 100 $\mu\text{mol}/\text{m}^2/\text{s}$ light intensity by keeping 80% of the water holding capacity for one week. Afterwards, 4 seedlings of green beans (*Phaseolus vulgaris*) previously cultivated in trays with wet vermiculite for one week were transferred into each pot. All

Chapter 5

experiments were conducted in triplicate. The soil pore water was collected every week through extraction using Rhizon soil moisture samplers for total Se and Zn measurement. pH, TC and TOC of the extracted soil pore water were determined before harvest. The bean plants were harvested after 6 weeks of growth, washed and separated into different tissues (root, stem, leaf and seed) for biomass, Se and Zn concentration analysis. Se speciation analysis was performed on selected bean seeds (at the SeZnDW1 and SeZnSL1 amendment for sandy soil, and sandy soil blank) after lyophilization.

5.2.3 Analytical methods

5.2.3.1 Total Se and Zn concentration in plants

The digestion method used to decompose the plant matrix prior to Se and Zn analysis has been described in Chapter 3. Total Se and Zn were determined in the digests by inductively coupled plasma mass spectrometry (ICP-MS, ELAN DRC-e, PerkinElmer, Waltham, MA, USA). Internal standards ($10\text{ }\mu\text{g/L}$ ^{103}Rh and ^{69}Ga) and an external multi-element standard solution were used during ICP-MS analysis to validate the accuracy of Se and Zn measurements. White clover samples (BCR-CRM, $6.7 \pm 0.25\text{ mg Se/kg}$ dry weight) and sea Lettuce ($51.3 \pm 1.3\text{ mg Zn/kg}$ dry weight) were included as certified reference materials for Se and Zn, respectively, in each analytical batch as quality control with recoveries of $107 (\pm 6\%)$ and $97 (\pm 7\%)$.

5.2.3.2 Selenium speciation analysis

Selenium speciation in the SeZn-enriched duckweed and harvested seeds of beans was analyzed by ICP-MS coupled to high-performance liquid chromatography (Series 200 HPLC, Perkin Elmer, Waltham, MA, USA). Selenium speciation in the SeZn-enriched sludge was determined by extended X-ray absorption fine structure (EXAFS) and X-ray absorption near edge structure (XANES). Additional details on the analysis can be found in Chapter 4.

5.2.3.3 pH, TC and TOC analysis of the extracted soil pore water

pH, TC and TOC of the extracted soil pore water at the end of the experiment were analyzed as described in Chapter 4.

Chapter 5

5.2.4. Statistical analysis

Descriptive statistics were performed using Sigma plot 13, Excel 2016 and SPSS 20.0. Results are expressed as mean $\pm$ SD. The different treatments were compared with a one-way analysis of variance (ANOVA) and Duncan's multiple comparison tests ($P < 0.05$).

5.3 Results and discussion

5.3.1 Characterization of SeZn-enriched biomaterials

The Se species present in the SeZn-enriched duckweed and granular sludge are shown in Figure 5.1a and 5.1b, respectively. Se in the SeZn-enriched duckweed was composed of 92% selenate (Se(VI)), 1.3% selenite (Se(IV)), 0.4% Se-cystine (SeCys₂), 3.7% methylselenocysteine (SeMetSeCys) and 4.9% Se-methionine (SeMet). Differently, the predominant Se species in SeZn-enriched sludge was the zerovalent form (72.9%) of Se, followed by Se(IV) (27.5%).

5.3.2 pH and TOC content of the extracted soil pore water

The amendment of SeZn-enriched duckweed and sludge notably increased the pH of the soil pore water extracted from the two soils (Table 5.1) ($P < 0.05$). Likewise, the TOC content of the soil pore water was increased with the amendment of SeZn-enriched duckweed, while it remained unchanged upon the application of SeZn-enriched sludge in both soils (Table 5.1). The maximum TOC concentration was observed upon the application of SeZnDW5, being 11 and 4 times higher than that of the blank in the sandy and loamy soil, respectively. The increase of soil pH with biomaterials supplementation may be partially explained by proton (H⁺) exchange between the soil and the added biomaterials. In biomaterials, anionic function groups are present and during the decomposition of the applied biomaterials in the soil, these may consume protons and increase the soil pH. Previous studies have also shown that application of organic materials, such as plant materials, animal manure, household compost, and sewage sludge, can lead to soil pH increase (Naramabuye & Haynes, 2006; Opala et al., 2012).

Chapter 5




Figure 5.1. (a) Chromatogram of Se speciation in the SeZn-enriched duckweed compared with that of a Se standard solution of 100 µg/L of each species (1) SeCys₂, (2) SeMetSeCys, (3) selenite (Se(IV)), (4) SeMet, (5) selenate (Se(VI)), and (b) XANES spectra of the Se species in the SeZn-enriched sludge (green) as compared to standards.

Chapter 5

Table 5.1. Characterization of Rhizon extracts after 42 days of SeZn-enriched biomaterials amended in the two soils. Different lower case or upper case letters in the same column, respectively, indicate statistically significant differences between different amounts of SeZn-enriched duckweed or sludge application according to Duncan's multiple comparison tests ($P < 0.05$). No letters indicate $P > 0.05$.

Amendments	Sandy soil		Loamy soil	
	pH	TOC (mg/L)	pH	TOC (mg/L)
Blank	4.9 ± 0.5 c B	34 ± 5.2 b ^A	6.2 ± 0.2 b B	67 ± 16.2 b ^A
SeZnDW1	5.8 ± 0.5 b	62 ± 11.4 b	6.3 ± 0.5 b	83 ± 24.5 b
SeZnDW5	7.8 ± 0.2 a	378 ± 53.3 a	6.9 ± 0.1 a	256 ± 43.4 a
SeZnSL1	6.9 ± 0.4 A	51 ± 9.7 ^A	7.1 ± 0.3 A	46 ± 10.6 ^A
SeZnSL5	7.5 ± 0.1 A	69 ± 13.5 ^A	7.3 ± 0.0 A	40 ± 7.2 ^A

5.3.3 Selenium content in the soil pore water during plant growth

Amendment with SeZn-enriched duckweed and sludge significantly increased the Se concentration in the pore water of both soils compared to the blank (Fig. 5.2). The effect of the Se dosage was also evident from the higher Se concentration in the soil pore water for the most Se-enriched biomaterials (SeZnDW5 and SeZnSL5). After 7 days of incubation (before planting), the supplement of SeZnDW1 and SeZnDW5 increased the Se concentration in the pore water from the soil background values (around 1.0 $\mu\text{g/L}$) to 2491 and 3789 $\mu\text{g/L}$ for the sandy soil, and 2364 and 2540 $\mu\text{g/L}$ for the loamy soil, respectively (Fig 5.2a and 5.2b). The supplement of SeZnSL1 and SeZnSL5 raised the Se content in the pore water to 49 and 175 $\mu\text{g/L}$ for the sandy soil, and 30 and 190 $\mu\text{g/L}$ in the loamy soil pore water, respectively (Fig 5.2c and 5.2d).

The Se concentrations in the soil pore water significantly decreased along with the incubation time for the SeZn-enriched duckweed amendment, while they slightly declined for the SeZn-enriched sludge amendment. During the entire experiment (6 weeks), an approximate 93–98% decrease of the Se concentrations in the soil pore water was observed for the SeZn-enriched duckweed amendment, while the Se concentrations in the soil pore water decreased by 21–59% for the SeZn-enriched sludge amendment. It should be noted that the decrease of the Se concentration in the soil pore water with the amendment of SeZnDW5 was sharper than that of SeZnDW1

Chapter 5

in the initial phase of the experiment (Fig. 5.2a and 5.2b), resulting in a lower Se concentration in the soil pore water with the SeZnDW5 application in comparison with the SeZnDW1 application after 21 days. The Se concentration in the soil pore water with SeZnDW5 supplement decreased by 97 and 90% in the first 21 days for the sandy and loamy soil, respectively, and remained almost stable in the remaining part of the experiment.

The Se concentration in the pore water of the two soil types supplied with SeZn-enriched duckweed was significantly higher than that of the soils supplied with SeZn-enriched sludge (Fig. 5.2a and 5.2b versus Fig. 5.2c and 5.2d). In addition, the Se concentrations in the sandy soil pore water were slightly higher than those in the loamy soil pore water, which could be the result of the different soil characteristics. The loamy soil has a higher content of organic matter (OM), CEC and EC, and a higher percentage of silt and clay, in comparison with the sandy soil (Table 4.1 in chapter 4). Generally, there are more oxygenic groups in soil OM, such as phenolic hydroxyl and carboxyl groups (Li et al., 2015b). These can complex or chelate soluble Se in soil, thereby decreasing the Se content in the soil pore water (Coppin et al., 2006; Li et al., 2015b).

The increased Se content in the soil (or pore water) upon Se-enriched biomaterials amendment has also been observed in our previous work (Chapter 4) and in other studies (Bañuelos et al., 2016; Wang et al., 2018). In line with our previous results (Chapter 4), the remarkable decrease of the Se content in the soil pore water with the amendment of SeZn-enriched duckweed along with the growth time could be attributed to (1) the plant uptake; (2) the immobilization of the released Se by the soil matrix; and (3) the adsorption of the extra OM introduced by the duckweed supplementation. The stable Se content in the soil pore water with the supplementation of SeZn-enriched sludge during the entire growth period could be due to the Se(0) species predominating in the sludge samples (Fig. 5.1), as Se(0) is highly stable in the environment (Hu et al., 2018; Zahedi et al., 2019).

Chapter 5




Figure 5.2. Evolution of the Se concentration in the pore water of the sandy and loamy soils with biomaterials supplementation: (a) sandy soil with SeZn-enriched duckweed supplementation, (b) loamy soil with SeZn-enriched duckweed supplementation, (c) sandy soil with SeZn-enriched sludge supplementation, and (d) loamy soil with SeZn-enriched sludge supplementation. Values are mean $\pm$ standard deviation (n=3).

5.3.4 Zinc content in the soil pore water during beans growth

Both SeZn-enriched biomaterials were effective for improving the availability of Zn in the two types of soil, which is reflected in a significantly higher amount of Zn in the pore water of soils amended with SeZn-enriched duckweed and sludge compared to that of the blank at the end of the experiment (Fig. 5.3). The application of SeZnDW5 showed the highest Zn content in pore water of both soils among all amendments in the first 3 days, indicating the fastest Zn release from this biomaterial (Fig. 5.3a and 5.3b). Along with the growth time, the Zn content rapidly decreased within 21 days and slightly decreased thereafter for this amendment (SeZnDW5), resulting in a 65% (Fig. 5.3a) and 35% (Fig. 5.3b) decrease of the Zn content in the sandy and loamy soil, respectively, over the entire growth period. On the contrary, the Zn content in the pore water of both soils increased (approximately 4 times) along with the growth time for the amendment of SeZnDW1, which even had no significant difference in comparison with

Chapter 5

that of the SeZnDW5 supplement after 21 days (Figs. 5.3a and 5.3b). Similarly, the supplement of SeZn-enriched sludge increased the Zn content in the soil pore water by 4–15 times from day 3 to day 42 (Figs. 5.3c and 5.3d). These results indicate that Zn is being slowly released from the SeZn-enriched biomaterials, except for the amendment of SeZnDW5. The reason for the noticeable decrease of the Zn content over time in the soil pore water after SeZnDW5 supplementation could be similar to that of the Se change (Fig. 5.2), i.e. the adsorption onto the extra organic matter (introduced by the high amount of SeZn-enriched materials application) (Table 5.1).




Figure 5.3. Evolution of Zn concentration in the pore water of sand and loamy soils amended with SeZn-enriched biomaterials: (a) sandy soil with SeZn-enriched duckweed supplementation, (b) loamy soil with SeZn-enriched duckweed supplementation, (c) sandy soil with SeZn-enriched sludge supplementation, and (d) loamy soil with SeZn-enriched sludge supplementation. Values are mean $\pm$ standard deviation (n=3).

Additionally, the Zn content of the pore water of soils supplied with SeZn-enriched duckweed was significantly higher than that of the soils supplied with SeZn-enriched sludge (Fig. 5.3 a and b versus Fig 5.3 c and d), which corresponds to the significantly higher Zn content in the SeZn-enriched duckweed (2289 mg Zn/kg dry weight)

Chapter 5

compared to that in the SeZn-enriched sludge (563 mg Zn/kg dry weight). The Zn concentration in the pore water of the loamy soil was higher than that of the sandy soil (1.1–2.6 times and 1.1–5.6 times for the supplementation of SeZn-enriched duckweed and sludge, respectively), except for the SeZn-enriched duckweed application at the beginning, i.e. day 7 (Fig. 5.3).

5.3.5 Biomass of beans

Fig. 5.4 shows the biomass of beans grown in soils amended with Se-enriched biomaterials, non-enriched biomaterials (the corresponding controls), and without any biomaterials (Blank). The biomass yield of beans was related to the duckweed application dose ($P < 0.05$). Specifically, no significant influence on the dry biomass weight was observed for the sandy soil amended with SeZnDW1 and control-SeZnDW1 in comparison with the blank, while the duckweed amendment SeZnDW5 significantly decreased the dry weight of each beans tissue by 51-85% and an even over 85% decrease was noted for the control-SeZnDW5 (Figure 5.4a). Likewise, for the loamy soil, the amendment SeZnDW1 significantly increased the beans biomass of each tissue (up to 159%), while an over 48 and 72% decrease was observed for the amendment of SeZnDW5 and control-SeZnDW5, respectively (Figure 5.4b). On the other hand, there was no remarkable difference between the blank and the different amounts of SeZn-enriched sludge applied (Figure 5.4c and 5.4d). It should be noted that the Se and Zn present in the biomaterials seem to counteract the negative effects of the biomaterials themselves on the growth of beans, which was reflected in the slightly higher biomass observed for the SeZn-enriched biomaterials amendment compared to that of the control-biomaterials amendment (Fig. 5.4).

The remarkable decrease of the beans biomass upon amendment of SeZnDW5 might be due to the high amount of duckweed (24g in 0.5kg soil) applied to the soil. Some phytotoxic substances such as low molecular weight organic acids, phenols and other allelochemicals may be released at the early stage of the fresh duckweed decomposition, which may inhibit the growth of the beans (Dinh et al., 2020; Jin et al., 2020). Besides, a high amount of duckweed applied to the soils may increase the C/N ratios, promoting the use of mineral nitrogen from the soil by microbial activities and resulting in competition for nitrogen between the soil microorganisms and the crop (Jin et al., 2020; Witt et al., 2000), thus reducing the beans plant biomass yields. Moreover,

Chapter 5

accumulation of a large amount of duckweed in the soil may prevent the roots of the beans plant to completely contact the soil, thus hindering plants to take up nutrients from soil and seriously affecting the growth (Jin et al., 2020). Additionally, the decrease of plant biomass with the higher amount of biomaterials may be attributed to the introduction of excessive soil alkalinity which could cause salinization after biomaterials fertilization, or a disturbance of the ionic balance in the soil solution thus affecting productivity. This is consistent with the results of Antonkiewicz et al (2020) and Schumann & Sumner (2004). Nevertheless, future research is needed to further reveal the causes of the lower bean biomass yield upon application of high amounts of duckweed.




Figure 5.4. Dry biomass of the bean plant tissues grown on the sandy and loamy soils with SeZn-enriched biomaterials supplementation: (a) sandy soil with SeZn-enriched duckweed supplementation, (b) loamy soil with SeZn-enriched duckweed supplementation, (c) sandy soil with SeZn-enriched sludge supplementation, and (d) loamy soil with SeZn-enriched sludge supplementation. Values are mean $\pm$ standard deviation ($n=3$). Different lower-case letters indicate statistically significant differences between amendments within the same tissues according to Duncan's multiple comparison tests ($P < 0.05$). Different upper-case letters indicate the statistically

Chapter 5

significant differences in the total dry weight between amendments according to the Duncan's multiple comparison tests ($P < 0.05$).

5.3.6 Selenium content in the beans grown on soils amended with SeZn-enriched biomaterials

The amendment of SeZn-enriched duckweed and sludge remarkably increased the Se concentration in the different tissues of beans (seed, leaf, stem and root) in comparison with the blank (Fig. 5.5). More specifically, increasing the sludge amendment from SeZnSL1 to SeZnSL5 increased the Se concentration in the plant tissues by 3.1–4.3 times. However, the Se content in all tissues of the beans, except for the roots, decreased by 1.1–4.0 times when the SeZn-enriched duckweed amendment was increased from SeZnDW1 to SeZnDW5.

The Se concentration in all tissues of the beans amended with SeZnDW1 was 2.3–4.1 times higher than in those amended with SeZnSL1, except for in the stems (7–9 times) and similar values in the roots. However, the amendment of SeZnSL5 resulted in an approximately 1.6–3.0 times higher Se content in all beans tissues compared to the amendment of SeZnDW5. Selenium was mainly accumulated in the roots after being taken up by the bean plants, especially for the amendment of SeZnSL1 and SeZnSL5, while Se was more rapidly translocated from the roots to aboveground biomass when SeZnDW1 was amended.

Similarly, our previous work (chapter 4) and some other studies (Bañuelos et al., 2015; Bañuelos et al., 2016) also indicated that micronutrient (Se)-enriched supplements can successfully increase the Se content in crops and plants species after being applied into soils, such as the supplementation of the Se-enriched hyperaccumulator *Stanleya pinnata* biomass to sandy-loam soil. However, the higher Se content in the tissues of beans upon amendment of SeZnDW1 compared to SeZnDW5 was attributed to the negative effects of the high amount of duckweed (SeZnDW5) on the plant growth and nutrients accumulation, which was indicated by the remarkable decrease of beans biomass (Fig. 5.4). The different translocation rate of Se from the roots to aboveground biomass (stem, leaf, and seed) between the supplement of SeZn-enriched duckweed and sludge may be due to the different Se species dominating in the two biomaterials. Specifically, 92% Se(VI) being present in the SeZn-enriched duckweed (Fig. 5.1a) may

Chapter 5

result in a high amount of Se(VI) released in the soil pore water. Se(VI) is highly soluble and readily translocated from underground to aboveground plant parts after being taken up by plants (Peng et al., 2019). In contrast, the zerovalent form (72.9%) of Se was predominant in the SeZn-enriched sludge, which may be released as such or converted into Se(IV) under toxic conditions and mainly accumulate in the roots.




Figure 5.5 Se concentration in the different tissues of the harvested beans grown on sandy and loamy soils amended with SeZn-enriched biomaterials: (a) sandy soil with SeZn-enriched duckweed supplementation, (b) loamy soil with SeZn-enriched duckweed supplementation, (c) sandy soil with SeZn-enriched sludge supplementation, and (d) loamy soil with SeZn-enriched sludge supplementation. Values are mean $\pm$ standard deviation (n=3). Different letters indicate statistically significant differences between the different doses within the same tissues according to Duncan's multiple comparison tests ($P < 0.05$).

Additionally, the Se concentration ranged from 5.1 to 13.5 mg/kg and 3.5 to 17 mg/kg in the seeds of beans with the supplementation of SeZn-enriched duckweed and sludge, respectively. In this regard, the Superior Health Council in Belgium implemented a daily dietary allowance (RDA) of 70 μ g Se/day for human diets. If we

Chapter 5

assume that a daily serving size of the Se-enriched beans seeds is 96 g fresh weight (or 7.2 g dry weight due to its 92.5% water content calculated in this study), consumption of the highest Se containing bean seeds (13.5 and 17 mg Se/kg as observed in SeZnDW1 and SeZnSL5) would result in the ingestion of 97.2 and 122.4 µg Se. This means that the recommended daily Se requirements would be safely met and not substantially exceeded.

5.3.7 Zinc content in the beans grown on soils amended with SeZn-enriched biomaterials

Fig. 5.6 illustrates the Zn content in different tissues of the beans grown on sandy and loamy soil amended with SeZn-enriched duckweed (Fig. 5.6a and 5.6b) and sludge (Fig. 5.6c and 5.6d). An increasing Zn content in the different bean tissues was associated with the amendment of SeZn-enriched duckweed, while contrary results were observed with the amendment of SeZn-enriched sludge in all bean tissues except for the roots. Specifically, increasing the SeZn-enriched duckweed application from the blank to SeZnDW5 increased the Zn content in the roots stepwise from 168 to 780 mg/kg for the sandy soil and from 541 to 931 mg/kg for the loamy soil (Fig. 5.6a and 5.6b). On the other hand, the application of SeZn-enriched sludge declined the Zn concentrations by 20–34% in the stems, leaves and seeds. Besides, no significant difference of the Zn content in the roots between the bank and the amendment of SeZn-enriched sludge was observed (Fig. 5.6c and 5.6d).

Zn mainly accumulated in the roots after being taken up by the beans with the amendment of SeZn-enriched duckweed, as shown by the higher ratio of Zn content between the roots and aboveground tissues (3–11 times for the sandy soil, 5–13 times for the loamy soil). This indicates that Zn was not easily translocated in the plants, which is consistent with other studies (Fontes et al., 2014; Mao et al., 2014; Montalvo et al., 2016). For instance, Mao et al. (2020) found that Zn applied to the soil in the form of Zn sulfate at 232.7 kg/ha did not increase the Zn concentrations in the edible parts of wheat, maize and soybean, due to the limited transportation ability of Zn by the phloem to the grain/seed.

The decrease of the Zn content in the bean tissues with the supplementation of SeZn-enriched sludge could be attributed to the competition between Zn and other ions (e.g.

Chapter 5

Se) and the increase of soil pH. Specifically, the higher amount of Se in the soil pore water (released from the SeZn-enriched sludge) (Fig. 5.3) and in the plants (Fig. 5.5) may inhibit the uptake and translocation of Zn by beans. The antagonistic effect of Se and Zn on plant uptake has also been observed in other plants, such as wheat, cabbage and potato (Mao et al., 2014; Singh & Singh, 1978). Besides, pH is the most important factor in determining Zn solubility (Montalvo et al., 2016). Increasing pH values could increase the complexation of Zn with dissolved organic matter (e.g. humic and fulvic acids), resulting in the lower bioavailability of Zn (Dinh et al., 2020). In this study, the pH of the soil pore water was remarkably increased by the supplementation of SeZn-enriched biomaterials (from 4.9 to 7.5 when sludge is amended in sandy soil, and from 6.2 to 7.3 in loamy soil) (Table 5.1), which may therefore partially explain the lower Zn content in the bean tissues.




Figure 5.6. Zn concentration in the different tissues of beans grown on sandy and loamy soils amended with SeZn-enriched biomaterials: (a) sandy soil with SeZn-enriched duckweed supplementation, (b) loamy soil with SeZn-enriched duckweed supplementation, (c) sandy soil with SeZn-enriched sludge supplementation, and (d) loamy soil with SeZn-enriched sludge supplementation. Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate the statistically significant differences among

Chapter 5

amendments within the same tissues according to Duncan's multiple comparison tests ($P < 0.05$).

The supplementation of the SeZn-enriched biomaterials did not successfully improve the Zn content in the seeds of beans (Fig. 5.6), and the biofortification purpose of Zn is thus not achieved. This could be attributed to the lower Zn translocation rate between plant roots and seeds, and the application of a too low Zn dose, particularly for the SeZn-enriched sludge application. As aforementioned, the lower translocation rate of Zn in plants was reflected in the remarkably higher Zn content in the bean roots after SeZn-enriched duckweed application, with no difference in aboveground biomass among all treatments (Fig. 5.6a and 5.6b). On the other hand, the Zn content in the SeZn-enriched sludge was only 563 mg/kg (vs 2289 mg/kg in SeZn-enriched duckweed), and the applied amounts of SeZnSL1 and SeZnSL5 were 1.7 and 8.7 g/pot, which may not be able to provide sufficient Zn for plant uptake. This suggests that the Zn content in the sludge should be reconsidered during its production.

5.3.8 Se species in the beans

The seeds of beans grown on the sandy soil without (blank) and with the amendment of SeZnDW1 and SeZnSL1 were selected for the measurement of Se speciation (Fig. 5.7 and Table 5.2). The recovery of Se in all seeds ranged from 82 to 99% of the total Se content after protease hydrolysis.

Organic Se (SeCys2, SeMetCys and SeMet) was the main Se species in all seed samples. SeMet accounted for the highest percentage of the total Se in the seeds for the blank (77.2%), SeZnDW1 (68.5%) and SeZnSL1 (82.3%). Besides, the percentage of inorganic Se species in the seeds with the SeZnDW1 amendment and blank was much higher compared to that of the SeZnSL1 amendment. Particularly for Se(VI), 13.7, 10.6, and 2.4% of the total Se was detected in the seeds of the blank, SeZnDW1 and SeZnSL1 amendment, respectively.

Organic Se species are beneficial for dietary intake by humans and animals, as they are more bioavailable and less toxic (Gómez-Jacinto et al., 2020) compared to inorganic Se species. SeMet and SeCys can be incorporated at the active sites of a wide range of selenoproteins involved in major metabolic pathways, such as thyroid hormone metabolism, antioxidant defense, and immune function (Malagoli et al., 2015).

Chapter 5

Therefore, the high percentage of organic Se species present in the seeds of the beans indicates that Se in the SeZn-enriched duckweed and sludge could be taken up and transformed into more valuable Se species. These results demonstrated the potential of SeZn-enriched biomaterials produced from wastewater as Se biofertilizers.

Table 5.2. Percentage of Se species in the seeds of beans grown on the sandy soil amended with SeZnDW1 and SeZnSL1 and without amendment (Blank) relative to total Se in bean seeds.

	Total Se	SeCys ₂	SeMetCys	SeMet	Se(IV)	Se(VI)	Se species recovery
	(mg/kg)			(%)			
Blank	0.1	3.2	3.5	77.2	2.4	13.7	99%
SeZnDW1	11.6	1.1	2.1	68.5	0.3	10.6	82%
SeZnSL1	4.1	1.4	4.5	82.3	0.4	2.4	91%

Chapter 5




Figure 5.7. Se speciation in harvested seeds of beans after 42 days of growth on sandy soil amended with (a) SeZnDW1 and (b) SeZnSL1 and the (c) Blank soil (higher magnification in insert) together with (d) Se standard solution of 10 µg/L of each Se species.

5.4 Conclusions

The potential of SeZn-enriched biomaterials produced from simulated wastewater as Se biofertilizers has been proven, as illustrated by the remarkably higher Se content in the soil pore water and plant tissues upon amendment of soil with these biomaterials. The highest dose of SeZn-enriched duckweed (SeZnDW5) inhibited the plant growth and noticeably raised the pH and TOC content of soils, confirming that appropriate amounts of biomaterials should be applied. SeZn-enriched sludge is considered as the preferred Se biofertilizer in comparison with SeZn-enriched duckweed, as the Se slowly released from the SeZn-enriched sludge remained stable in the soil pore water, which could slowly provide sufficient Se for plant uptake and transformation to selenoamino acids (SeMet). On the other hand, the agronomic application of the SeZn-enriched duckweed and sludge may not be an effective alternative for Zn biofortification, as reflected by the lower Zn content in the bean seeds. This should be optimized by increasing Zn content in the SeZn-enriched biomaterials.

**Chapter 6 Production of selenium-enriched microalgae as potential
feed supplement in high rate algal ponds treating domestic
wastewater**

This chapter was redrafted after:

Li, J., Otero-Gonzalez, L., Michiels, J., Lens, P.N.L., Du Laing, G., Ferrer, I., 2021. Production of Se-enriched microalgae as potential feed supplement in high rate algae ponds treating domestic wastewater. *Bioresource Technology*. 333, 125239.

Abstract

This study assessed the selenium (Se) removal efficiency of two pilot-scale high rate algal ponds (HRAPs) treating domestic wastewater and investigated the production of Se-enriched microalgae as potential feed supplement. The HRAPs were operated for 3 months under two hydraulic retention times (8 and 4 days) with corresponding average organic loading rates of 66 and 127 mg COD/L·day, respectively. The HRAP-Se had a selenium loading rate of 6.28 µg Se/L·day, while the HRAP without Se spiking served as control (HRAP-C). The wastewater treatment efficiency of the HRAPs, the Se content and speciation, in vitro digestibility, nutritional value, and pathogen load of the microalgae grown in the HRAP were evaluated. The HRAP-Se had an average Se, NH₄⁺-N, total phosphorus, total COD, total carbon removal efficiency of, respectively, 43%, 93%, 77%, 70% and 67%. Inorganic Se taken up by the microalgae was mainly (91%) transformed into valuable selenomethionine (SeMet), and 49–63% of Se in the Se-enriched microalgae was bioaccessible for animals through in vitro digestibility tests. Besides, the crude protein content (around 48%) of the microalgae was higher than that of conventional plant-based protein sources in feed (soybeans and soybean meals), whereas the essential amino acid content of the microalgae was comparable to that of soybeans. Fatty acid profile analysis demonstrated that Se may induce the production of the polyunsaturated fatty acids omega-3 and omega-6 in microalgae, particularly for eicosapentaenoic (EPA). Microbiological analysis indicated that downstream drying processes of microalgae could avoid pathogen contamination. Although more research is still further needed to confirm the results, this study showed that the production of Se-enriched microalgae in HRAPs may offer a promising alternative for upgrading of low-value recovered resources into high-value feed supplements, supporting the drive to a circular economy.

Keywords: Selenium, algae, HRAPs, photobioreactor, resource recovery, wastewater treatment

Chapter 6

6.1 Introduction

Nowadays, microalgae-based wastewater treatment technologies are attracting considerable attention, as they are low-cost, low-energy consuming and easily implemented in regions with high temperatures and sunlight exposure (Arashiro et al., 2019). Microalgae have a great capacity to remove/take up excess nutrients from the corresponding growth medium, as their cultivation requires high amounts of macro-nutrients (such as nitrogen and phosphorus) and micro-nutrients (Arashiro et al., 2020a; Gan et al., 2019). Furthermore, microalgae are a potential source for the production of protein-rich biomass and numerous other high-value compounds, e.g. fatty acids, pigments and vitamins (Markou et al., 2018). Microalgae based products grown in nutrient medium are adding value to the market. On the other hand, wastewater contains nutrients that could serve as feedstock, providing free nutrients for microalgae growth. In this context, the cultivation of microalgae on wastewater with nitrogen, phosphorus and organic matter removal does not only provide an option to treat wastewater, but also significantly reduces the cost and carbon footprint of conventional microalgae production systems that does not use wastewater as a growth medium, by converting low-value resources in wastewater into high value-added bioproducts (Borowitzka & Moheimani, 2013; Silambarasan et al., 2021).

Furthermore, Se supplementation of feed and food to overcome the Se deficiency received much attention in recent years. Microalgae may also have the ability to take up inorganic Se and incorporate it into amino acids forming selenoamino acids, such as selenomethionine (SeMet) and Se-cystine (SeCys₂), which are beneficial for animal and human health (Umysova et al., 2009; Winkel et al., 2015). In this context, high-value microalgae biomass with high Se content may be generated during wastewater treatment processes and subsequently added to animal feed for alleviating Se deficiency issues in animals. The Se source for this process could be Se-rich wastewaters, but Se could also be added from a primary source to produce Se-rich microalgae on domestic wastewater, providing the macronutrients for microalgae growth, as feedstock. Accordingly, a higher-value animal feed product could be produced from wastewater, while recovering resources.

Chapter 6

Therefore, in this study, microalgae were grown in two pilot-scale high rate algal ponds (HRAPs) treating domestic wastewater with and without Se spiking in order to: (1) investigate the Se removal efficiency of pilot-scale HRAPs treating domestic wastewater, (2) evaluate the possible use of domestic wastewater as a nutrient source for microalgae growth in HRAPs to produce high-value Se-enriched microalgae, and (3) assess the potential use of upgraded Se-enriched microalgae as feed supplement by examining their Se content and speciation, digestibility, biochemical properties and nutritional profile.

6.2 Materials and methods

6.2.1 Source of biomass and wastewater

The microalgae inoculum was collected from a demonstrative-scale photobioreactor treating agricultural runoff (90%) and domestic wastewater (10%) located outdoors at the Agròpolis experimental campus of the Universitat Politècnica de Catalunya-BarcelonaTech (Viladecans, Spain). Operational details of the photobioreactor and characteristics of the biomass were presented by García et al. (2018).

The wastewater used in this study was real domestic wastewater from a residential area close to the Universitat Politècnica de Catalunya-BarcelonaTech (Barcelona, Spain), as described by Arashiro et al. (2019). The experimental set-up was located outdoors. Domestic wastewater received a screening pretreatment before being pumped into a 1 m³ homogenization tank that was continuously stirred to avoid solids sedimentation, followed by a 3 L primary settler (diameter: 18 cm, height: 30 cm) with a hydraulic retention time (HRT) of 41 min. The effluent from the primary settler (noted as primary effluent) was collected (Fig. 6.1) for the subsequent batch experiments or pumped into two parallel HRAPs (0.5 m³ each) as influent of the continuous system. Each HRAP, constructed from PVC, had a surface area of 1.54 m², a water depth of 0.3 m, a working volume of 0.47 m³ and a paddle-wheel constantly stirring the mixed liquor at an average velocity of 10 cm/s. Two secondary clarifiers (10 L) followed the two HRAPs to separate the effluent and biomass. The scheme of the HRAPs is shown in Fig. 6.1.

The average values of the main parameters (e.g. pH, total suspended solids (TSS), volatile suspended solids (VSS), chemical oxygen demand (COD) and ammonium

Chapter 6

nitrogen ($\text{NH}_4^+\text{-N}$), among others) in the primary effluent that was pumped to the HRAPs and the secondary effluent from the HRAPs clarifiers through the entire experimental period (3 months) are presented in Fig. 6.2 and Table 6.S1 in supplementary information.




Figure 6.1. Scheme of the HRAPs treating domestic wastewater. HRAP-Se is the line with Se spiking and HRAP-C is the line without Se spiking, which served as control. Sampling points are 1: primary effluent (also called influent of the HRAPs), 2: mixed liquor of the HRAPs, 3: secondary effluent.

6.2.2 Se removal by microalgae in batch experiments

The mixed microalgae consortium was cultivated in a 3-L batch photobioreactor fed with the primary effluent for 2 weeks, which served as the microalgae inoculum for the subsequent batch experiments. A photon flux density of $120 \mu\text{mol}/\text{m}^2/\text{s}$ was provided by two cool-white fluorescent lamps with a 12 h/12 h of light/darkness photoperiod at 25°C . The microalga biomass was continuously mixed with a magnetic stirrer. pH was continuously monitored with a pH sensor (HI1001, HANNA, U.S.A.) and maintained at 7.8 with a pH controller (HI 8711, HANNA, U.S.A.) by the automated addition of 0.1 M HCl and NaOH. This lab-scale set-up was located indoors.

Harvested biomass from the photobioreactor was thickened by gravity settling in Imhoff cones and then the cell number of the thickened biomass was counted by microscopy

Chapter 6

(BA310, Motic, China). The thickened biomass was added into 300 mL Erlenmeyer flasks containing 200 mL of the primary effluent to make cultures with an initial density of 1×10^6 cells/mL. Sodium selenite (Na_2SeO_3) or sodium selenate (Na_2SeO_4) was added to the Erlenmeyer flasks before inoculation at a Se concentration of 0, 10, 25, 50, 100, 200 and 500 $\mu\text{g/L}$. The cultures were mixed with magnetic stirrers and incubated for 7 days under the same light intensity and photoperiod as described above. All experiments were conducted in duplicate. pH and turbidity were monitored daily. 10 mL of medium was collected and filtered every other day for Se concentration analysis. After 7 days of incubation, the biomass was centrifuged and dried for total Se measurement.

6.2.3 Se removal in HRAPs and production of Se-enriched biomass

Experiments were carried out in an outdoor pilot plant (May 2019–July 2019) as described in detail by Arashiro et al. (2019) with some modifications. The microalgae species in the HRAPs were observed microscopically (BA310, Motic, China) every week. The medium was mainly composed of *Chlorella* sp.. The effluent from the primary settler (noted as primary effluent) was pumped into two parallel HRAPs: one with continuous spiking with Na_2SeO_3 (HRAP-Se) and another one without Se spiking as a control (HRAP-C).

The two HRAPs received the corresponding influents (53 L/day of wastewater and 6 L/day of Se stock solution (500 $\mu\text{g Se/L}$) for the HRAP-Se, and 59 L/day of wastewater for the HRAP-C) with an HRT of 8 days during the first 1.5 months (in spring season). Afterwards, the HRT was adjusted to 4 days (in summer season) until the end of the experiment, by increasing the influent flow rates to twice the level previously mentioned. The change of HRT from 8 (in spring season) to 4 days (in summer season) has been demonstrated to improve the wastewater treatments performance by HRAP by the previous studies (Arashiro et al., 2019). The flow rates of Se spiking and wastewater in the HRAP-Se were monitored daily to accurately quantify the Se concentration in the influent. The flow rates of the Se spike and the wastewater in the HRAP-Se were monitored daily to accurately quantify the Se concentration in the influent. The effluent was collected daily for total Se analysis. The biomass in the secondary clarifiers was accumulated and collected every week.

Chapter 6

6.2.4 Wastewater characterization in the HRAP systems

The wastewater treatment performance was monitored for 3 months. Samples from the influent, effluent and mixed liquor of the two HRAPs (Fig. 6.1) were collected twice per week for analysis of the following parameters: pH, dissolved oxygen (DO), turbidity, TSS, VSS, total and soluble COD (COD_{tot} and COD_{sol}), total and soluble P (TP and SP), nitrite (NO_2^-), and nitrate (NO_3^-); these parameters were analyzed according to standard methods (APHA-AWWA-WEF, 2012). $\text{NH}_4^+\text{-N}$ was measured according to the Solórzano method (Solórzano, 1969). Total carbon (TC) and total nitrogen (TN) were measured by a N/C-analyzer (multi N/C 2100S, Analytik Jena, Germany) as described by Arashiro et al. (2019). All analyses were conducted in triplicate. Selenium concentration in wastewater was measured using inductively coupled plasma-mass spectrometry (ICP-MS, ELAN DRC-e, PerkinElmer, Waltham, MA, USA) after being filtered over a 0.45- μm syringe PVDF membrane filter.

6.2.5 Nutritional parameters of the microalgae

The microalgae biomass collected in the secondary clarifier of the HRAPs at operational week 7 (from day 43 to 50) was rinsed with deionized (DI) water and centrifuged at 4200 rpm for 5 minutes. The centrifuged paste was frozen at $-80\text{ }^\circ\text{C}$ overnight and then lyophilized for 24 h. The freeze-dried biomass was stored in a $-20\text{ }^\circ\text{C}$ freezer for the subsequent analysis and experiments.

6.2.5.1 Se speciation, Se bioaccessibility and total Se analysis

Selenium speciation and total Se of the freeze-dried microalgae were determined as in the previous Chapter (Chapter 3). Besides, the bioaccessibility of Se in raw and bead milled microalgae for pigs was simulated in vitro in a two-step incubation based on the method described by Moheimani et al. (2018) and Vu et al. (2019) with minor modifications. Briefly, an amount of freeze-dried sample equivalent to 150 mg protein was weighed into a 100-mL centrifuge tube with 20 mL of simulated gastric juice (1 g pepsin dissolved into 500 mL of 0.075 M HCl) and one drop of 50 g/L thimerosal. The mixture was shaken in a reciprocating thermostatic shaking water bath at $37\text{ }^\circ\text{C}$ for 4 h. After gastric digestion, the mixture was cooled down and the pH was adjusted to 7.5 using 0.2 M NaOH followed by adding 15 mL pancreatin solution (375 mg pancreatin dissolved into 250 mL phosphate buffer) to simulate small intestine digestion. The

Chapter 6

mixture was shaken in a water bath at 37 °C for 4 h, followed by adding 7.5 mL of 0.02 M phosphotungstic acid for deproteination, and afterwards centrifuged at 3000 rpm for 10 min. The supernatant was collected and filtered by a 0.45-µm syringe PVDF membrane filter for analysis of the Se content, which was considered to represent the digestibility in the gastric and intestine phase. Se bioaccessibility was determined by the ratio of Se obtained from the gastrointestinal digestion divided by the total amount of Se in the corresponding biomass.

6.2.5.2 Macromolecular characterization and protein extraction by different cell disruption methods

Microalgae macromolecular composition (i.e., lipid, carbohydrate and crude protein) was determined and calculated over the VSS content. Lipids were extracted by chloroform and methanol (2:1) according to the Soxhlet extraction method (Folch et al., 1957). Carbohydrates were measured by phenol-sulphuric acid method with acid hydrolysis (Dubois et al., 1951) and determined by spectrophotometry (Spectronic Genesys 8, Helsingborg, Sweden). Total crude proteins were measured and quantified according to the total Kjeldahl nitrogen (TKN) method (Kjeldahl, 1883) with a TKN/protein conversion factor of 5.95 (Arashiro et al., 2019).

For different protein extraction method tests, 0.5 g freeze-dried microalgae biomass was dispersed and mixed into 25 mL PBS buffer solution. Five cell disruption methods for the microalgae suspension were investigated and compared: (a) freeze-thawing at -80 °C and 4 °C with 5 cycles; (b) combination of freeze-thawing and ultrasonication (Bandelin Sonouls HD2070, 20 kHz and 2 mm probe) for 30 min with 30 s on/off intervals at 70% amplitude; (c) high-pressure cell disruption (constant cell disruption systems with one-shot model, Northants, UK) at 2.4 kpsi; (d) ball milling (MM 400, Retsch, Haan, Germany) for 10 min at 30 Hz; and (e) bead milling (Powerlyzer 24, MO BIO Laboratories, Carlsbad, CA, USA) at 2000 rpm for 10 min. All experiments were performed in triplicate.

The protein content of the microalgae after each disruption was quantified by the Lowry method with minor modification (Lowry et al., 1951). In brief, 1.0 mL of cell suspension after disruption was vortex mixed with either 3 mL of 7.0% sodium dodecyl sulfate (SDS) solution or 3 mL DI water. The mixture was incubated at 100 °C for 5 min and

Chapter 6

cooled down before centrifuging at 4000 rpm for 10 min. 1.0 mL of the supernatant after centrifugation was collected and vortex mixed with 5.0 mL alkaline copper reagent. After 10 min, 0.5 mL Folin solution was added to react for 30 min in the dark place. A spectrophotometer (Spectronic Genesys 8, Helsingborg, Sweden) was used to measure the protein content by measuring the absorbance at 750 nm. A calibration curve was prepared using bovine serum albumin (BSA).

6.2.5.3 Fatty acid and amino acid profiles analysis

Fatty acids of the microalgae were analyzed as described by Michiels et al. (2014). Amino acids of the microalgae were analyzed by the lab of nutriFOODchem (Gent University, Belgium). Briefly, the freeze-dried microalgae sample was hydrolyzed with 6 M HCl for 24 h. After neutralization, the amino acids were derivatized in the injector of the HPLC, separated on a C₁₈ column and detected fluorometrically. Cysteine was derivatized in the injector of the HPLC with iodoacetic acid (IDA) and o-phthaldialdehyde (OPA), separated on a C₁₈ column and detected fluorometrically. All samples were analyzed in duplicate.

6.2.6 Pathogenic bacterial content

Pathogen loads (aerobic bacteria, coliforms, *E. coli*, *Listeria* and *Salmonella*) of fresh, oven-dried (at 70 °C until constant weight) and freeze-dried microalgae were evaluated according to the method of Montville and Matthews (2008) with three replicates. For fresh microalgae analysis, the biomass was collected, rinsed with DI water, and analyzed immediately

6.2.7 Statistical analysis

Descriptive statistics were performed using Sigma plot 13, Excel 2016 and SPSS 20.0. Results are expressed as mean $\pm$ standard deviation (SD).

6.3 Results and discussion

6.3.1 Microalgae growth, Se accumulation and Se removal in batch experiments

Fig. 6.2 shows the turbidity of the microalgae suspension when exposed to different Se concentrations (0–500 µg Se/L). The highest turbidity in both selenite (Se(IV)) (885 NTU) and selenate (Se(VI)) (1614 NTU) treatments was observed at 50 µg/L of Se exposure after 7 days of cultivation, being significantly ($p<0.05$) different from the control, which demonstrates that low Se application may stimulate microalgae growth. Fig. 6.2 further demonstrates that the turbidity significantly increased with incubation time, and a similar turbidity value was observed for the control treatments and the 500 µg Se/L selenite and selenate treatments, indicating that microalgae growing on domestic wastewater treatment could tolerate such high concentrations of Se.

Similarly, Li et al. (2003) reported that sodium selenite has either stimulating (0.5 mg Se/L) or toxic (500 mg Se/L) effects on *Spirulina platensis* growing in Zarrouk medium. Reunova et al. (2007) reported positive impacts of selenite on the unicellular alga *Dunaliella salina* (e.g., stimulation of cell growth) after exposure to 0.01 and 0.5 mg/L of Se dosed as sodium selenite in nutrient medium prepared in 32‰ seawater. Conversely, the inhibition of cell growth together with an increasing number of destroyed cells and cells with damaged organoids were observed after exposure to Se concentrations higher than 1.0 mg/L. Sun et al. (2014) found that Se(IV) concentrations lower than 75 mg Se/L in BG11 medium promoted *Chlorella vulgaris* growth and acted as an antioxidant by inhibiting lipid peroxidation and formation of intracellular reactive oxygen species (ROS). Accordingly, the growth-stimulating effects of Se for microalgae in this study may be also related to the enhancement of the antioxidant activity in cells, as Se can increase the activity of antioxidant enzymes (e.g., glutathione peroxidases, superoxide dismutase and methionine sulfoxide reductase) and the synthesis of metabolites (such as phytochelatins and ascorbate), resulting in higher ROS scavenging capacity of cells (Schiavon et al., 2017b; Sun et al., 2014), and eventually promoting microalgae growth.

Chapter 6




Figure 6.2. Biomass growth, measured as turbidity (NTU), during batch incubation in domestic wastewater supplemented with varying Se concentrations ($\mu\text{g/L}$), (a) selenite and (b) selenate. Values are mean $\pm$ standard deviation ($n=3$).

The Se concentration in the microalgae differed significantly depending on the chemical form and concentration of the applied Se (Fig. 6.3). Generally, increasing the Se dosage in the wastewater resulted in a higher Se concentration in the microalgae biomass. The microalgae had a higher ability to take up Se(IV) compared with Se(VI), which is reflected in the around 3 times higher Se content in microalgae cultivated in the selenite amended wastewater compared to the Se(VI) amended wastewater (Fig. 6.3). The maximum Se content in the microalgae biomass was 67 and 24 mg/kg when exposed to 500 $\mu\text{g Se/L}$ of Se(IV) and Se(VI), respectively. These values are much

Chapter 6

higher than the Se accumulation in the microalga *Spirulina platensis* (< 22 mg/kg) exposed to nutrient growth medium containing 500 µg Se/L of Se(IV) (Li et al., 2003), and also higher than the Se accumulation in the macroalga *Ulva australis* (around 20 mg/kg) exposed to 50 µM (equivalent to 4.0 mg/L) of Se(IV) or Se(VI) supplemented seawater after 7 days of incubation (Schiavon et al., 2016). Besides, the linear correlation ($R^2 > 0.99$) between the Se concentration in the microalgae and Se application dose indicates that the microalgae may still have the capacity to accumulate higher amounts of Se (Fig. 6.3).

A higher Se accumulation, when exposed to Se(IV) compared to Se(VI), has also been observed in other algae species. Vriens et al. (2016) reported 10 times more accumulation of Se(IV) than Se(VI) by the microalga *Chlamydomonas reinhardtii* when grown in 100 µM of Se (equivalent to 8.0 mg/L) nutrient growth medium for 24 h. The wild type of microalgae *Scenedesmus quadricauda* took up 2 times more Se(IV) than Se(VI) after exposure to 50 mg/L Se in mineral medium (Vitova et al., 2011). However, some other reports indicated a reversed result, for instance, Simmons and Wallschager (2011) found that *Chlorella vulgaris* had around 5-fold preference for Se(VI) uptake over Se(IV) upon 10 µg/L of Se exposure in 10% Bold's basal medium. These discrepancies are likely due to the different algal genus and species (Schiavon et al., 2017a; Simmons & Wallschager, 2011).

In this study, the higher uptake of Se(IV) compared to Se(VI) may be attributed to the different uptake mechanisms and metabolism by microalgae, partially similar to those in plants. Se(IV) is mostly taken up in a low-affinity passive way and quickly converted into organic Se forms (e.g., SeMet and SeCys₂) in algae (de Oliveira et al., 2017; Li et al., 2020b; Schiavon et al., 2017a). In contrast, Se(VI) is taken up in a high-affinity active way through the facilitation of a sulfur transporter, reduced to Se(IV) in cells and then converted into organic Se compounds (Arvy, 1993; Li et al., 2008). The Se(VI) reduction is an ATP-consuming process and the rate-limiting step, which eventually results in a lower Se uptake by microalgae (Schiavon et al., 2017a; Van Hoewyk, 2013). Besides, the lower Se(VI) uptake of microalgae could be partially attributed to the competition between Se(VI) and sulfate (164 mg/L) in the wastewater.

Chapter 6

The efficiency of Se removal by the microalgae is presented in Fig. 6.S1 of the supplementary information (SI). A decreasing trend was observed in the Se removal efficiency with increasing Se dose. Accordingly, when microalgae were exposed to Se(VI), the Se removal efficiency was much lower compared to Se(IV) exposure, which was associated with the lower Se(VI) uptake and accumulation in the microalgae cells (Fig. 6.3). Next to Se accumulation by microalgae biomass, Se volatilization could also partially contribute to the Se removal. The highest Se removal efficiency was 56 and 19% when microalgae were exposed to 10 $\mu\text{g/L}$ of Se(IV) and Se(VI), respectively. Se(IV) was therefore selected for the subsequent pilot-scale experiment due to the higher Se accumulation ability.




Figure 6.3. Se concentration in the microalgal biomass grown in wastewater with different selenite and selenate concentrations. Values are mean $\pm$ standard deviation ($n=3$).

6.3.2 Wastewater treatment efficiency and Se removal in HRAPs

The temporal variation and average values of the main parameters in HRAP-Se and HRAP-C over a period of 3 months are shown in Fig. 6.4 and Table 6.S1. A summary of the average removal efficiencies for the main water quality parameters is calculated and presented in Table S2. Likewise, the variation of Se content in the influent and effluent of HRAP-Se over the monitoring period is shown in Fig. 6.5. No significant differences were observed in the turbidity, TSS, VSS, total and soluble COD, $\text{NH}_4^+\text{-N}$, TN, TC, TP and SP removal efficiency throughout the entire experimental period

Chapter 6

between the HRAP-Se and HRAP-C (Fig. 6.4 and Table S1-2). The HRAP systems showed high nutrients and organic matter removal efficiencies. Specifically, the average $\text{NH}_4^+\text{-N}$ and turbidity removal efficiency reached 93% and 91%, respectively. The COD_{tot} and TC removal efficiency ranged between 70 and 66% in the HRAP-Se and HRAP-C throughout the whole experimental period. The average removal efficiencies of TP in HRAP-Se and HRAP-C were up to 77% and 72%, respectively. Despite the very high removal efficiency of $\text{NH}_4^+\text{-N}$ in the HRAPs, the TN removal efficiencies were lower (around 65%). This was attributed to the conversion of some $\text{NH}_4^+\text{-N}$ into $\text{NO}_3^-\text{-N}$ and $\text{NO}_2^-\text{-N}$ (e.g. nitrification), which has also been observed in a previous study using the same HRAPs (Arashiro et al., 2019). In terms of the HRT influence, no significant differences in removal efficiencies between 8 d and 4 d were observed. The results of the wastewater treatment efficiency are in accordance with those of previous studies using HRAPs for wastewater treatment (Arashiro et al., 2019; Gutierrez et al., 2016).

The main mechanism of pollutants (e.g., N, P and C) removal by microalgae in HRAP is biomass assimilation, as microalgae can utilize them for the production of cellular components, such as the synthesis of proteins, nucleic acids and carbohydrates (Hoffmann, 1998). Furthermore, the high removal efficiencies of NH_4^+ and TN can also be partially attributed to ammonia volatilization and organic N settlement, respectively (Chen et al., 2006). Likewise, previous studies have demonstrated that precipitation of phosphate with other ions, such as calcium and magnesium in HRAPs also contributes to P removal (Chen et al., 2006; Delgadillo-Mirquez et al., 2016). For organic matter, the decomposition of complex organic carbon compounds by heterotrophic microorganisms in HRAPs (i.e. bacteria and fungi) has been proposed to partially facilitate organic C removal (Mohsenpour et al., 2021).

As far as the Se removal efficiency is concerned, no significant difference between an HRT of 8 days (Se removal average 43%) and 4 days (Se removal average of 46%) was observed (Fig. 6.5). Liu et al. (2019) studied the Se removal efficiency by *Chlorella vulgaris* after exposure to different selenite concentrations in BG11 nutrient medium and found that approximately 51 and 90% of Se was removed upon 500 and 1000-3000 $\mu\text{g/L}$ of Se exposure. This removal was mainly achieved through Se volatilization by facilitating Se methylation by algae upon exposure to highly toxic Se levels (also called Se detoxification mechanism). Besides, Liu et al. (2019) further studied the effect

Chapter 6

of *Chlorella vulgaris* biomass density on selenite removal under 1580 µg/L of Se exposure after 3 days of cultivation and concluded that Se accumulation became the main Se removal mechanism at algal densities between 0.75 and 4.03 g dry weight/L, with an average Se removal of 49–62%, which is close to the Se removal efficiency observed in this study (43–46%). Likewise, it might be deduced that the Se removal in this study was mainly via microalgae Se accumulation, as reflected by the suitable biomass density (around 0.42 g DW/L in the HRAPs) and the lower Se exposure dosage (approximately 25–60 µg Se/L) without toxic effects. Additionally, the Se removal efficiency observed in this study was similar to those reported by Gerhardt et al. (1991), who found an average selenate removal of 45% in high-rate aerobic (algae)–anoxic (anaerobic bacteria) ponds treating agricultural drainage water over two years. The Se removal efficiency by HRAP is not very high, so enhancement of Se accumulation by HRAP is necessary, such as through optimization of growth conditions (e.g. optimal temperature and light), screening specific microalgae species with high Se accumulation ability, conducting pretreatment processes aimed to remove other components (e.g. S and P), which could interfere Se uptake, and inoculation of microalgae with specific bacteria facilitating Se removal.

Chapter 6




Figure 6.4. Influent (●) and effluent (■) concentration of turbidity, total suspended solids (TSS), volatile suspended solids (VSS), total carbon (TC), total and soluble chemical oxygen demand (COD_{tot} and COD_{sol}), total and soluble P (SP and TP) and NH₄⁺-N monitored in the HRAP-Se (with Se spiking, left) and HRAP-C (without Se spiking, right) systems over the experimental period.




Figure 6.5. Influent and effluent concentrations of total Se in the HRAP-Se with continuous selenite spiking during the experimental period. HRT was reduced from 8 to 4 days after 50 days.

6.3.3 Nutritional value of microalgae grown in HRAPs

6.3.3.1 Selenium species in Se-enriched microalgae

Fig. 6.6 shows the chromatogram of Se species in microalgae grown in the HRAP-Se collected at the operational days 43–50. Se-methyl-selenocysteine (SeMetSeCys), Se-methionine (SeMet), Se(IV) and Se(VI) were observed in the sample. 95% of the accumulated Se in the microalgae was converted into organic Se forms. SeMet accounted for the highest proportion (91%) of the identified Se species, whereas the percentage of inorganic Se(IV) and Se(VI) was only 1.9% and 3.0%, respectively. This is consistent with many previous results. For instance, Gómez-Jacinto et al. (2020) found that 95% of the Se taken up by *Chlorella sorokiniana* was transformed into organic Se, and SeMet accounted for 79% of the total Se, when cultivated in Basal medium containing 50 mg/L selenate. Vu et al. (2019) demonstrated that SeMet and SeMetSeCys were predominating Se species in Se-enriched *Chlorella vulgaris* upon selenite (2.25–4.5mg/L) exposure, while Umysova et al. (2009) reported that SeMet made up only 30–40% of the total Se in *Scenedesmus quadricauda* after selenite (10 mg/L) or selenate (20–50 mg/L) exposure.

Chapter 6




Figure 6.6. Chromatograms of Se speciation of (a) Se standard solution containing 100 µg/L of each Se species and (b) an extract of Se-enriched microalgae (diluted 20 times) grown in the HRAP-Se at the operational week 7.

SeMet, a type of selenoamino acid, is one of the major nutritional source of Se for higher animals and humans, as these are unable to synthesize SeMet in their organs (Schrauzer, 2003). Importantly, SeMet is more bioavailable to provide higher Se concentrations in tissues than inorganic Se and is beneficial for human and animal health, which is thus claimed as the most suitable form of Se for nutritional supplementation (Gómez-Jacinto et al., 2020). Our results indicate that microalgae cells are capable of accumulating and transforming less-valuable inorganic Se into

Chapter 6

more-valuable selenoamino acids efficiently. Accordingly, the microalgae enriched with SeMet produced in this study might be a potential and preferable alternative Se source for feed supplementation without utilizing other external nutrients for microalgae growth in domestic wastewater.

6.3.3.2 Selenium bioaccessibility of microalgae

Bioaccessibility measures the fraction of a substance released from products into the gastrointestinal tract by mimicking the gastric and intestinal digestion through *in vitro* tests (Vu et al., 2019). The digestion model in this study comprised a simulation of both the stomach and intestinal physiology of the pig. According to the results, the bioaccessibility of Se in the ball-milled sample was significantly higher than that in the raw sample (Fig. 6.7). This result was expected, as the ball milling would disrupt microalgae cell walls and therefore enhance the Se release from biomass during the gastrointestinal digestion, indicating the importance of pretreatments (i.e. cell disruption) for improving nutrient bioaccessibility. 49 and 63% of the Se in the raw and ball-milled Se-enriched microalgae were solubilized under the gastrointestinal conditions and were thus potentially bioavailable, while the *in vitro* digestibility of Se in the raw and ball-milled microalgae grown in the HRAP-C (control) was 69 and 95%, respectively. The lower digestibility of Se in the Se-enriched microalgae biomass may be attributed to the significantly higher total Se content in the Se-enriched biomass in comparison with the control microalgae, resulting in the incorporation of part of the extra Se in the less digestible microalgae fraction, such as in the hemicellulosic cell wall structure (Gómez-Jacinto et al., 2020).

A similar Se bioaccessibility (~49%) was found in Se-enriched *Chlorella vulgaris* (Vu et al., 2019), while it should be noted that Se bioaccessibility in Se-rich yeast, the most popular supplement of organic Se, is generally higher in comparison with that of the HRAP-Se grown microalgae. For instance, Lavu et al. (2016) reported that approximately 70% of total Se was bioaccessible in SelenoPrecise tablet (Se-rich yeast); Reyes et al. (2006) observed that 89% of total Se in SEAS 6 (Se-rich yeast, Pharma Nord, Denmark) was extracted after gastrointestinal digestion; Thiry et al. (2013) demonstrated that 57% of Se from Se-yeast (LepiVits, Vincennes, France) was liberated through the *in vitro* gastrointestinal digestion. Moreover, a Se bioaccessibility of 81% was previously observed in the Se-enriched microalga *Chlorella sorokiniana*

Chapter 6

(Gómez-Jacinto et al., 2020), which is higher than that in the current study. The higher Se bioaccessibility observed in the above studies might be due to the different digestion methods (e.g. different amount and type of enzymes). Specifically, extra bile salts were adopted in the intestinal juice of the all studies referred to above, while they were not added in the current study. Bile salts assist in the solubilisation of fat, which may result in the dissolution of lipid-bound Se and thus a higher Se bioaccessibility. More enzymes (e.g., extra addition of amylase) and chemicals (e.g., NaCl) were also included in the intestinal juice used by Gómez-Jacinto et al. (2020) and Reyes et al. (2006) to simulate human gastrointestinal digestion in comparison with the juice used in this study. Taking into account that animals could also excrete bile and amylase, further assessment of Se bioaccessibility involving also bile salts and amylase is proposed for the Se-enriched microalgae tested in the current study. Besides, in this study, phosphotungstic acid was added into the extract after gastrointestinal digestion to precipitate undigested protein or peptide, which could correspondingly result in the precipitation of protein/peptide/macromolecule-bound Se and eventually lead to a lower Se bioaccessibility. Also results reported by Reyes et al. (2006) point in this direction, as they observed that 89% of total Se in SEAS 6 was extracted after gastrointestinal digestion, but only 34% of the extracted Se was free SeMet and most of the Se-species seemed to be peptide-bound. Additionally, the difference in microalgae species, Se concentrations or species in the growth medium and microalgae biomass could partially contribute to differences in results between this study and the study of Gómez-Jacinto et al. (2020).




Figure 6.7. Bioaccessibility of Se in the raw and ball-milled microalgae grown in both the HRAP-C (Control) and HRAP-Se (Se-enriched microalgae). Values are mean $\pm$ standard deviation (n=3).

6.3.3.3 Protein extraction and macromolecular characterization of microalgae

The total crude Kjeldahl-protein (TKN) content of the microalgae grown in both HRAPs was about 48% (Table 6.1), which is within the range reported in the literature for microalgae species (Arashiro et al., 2019; Rasouli et al., 2018). This is comparable to that of soybean (38% in full-fat soybeans, 48% for dehulled soybean meal and 44% for non-dehulled soybean meal) (Moheimani et al., 2018), which is currently the primary source of protein for pigs around the world (Moheimani et al., 2018). Additionally, a slightly higher content of carbohydrates and lipids was observed for the biomass grown in the HRAP-Se than that grown in the HRAP-C. Specifically, the biomass grown in the HRAP-C and HRAP-Se was composed of 21% and 32% carbohydrates, and 19% and 21% lipids, respectively, indicating that Se may have the potential to stimulate the biosynthesis of these compounds in microalgae, which is consistent with the results of Gan et al. (2019). The increased carbohydrate levels in microalgae grown in HRAP-Se may be attributed to the enhancement of photosynthesis. Indeed, several studies have demonstrated that Se has the ability to improve the photosynthetic performance of microalgae, as shown by the increase of photosynthetic pigments content (e.g., carotenoid and chlorophyll) and the enhancement of electron transport system activity (Zhong & Cheng, 2017). Similarly, Se may stimulate the enzyme activity of fatty acid synthesis (FAS) in microalgae cells, eventually resulting in higher lipid production in

Chapter 6

HRAP-Se. This stimulation effect of Se to FAS has also been observed in yeast cells (Kieliszek et al., 2019).

For protein extraction, the protein content obtained by SDS extraction was much higher than the protein content obtained by DI water extraction (Table 6.1). The lowest protein content (12–14% for SDS extraction) was observed for the biomass after cell disruption by freeze-thawing, while ball and bead milling of the biomass favored the highest release of protein, i.e. 46–48% upon SDS extraction. This result indicates that ball and bead milling disruption in combination with SDS extraction results is the most efficient solubilization and quantification of proteins in microalgae by the Lowry method, which could provide a reference for protein extraction of microalgae.

Table 6.1. Protein content of microalgae subjected to different cell disruption techniques and biochemical composition (%) of microalgae grown in the HRAPs. Results are reported as percentage of the total volatile suspended solids (VSS). Values are mean $\pm$ standard deviation (n=3).

		HRAP-C		HRAP-Se	
		H ₂ O- Lowry ^a	SDS- Lowry ^b	H ₂ O- Lowry ^a	SDS- Lowry ^b
Protein content after application of different cell disruption techniques	Freeze-thawing	3.2 $\pm$ 0.0	14 $\pm$ 0.7	2.4 $\pm$ 0.1	12 $\pm$ 2.5
	Sonication	9.4 $\pm$ 0.7	18 $\pm$ 1.4	10 $\pm$ 0.3	16 $\pm$ 0.8
	High pressure cell disruption	7.5 $\pm$ 0.5	24 $\pm$ 5.7	10 $\pm$ 0.4	32 $\pm$ 1.7
	Ball milling	4.5 $\pm$ 0.3	46 $\pm$ 1.8	5.9 $\pm$ 0.4	48 $\pm$ 1.2
	Bead milling	10 $\pm$ 2.3	47 $\pm$ 4.8	7.6 $\pm$ 1.9	48 $\pm$ 6.3
Macromolecular composition	Kjeldahl-protein ^c	47.6		48.4	
	Carbohydrates	20.5		31.9	
	Lipids	18.7		20.9	

Note: ^a Disrupted microalgae cell suspension was incubated at 100 °C for 5 min with DI water followed by Lowry protein measurement.

^b Disrupted microalgae cell suspension was incubated at 100 °C for 5 min with SDS solution followed by Lowry protein measurement.

^c Total protein content was calculated by multiplying the total Kjeldahl nitrogen by 5.95.

Chapter 6

6.3.3.4 Amino acids in the microalgae

Table 6.2 compares the amino acid content of microalgae grown in both HRAPs with that of soybeans and soybean meal. The amino acid content of the microalgae grown in both HRAPs was close to that of soybeans (with the exception of glutamic acid), while it was slightly lower than that in soybean meal, except for glycine, threonine, and alanine contents which were higher in the microalgae (Table 6.2). This result showed that microalgae could be a source of some essential amino acids for animals, such as lysine, threonine, methionine, cystine, isoleucine, histidine, valine, arginine, phenylalanine and tyrosine, which must be provided in some animals' diets (Mahan and Shields, 1998). Even though the content of some essential amino acids (e.g., arginine, lysine and cystine) in the Se-enriched microalgae was slightly lower than that in the soybean, the result still shows the potential of using the produced microalgae as feed additive in animal diets, offering a valid alternative taking into account the higher land, water, nutrient and carbon footprint of conventional vegetable protein production (Matassa et al., 2016). However, the Se content in the Se-enriched microalgae should be particularly addressed when using it as a feed additive because European Food Safety Authority regulated that the total Se content in the complete feed can be maximum 0.5 mg Se/kg (EFSA, 2016). Besides, further study should quantify the digestibility of amino acids in the microalgae, as it is also an important factor affecting amino acids utilization efficiency.

Chapter 6

Table 6.2. Amino acid contents of microalgae grown in HRAPs at day 50 compared with soybeans and soybean meal (SBM) for pigs. Values are mean $\pm$ standard deviation (n=3)

Amino acid	HRAP-C	HRAP-Se	Soybeans	Soybean meal
	[g/100g DW]			
Aspartic acid	3.11 $\pm$ 0.05	2.57 $\pm$ 0.03	3.89	4.88
Glutamic acid	3.65 $\pm$ 0.09	3.06 $\pm$ 0.04	6.05	7.87
Asparagine	<i>N.D.</i>	<i>N.D.</i>	--	--
Serine	1.48 $\pm$ 0.03	1.25 $\pm$ 0.01	1.67	2.14
Glutamine	0.15 $\pm$ 0.01	<i>N.D.</i>	--	--
Histidine	0.60 $\pm$ 0.00	0.56 $\pm$ 0.01	0.88	1.26
Glycine	2.05 $\pm$ 0.01	1.60 $\pm$ 0.03	1.52	1.89
Threonine	1.78 $\pm$ 0.04	1.49 $\pm$ 0.02	1.42	1.76
Citrulline	<i>N.D.</i>	<i>N.D.</i>	--	--
Arginine	1.91 $\pm$ 0.04	1.74 $\pm$ 0.03	2.45	3.17
Alanine	3.03 $\pm$ 0.05	2.15 $\pm$ 0.03	1.59	1.92
Tyrosine	1.25 $\pm$ 0.02	1.02 $\pm$ 0.02	1.20	1.55
Valine	1.81 $\pm$ 0.04	1.39 $\pm$ 0.04	1.73	1.93
Methionine	0.60 $\pm$ 0.01	0.51 $\pm$ 0.02	0.55	0.60
Phenylalanine	1.68 $\pm$ 0.03	1.38 $\pm$ 0.02	1.74	2.26
Isoleucine	1.45 $\pm$ 0.03	1.18 $\pm$ 0.04	1.60	1.96
Ornithine	0.19 $\pm$ 0.00	<i>N.D.</i>	--	--
Leucine	2.79 $\pm$ 0.06	2.18 $\pm$ 0.04	--	--
Lysine	1.76 $\pm$ 0.01	1.49 $\pm$ 0.02	2.23	2.76
Hydroxyproline	<i>N.D.</i>	<i>N.D.</i>	--	--
Proline	1.66 $\pm$ 0.02	1.54 $\pm$ 0.01	--	--
Cysteic acid	0.25 $\pm$ 0.00	0.25 $\pm$ 0.00	0.59	0.68
Total	31.24 $\pm$ 0.52	25.76 $\pm$ 0.41	--	--

Data of soybeans and soybean meal from Moheimani et al. (2018).

N.D.= not detected

-- = no data shown

6.3.3.5 Fatty acids in the microalgae

Chapter 6

The composition and contents of total saturated fatty acids (SFA), monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA) in the microalgae are shown in Table 6.3. The detailed composition and contents are listed in Table 6.S3. The data to some extent indicated that microalgae grown in the HRAP-C contained a higher percentage of total SFA and MUFA in comparison with those present in the HRAP-Se, while it had a lower percentage of PUFA omega-3 (ω 3) and omega-6 (ω 6) (Table 6.3). Specifically, SFA and MUFA accounted for 32.1 and 10.6% of the fatty acids for the biomass grown in HRAP-C, respectively, compared to 25.0 and 8.17% for biomass grown in the HRAP-Se, respectively. The percentage of PUFA ω 6 and ω 3 were 10.0 and 17.1% for the biomass grown in the HRAP-C, and 11.8 and 26.1% for the biomass grown in the HRAP-Se, respectively. Among PUFA ω 3, eicosapentaenoic (EPA, C20:5) showed the most significant increase in concentration when Se was added to the growth medium (13.2% in versus 24.7% of the fatty acids in HRAP-C and HRAP-Se, respectively) (Table 6.S3). Although further research is still needed to confirm these results because of the fewer sample points we collected, this may to some extent indicate that Se has the potential of contributing to the synthesis of PUFAs and the production of value-added biomass. Moreover, previous studies have also shown the ability of Se to activate fatty acid unsaturation and stimulate the formation of PUFA via inducing desaturase activity, such as Δ 12 and Δ 15 (Čertík et al., 2013; Kieliszek et al., 2019). Likewise, the increased percentage of PUFAs in microalgae cultivated in HRAP-Se of the current study could be due to the fact that Se may be involved in the induction of fatty acid Δ 12, Δ 15 and Δ 5 desaturases. Noticeably, the elevated PUFA ω 3 percentage of microalgae in the HRAP-Se, particularly for the EPA, could demonstrate this, as Δ 15 and Δ 5 are the main desaturases to catalyze PUFA ω 3 and EPA formation (Ma et al., 2016).

PUFA ω 3 and ω 6 are considered essential fatty acids, being beneficial for human health and livestock nourishment (Moheimani et al., 2018). They have a positive effect on cardio-circulatory diseases, atherosclerosis, coronary disease, degenerative diseases and anticancer (Otleş & Pire, 2001). Besides, the proportion of the PUFA ω 6 and ω 3 of the biomass in this study is also higher than that of microalgae grown on anaerobically digested piggery effluent (8.7% for ω 6 and 15.7% for ω 3) (Moheimani et al., 2018). Besides, among the different PUFAs ω 3 present in algae, eicosapentaenoic (EPA, C20:5) has the most important nutritional and health value (Becker, 2013). Its

Chapter 6

supplementation can be co-therapeutic (Doughman et al., 2007). In this study, EPA was dominant in the biomass grown in both the HRAP-C and HRAP-Se, accounting for 13.2 and 24.7% of the fatty acids, respectively (Table 6.S3). The EPA proportion of the biomass grown in the HRAP-Se is higher than that of commercial products on the market, such as salmon (14% EPA) and fish (18% EPA) oil (Otleş & Pire, 2001), which is considered favorable for animal and human nutrition. The value-added biomass produced from HRAPs spiked with Se could thus offer a promising alternative source of valuable PUFAs.

Table 6.3. Fatty acid composition and content of the microalgae grown in the control (HRAP-C) and Se spiked microalgae pond (HRAP-Se). SFA: saturated fatty acids; MUFA: monounsaturated fatty acids; PUFA: polyunsaturated fatty acids; ω 3 and 6: omega-3 and 6. Values are mean $\pm$ standard deviation (n=3).

	HRAP-C		HRAP-Se	
	[% relative fat]	[mg/100 g]	[% relative fat]	[mg/100 g]
Total SFA	32.1 $\pm$ 0.1	1359 $\pm$ 7.9	25.0 $\pm$ 0.1	2215 $\pm$ 4.4
Total MUFA	10.6 $\pm$ 0.1	450 $\pm$ 4.1	8.17 $\pm$ 0.0	773 $\pm$ 3.1
Total PUFA ω -6	10.0 $\pm$ 0.0	426 $\pm$ 2.8	11.8 $\pm$ 0.0	1048 $\pm$ 10.0
Total PUFA ω -3	17.1 $\pm$ 0.1	726 $\pm$ 4.1	26.1 $\pm$ 0.1	2310 $\pm$ 12.0

6.3.4 Pathogenic bacterial content

Microalgae grown on wastewater can harbor a risk of pathogen transfer when they are consumed as feed. Therefore, for the effective protection of human and animal health, microbiological regulations must be met (Montville & Matthews, 2008). Table 6.4 shows the content of selected manure-borne bacteria of fresh and dried HRAP grown biomass. A concentration of over 10^7 CFU/g of aerobic bacteria was found in the fresh microalgae samples, however, these were absent in the dried biomass (both oven-dried and freeze-dried). The coliform, *E. coli* (at a concentration $> 10^5$ CFU/g) and *Salmonella* were detected in the fresh microalgae samples grown in the HRAP-C, while the population of *Listeria* was below 25 CFU/g (tolerance level < 100 CFU/g (Montville & Matthews, 2008) in all types of samples. Table 6.4 indicates that further downstream processing after harvest, such as drying, could reduce the bacterial loads of microalgae

Chapter 6

and avoid the pathogen risk, supporting the application of microalgae grown on domestic wastewater as a potential feed supplement. This is similar to the finding of Moheimani et al. (2018), who studied the pathogen loads of dried microalgae cultivated on anaerobically digested piggery effluent. However, further risk assessment on the implementation of Se-enriched microalgae grown in domestic wastewater as feed supplement is still required, such as *in vivo* studies and quantification of other parameters potentially affecting food and feed safety (e.g., residues of mycotoxins, antibiotics, and nucleic acids).

Table 6.4. Pathogen loads on fresh (CFU/g wet weight) and dry (CFU/g dry weight) microalgae biomass grown in the control (HRAP-C) and Se spiked (HRAP-Se) microalgae ponds treating domestic wastewater.

		Aerobic bacteria	Coliforms	<i>E. coli</i>	<i>Listeria</i>	<i>Salmonella</i>
		[CFU/g]				
Fresh	HRAP-Se	2.35*10 ⁷	2.60*10 ⁶	N.D.	< 25	Absence
biomass	HRAP-C	9.93*10 ⁷	4.52*10 ⁷	2*10 ⁵	< 25	Presence
Oven-dry	HRAP-Se	N.D.	N.D.	N.D.	< 25	Absence
biomass	HRAP-C	N.D.	N.D.	N.D.	< 25	Absence
Freeze-dry	HRAP-Se	N.D.	N.D.	N.D.	< 25	Absence
biomass	HRAP-C	N.D.	N.D.	N.D.	< 25	Absence

N.D.= not detected

6.4 Conclusions

This study investigated the potential of microalgae to remove selenite and selenate from domestic wastewater and to recover and upgrade low-value resources into high-value products in HRAPs systems. This study clearly highlighted that HRAPs-grown microalgae are good candidates to upgrade nutrients and carbon dioxide into Se-enriched microalgae biomass that can be used as valuable feed supplements. Microalgae mainly accumulate organic SeMet (91%) in their cells after taking up inorganic Se from the solution, indicating the ability of microalgae to transform inorganic Se into selenoamino acids. Se release upon gastrointestinal digestion of

Chapter 6

microalgae was quite high and additional processing of the microalgae through ball-milling treatment further improved the digestibility significantly. Se may potentially assist the production of lipids and carbohydrates by microalgae, but further research with more analysis is still needed to confirm this. The nutritional properties of Se-enriched microalgae were comparable to commercially available soybean meals in terms of protein and amino acids content, while the fatty acid content of Se-enriched microalgae surpassed that of high-quality commercial fish oil. However, the Se bioaccessibility in the microalgae may still be further improved, e.g., by further testing and optimization of pretreatments. Moreover, standardization of assessment methods for bioaccessibility may enable more accurate comparison of bioaccessibility data reported by different researchers.

Chapter 6

Supplementary Information




Figure 6.S1. Se removal efficiency of microalgae grown in the batch system after 7d cultivation

Chapter 6

Table 6.S1. Summary of the main parameters (average $\pm$ standard deviation) monitored in the primary effluent (the influent of HRAPs) and the effluent of HRAPs throughout the entire experimental period.

	Primary effluent (=influent of HRAPs)	Effluent of HRAP-Se	Effluent of HRAP-C
pH	7.8 $\pm$ 0.2	8.7 $\pm$ 0.6	8.7 $\pm$ 0.5
Turbidity (NTU)	271 $\pm$ 241.9	14 $\pm$ 20.0	12 $\pm$ 16.0
TSS (mg/L)	424 $\pm$ 341.7	37 $\pm$ 48.0	22 $\pm$ 18.8
VSS (mg/L)	345 $\pm$ 254.6	35 $\pm$ 42.3	21 $\pm$ 15.1
COD _{tot} (mg/L)	497 $\pm$ 279.3	123 $\pm$ 97.2	117 $\pm$ 50.1
COD _{sol} (mg/L)	223 $\pm$ 139.7	97 $\pm$ 63.1	99 $\pm$ 54.6
TC (mg/L)	331 $\pm$ 176.2	97 $\pm$ 57.6	90 $\pm$ 39.0
TP (mg/L)	10 $\pm$ 5.5	2.9 $\pm$ 2.9	2.3 $\pm$ 1.6
SP (mg/L)	5.3 $\pm$ 2.7	1.6 $\pm$ 1.6	1.6 $\pm$ 1.2
TN (mg/L)	64 $\pm$ 19.8	26 $\pm$ 28.2	18 $\pm$ 13.1
NH ₄ ⁺ -N (mg/L)	27 $\pm$ 2.9	2.9 $\pm$ 2.8	2.4 $\pm$ 2.6
NO ₃ ⁻ -N (mg/L)	0.4 $\pm$ 0.9	1.3 $\pm$ 3.6	1.8 $\pm$ 6
NO ₂ ⁻ -N (mg/L)	0.8 $\pm$ 2.0	4.6 $\pm$ 3.7	5.8 $\pm$ 5.1
SO ₄ ²⁻ -S (mg/L)	55 $\pm$ 55.4	47 $\pm$ 17.4	49 $\pm$ 17.6

Chapter 6

Table 6.S2. Summary of the removal efficiency (%) of the main water quality parameters measured in influent and effluent of the two HRAPs (HRAP-Se and HRAP-C) (average $\pm$ standard deviation) for the entire experimental period as well as the two subperiods with a HRT of 8 days and 4 days, respectively.

	Entire experimental period		HRT= 8d		HRT= 4d	
	HRAP-Se	HRAP-C	HRAP-Se	HRAP-C	HRAP-Se	HRAP-C
Turbidity	91 $\pm$ 9.6	90 $\pm$ 19.7	92 $\pm$ 9.0	89 $\pm$ 24.2	89 $\pm$ 10.3	91 $\pm$ 12.4
TSS	86 $\pm$ 16.0	90 $\pm$ 14.1	94 $\pm$ 3.7	96 $\pm$ 2.5	80 $\pm$ 14.1	87 $\pm$ 6.4
VSS	88 $\pm$ 12.1	92 $\pm$ 6.9	93 $\pm$ 3.9	95 $\pm$ 3.0	82 $\pm$ 14.7	90 $\pm$ 8.4
COD _{tot}	70 $\pm$ 20.4	66 $\pm$ 23.1	74 $\pm$ 23.4	70 $\pm$ 22.0	64 $\pm$ 17.1	63 $\pm$ 24.7
COD _{sol}	49 $\pm$ 23.9	47 $\pm$ 23.7	46 $\pm$ 23.6	40 $\pm$ 26.0	52 $\pm$ 25.1	53 $\pm$ 20.9
TC	67 $\pm$ 17.6	65 $\pm$ 20.2	73 $\pm$ 13.9	73 $\pm$ 15.4	60 $\pm$ 19.7	59 $\pm$ 22.6
TP	77 $\pm$ 18.1	72 $\pm$ 25.5	84 $\pm$ 12.6	65 $\pm$ 20.4	70 $\pm$ 20.9	67 $\pm$ 29.2
SP	71 $\pm$ 21.6	72 $\pm$ 18.8	75 $\pm$ 21.8	77 $\pm$ 16.5	68 $\pm$ 22.1	69 $\pm$ 20.4
TN	65 $\pm$ 24.8	67 $\pm$ 27.3	76 $\pm$ 12.0	70 $\pm$ 21.9	53 $\pm$ 29.8	64 $\pm$ 32.6
NH ₄ ⁺ -N	93 $\pm$ 6.2	92 $\pm$ 7.7	94 $\pm$ 3.6	92 $\pm$ 9.3	91 $\pm$ 8.9	93 $\pm$ 6.2
Total Se	44 $\pm$ 6.5	N.D.	43 $\pm$ 7.3	N.D.	46 $\pm$ 4.3	N.D.

N.D.: Not determined

Chapter 6

Table 6.S3. Fatty acid composition and content of the microalgae grown in HRAPs the control (HRAP-C) and Se spiked microalgae ponds (HRAP-Se). Values are mean $\pm$ standard deviation (n=3).

	HRAP-C		HRAP-Se	
	[% relative fat]	[mg/100 g]	[% relative fat]	[mg/100 g]
C10:0	0.62 $\pm$ 0.00	26.4 $\pm$ 0.06	0.91 $\pm$ 0.14	80.5 $\pm$ 11.51
C12:0	0.77 $\pm$ 0.00	32.7 $\pm$ 0.43	0.44 $\pm$ 0.00	38.6 $\pm$ 0.35
C14:0	4.48 $\pm$ 0.00	190 $\pm$ 1.18	4.44 $\pm$ 0.00	393 $\pm$ 2.50
C15:0	0.60 $\pm$ 0.00	25.7 $\pm$ 0.18	0.32 $\pm$ 0.01	28.7 $\pm$ 0.65
C16:0	21.4 $\pm$ 0.10	908 $\pm$ 4.81	17.2 $\pm$ 0.03	1524 $\pm$ 13.44
C17:0	0.80 $\pm$ 0.01	33.8 $\pm$ 0.45	0.25 $\pm$ 0.01	22.2 $\pm$ 0.65
C18:0	2.34 $\pm$ 0.02	99.4 $\pm$ 1.12	1.05 $\pm$ 0.02	92.9 $\pm$ 1.02
C20:0	0.23 $\pm$ 0.02	9.66 $\pm$ 0.89	0.09 $\pm$ 0.00	8.06 $\pm$ 0.06
C22:0	0.38 $\pm$ 0.00	15.9 $\pm$ 0.04	0.15 $\pm$ 0.00	13.2 $\pm$ 0.71
C24:00	0.42 $\pm$ 0.03	17.9 $\pm$ 1.03	0.16 $\pm$ 0.01	13.9 $\pm$ 0.71
Total SFA	32.1 $\pm$ 0.10	1359 $\pm$ 7.94	25.0 $\pm$ 0.13	2215 $\pm$ 4.38
C14:1	0.08 $\pm$ 0.01	3.27 $\pm$ 0.32	0.07 $\pm$ 0.00	6.10 $\pm$ 0.39
C17:1	0.25 $\pm$ 0.00	10.7 $\pm$ 0.02	0.28 $\pm$ 0.01	25.2 $\pm$ 0.77
c9C18:1	6.40 $\pm$ 0.05	271 $\pm$ 2.85	6.25 $\pm$ 0.00	553 $\pm$ 4.14
c11C18:1	3.74 $\pm$ 0.03	159 $\pm$ 1.75	1.52 $\pm$ 0.01	134 $\pm$ 0.01
C20:1	0.15 $\pm$ 0.01	6.19 $\pm$ 0.23	N.D.	N.D.
C24:1	N.D.	N.D.	0.05 $\pm$ 0.00	4.52 $\pm$ 0.04
Total MUFA	10.6 $\pm$ 0.07	450 $\pm$ 4.08	8.17 $\pm$ 0.03	773 $\pm$ 3.01
C18:2 ω -6	6.5 $\pm$ 0.02	276 $\pm$ 1.70	6.60 $\pm$ 0.02	584 $\pm$ 5.81
C18:3 ω -6	0.34 $\pm$ 0.01	14.3 $\pm$ 0.37	0.50 $\pm$ 0.00	43.8 $\pm$ 0.56
C20:2 ω -6	0.18 $\pm$ 0.01	7.62 $\pm$ 0.32	0.07 $\pm$ 0.00	6.23 $\pm$ 0.17
C20:3 ω -6	0.18 $\pm$ 0.00	7.56 $\pm$ 0.21	0.21 $\pm$ 0.00	18.6 $\pm$ 0.58
C20:4 ω -6	2.74 $\pm$ 0.01	116 $\pm$ 0.91	4.42 $\pm$ 0.00	391 $\pm$ 3.15
C22:5 ω -6	0.08 $\pm$ 0.00	3.38 $\pm$ 0.04	0.05 $\pm$ 0.00	4.77 $\pm$ 0.04
Total PUFA ω -6	10.0 $\pm$ 0.04	426 $\pm$ 2.82	11.8 $\pm$ 0.03	1048 $\pm$ 10.32
C18:3 ω -3	3.62 $\pm$ 0.02	153 $\pm$ 1.32	1.23 $\pm$ 0.05	109 $\pm$ 3.46
C20:4 ω -3	0.12 $\pm$ 0.00	5.20 $\pm$ 0.02	0.12 $\pm$ 0.02	10.4 $\pm$ 1.75
C20:5 ω -3	13.2 $\pm$ 0.03	558 $\pm$ 2.83	24.7 $\pm$ 0.03	2182 $\pm$ 13.72
C22:5 ω -3	0.09 $\pm$ 0.01	3.81 $\pm$ 0.22	0.04 $\pm$ 0.00	3.29 $\pm$ 0.20
C22:6 ω -3	0.13 $\pm$ 0.01	5.63 $\pm$ 0.24	0.06 $\pm$ 0.00	5.52 $\pm$ 0.15
Total PUFA ω -3	17.1 $\pm$ 0.05	726 $\pm$ 4.15	26.1 $\pm$ 0.06	2310 $\pm$ 11.97

N.D.= not detected

Chapter 7 Production of Se-enriched microalgae in raceway ponds treating domestic wastewater as biostimulant and biofertilizer

This chapter was redrafted after:

Li, J., Ferrer, I., Lens, P.N.L., Du Laing, G., 2021. Evaluation of selenium-enriched microalgae produced on domestic wastewater as biostimulant and biofertilizer for growth of selenium-enriched crops. *Journal of Applied Phycology* (In press).

Abstract

This study assessed selenium (Se)-enriched microalgae biomass produced in a pilot-scale raceway pond treating domestic wastewater and an extract thereof as biostimulant and biofertilizer. After producing the Se-enriched microalgae in a raceway pond treating domestic wastewater, the effect of Se-enriched microalgae biomass and an extract thereof on seed germination, growth and yield of beans (*Phaseolus vulgaris* L.) was studied by conducting a germination test as well as foliar and soil applications in pot experiments. Furthermore, the potential of the Se-enriched microalgae dry biomass and extract to elevate the Se concentration of beans, leading to a biofortified crop, was also assessed in a pot experiment. Presoaking seeds in the Se-enriched microalgae extract at low concentration (1%) enhanced their germination, as measured by the significant increase of seedling length and vigor index. Application of the Se-enriched microalgae extract as foliar spray was more effective in stimulating the growth of beans and increasing the Se concentration in the seeds compared to its application as soil drench. Foliar spray resulted in a 3.5 times increase of the dry biomass of the seeds (at 1%) and 1.8 times Se increment in the seeds (at 5%). Additionally, amendment of the soil with Se-enriched microalgae biomass (at 5%) enhanced the growth of beans (3.2 times for seeds) and Se concentration in the bean plants (1.8 times for seeds), simultaneously. These results indicate that microalgae cultivated in Se-rich wastewater could be used as a microalgae-based biofertilizer or biostimulant to improve the bean seeds yield as well as the Se content in the beans, leading to beans with a higher market value. This may also offer an environmental-friendly and sustainable approach to biofortify food crops in Se-deficient regions.

Keywords: Algae, Fertilizer, Biofortification, Beans, Selenium, Wastewater

7.1 Introduction

Se deficiency exists worldwide, resulting in negative health effects and even causing Se-deficiency diseases, e.g. endemic Keshan disease in China (Tan et al., 2002; Wu et al., 2015). It is estimated that over 1 billion people may consume less Se than required for optimal protection against cancer and cardiovascular disease (Haug et al., 2007). The low dietary Se intake is generally associated with the consumption of food containing a low Se content, usually due to the low Se concentration in the soils on which the crops are grown. Biofortification is the possible solution for Se deficiency (Boldrin et al., 2013; Li et al., 2020a). However, the adverse effects of applying conventional inorganic Se fertilizers on soils and the environment is leading to the exploration of alternative Se fertilizers.

Microalgae can be cultivated in wastewater and agricultural runoff, recovering excess nutrients, including Se, while reclaiming the wastewater (Gan et al., 2019; Garcia-Gonzalez & Sommerfeld, 2016). The generated microalgae biomass with high nutrient content is not only considered as a valuable ingredient for food and animal feed, but has also a potential as biofertilizers or biostimulants (Ronga et al., 2019). Nowadays, the use of microalgae in agricultural production as biofertilizer or biostimulant is attracting the interest of growers and agrochemical industries aiming to improve the sustainability of crop production (Calvo et al., 2014; Grzesik & Romanowska-Duda, 2014; Ronga et al., 2019). Biostimulants are substance that are able to enhance physiological processes and stimulate the growth of plants under both optimal and stressful conditions after being applied to the plants or soils. Biofertilizers are bioproducts that deliver nutrients for plant growth, thus stimulating plant growth (Ronga et al., 2019).

Microalgae biomass contains several plant growth-promoting substances, such as phytohormones, vitamins, carotenoids, amino acids, and antifungal substances (Coppens et al., 2015), which could serve as potential biostimulant. A few studies have established an association between greater crop yields and the application of microalgal cellular extracts as biostimulant or microalgae biomass as biofertilizer, respectively. For instance, the application of 1.5 L/ha of the *Spirulina* extract obtained by supercritical fluid extraction on the field has been found to significantly raise the number of grains in ear and shank length of wheat (variety *Akteur*) (Michalak et al.,

Chapter 7

2016). The addition of the microalga *Chlorella vulgaris* biomass to soil (2–3 g dry algae/kg soil) significantly increased the fresh and dry weight of lettuce (Faheed & Abd-El Fattah, 2008). Similarly, the microalga *Acutodesmus dimorphus* dry biomass and its cellular extracts could trigger faster germination and enhance the plant growth and floral production of Roma tomato (Garcia-Gonzalez & Sommerfeld, 2016). The use of microalgae biomass as a slow-release biofertilizer resulted in a higher quality of tomatoes with increased carotenoid and sugar levels (Coppens et al., 2015).

When producing Se-enriched microalgae during wastewater treatment, the produced microalgae may not only be used as biofertilizer or biostimulant to improve crop growth, but also to biofortify the crops by enriching them with Se. Therefore, we explored whether Se-enriched microalgae biomass produced in raceway ponds from domestic wastewater and an extract thereof can be potentially used as an organic nutrient-rich biofertilizer or biostimulant in biofortification strategies to enhance the crop yield, meanwhile improving also the Se levels in the crops. This approach may be beneficial to save Se resources and avoid the introduction of chemicals contamination into the environment in biofortification schemes through the replacement of synthetic chemical fertilizers by Se-enriched microalgae biomass or its extract.

To this end, the influence of the produced Se-enriched microalgae dry biomass and an extract thereof on the germination, growth and yield of green beans (*Phaseolus vulgaris* L.) was assessed, along with the Se concentration in the beans. This is the first study to assess the application of Se-enriched microalgae biomass and an extract thereof as Se biofertilizer and biostimulant to biofortify green beans.

7.2 Materials and methods

7.2.1 Se-enriched microalgae production in a pilot-scale raceway pond

Se-enriched microalgae were produced as described in our previous study (Chapter 6). The microalgal biomass was dominated by *Chlorella* sp.. The Se-enriched biomass collected from the secondary clarifier was thickened by centrifugation and washed twice with deionized (DI) water. The centrifuged paste was frozen at -80 °C overnight, lyophilized and stored at -20 °C for subsequent experiments.

Chapter 7

Qualitative microscopic evaluation of the biomass indicated that the microalgae biomass was dominated by *Chlorella* sp. (Fig. 7.S1, Supplementary Information), which has been previously reported in studies using the same ponds during the same period of the year (Arashiro et al., 2019; Gutierrez et al., 2016).

7.2.2 Preparation of liquid Se-enriched microalgae extracts

Freeze-dried biomass (15 g) was ground by ball-milling (MM 400, Retsch, Haan, Germany) for 10 min at 30 Hz. The ground biomass was suspended into 90 mL DI water and 10 ml 10% sodium dodecyl sulfate (SDS). The final SDS concentration in the extract was 1%. SDS has a significant effect on improving the microalgae extraction efficiency as shown in the previous experiment (Chapter 6). The suspension was stirred on a stirring plate for 10 min to allow the biomass to dissociate, and then incubated at 100 °C for 5 min to obtain the extract. The hot extract was cooled down at room temperature and centrifuged at 5000 rpm for 10 min to separate the cell extracts from the biomass residue. The microalgae extract was stored at 4 °C for further trials. The composition of the Se-enriched microalgae biomass and the extract is shown in Table 7.1.

7.2.3 Germination test with Se-enriched microalgae extract

Bean seeds (*Phaseolus vulgaris*) with uniform shape, size and weight were selected for the germination test. The germination test was conducted as described previously (Garcia-Gonzalez & Sommerfeld, 2016; Hernández-Herrera et al., 2013) with minor modifications. Each treatment (Table 7.2) was replicated three times with 15 seeds per replicate. The seeds were surface-sterilized with 4 % sodium hypochlorite solution for 10 min and subsequently rinsed twice with DI water prior to soaking in 10 mL of the different concentrations of Se-enriched microalgae cell extracts (Table 7.2) for 24 h. After the 24 h soaking, the seeds were placed on 42.5 mm Whatman no. 1 filter papers and then allowed to dry for 12 h at room temperature. The treated seeds were then transferred into 100 mm Petri plates containing moist filter paper (5 mL of DI water) and incubated in an incubator at 27 °C. The filter paper was kept moist by the regular addition of DI water. Seeds germination was counted daily for one week. The seedling length was monitored every other day.

Chapter 7

Table 7.1. Characteristics of the Se-enriched microalgae biomass and extract (average $\pm$ standard deviation, n=3).

	Biomass	Extracts
pH	N.D.	5.71
EC (mS/cm)	N.D.	6.10
Elements	[mg/kg dry matter]	[mg/L]
Se	29 $\pm$ 0.6	0.67 $\pm$ 0.0
P	13935 $\pm$ 323.5	445 $\pm$ 5.5
S	6995 $\pm$ 148.4	378 $\pm$ 4.3
Zn	520 $\pm$ 5.1	1.45 $\pm$ 0.0
Cu	145 $\pm$ 3.0	1.36 $\pm$ 0.0
Ca	51215 $\pm$ 1478.3	280 $\pm$ 4.8
Mg	8199 $\pm$ 175.2	230 $\pm$ 4.0
Na	1640 $\pm$ 56.7	616 $\pm$ 15.0
K	4799 $\pm$ 195.5	476 $\pm$ 9.5
Ni	15 $\pm$ 0.5	1.01 $\pm$ 0.0
Cr	31 $\pm$ 3.0	<LOQ
Cd	<LOQ	<LOQ
Pb	19 $\pm$ 0.3	<LOQ
Hg	0.32 $\pm$ 0.1	<LOQ
As	4.4 $\pm$ 0.3	N.D.
Co	<LOQ	<LOQ

N.D.: Not determined

< LOQ: values lower than the limited of quantification

Chapter 7

Table 7.2. Concentration of Se-enriched microalgae extracts in each treatment.

Treatment	Concentration (%)	Preparation	Final SDS concentration (%)
Control	0.0	10 mL of DI water	0
T1	0.5	0.05 mL cell extract in 9.95 mL DI water	0.005
T2	1.0	0.1 mL cell extract in 9.9 mL DI water	0.01
T3	5.0	0.5 mL cell extract in 9.5 mL DI water	0.05
T4	10	1 mL cell extract in 9 mL DI water	0.1
T5	25	2.5mL cell extract in 7.5mL DI water	0.25
T6	50	5 mL cell extract in 5 mL DI water	0.5
T7	75	7.5 mL cell extract in 2.5 mL DI water	0.75
T8	100	10 mL cell extract	1.0

7.2.4 Beans production with foliar spray or soil drench of Se-enriched microalgae extract

Non-contaminated soil classified as sandy was collected at a depth of 0–20 cm from a field in Evergem (51°6′57" N, 3°39′40" E), Belgium. The physicochemical properties of the soil were described previously (Chapter 4). The soil was dried, homogenized and passed through a 2 mm sieve mesh. 0.5 kg of the soil was weighed and placed into a 10 cm x 10 cm pot.

Bean seeds (*Phaseolus vulgaris*) were pre-cultivated in trays with wet vermiculite at 27 °C for one week to achieve bean seedlings. Five of the bean seedlings were then transplanted into each pot. Potted plants were grown for 6 weeks indoors (at 24 °C, 53% relative humidity and 100 $\mu\text{mol}/\text{m}^2/\text{s}$ light intensity) with 80% of the water holding capacity. A total amount of 50 mL of different concentrations of the Se-enriched microalgae extracts was applied to each pot every week by foliar spray or soil drench, except for the first and last week of the growth period. A 100 mL small spray bottle was used for the foliar application. The concentrations of the Se-enriched microalgae cell extracts were 0%, 0.5%, 1%, 5% and 10%, which were derived from the previous germination test (in the range of 0–100%). This experiment was conducted in triplicate. During foliar application, the soil surface was covered with aluminum foil to prevent

Chapter 7

spray runoff from coming in contact with the potting soil and thus potentially being taken up by the roots. The bean plants were harvested, washed and separated into different tissues (root, stem, leaf and seed) for biomass and Se concentration analysis.

7.2.5 Beans production on soil amendment with Se-enriched microalgae biomass

Different amount (0 g (0%), 0.225 g (0.5%), 0.45 g (1.0%), 2.25 g (5%) and 4.5 g (10%)) of freeze-dried Se-enriched microalgae biomass were mixed thoroughly with 0.5 kg of sandy soil. Five bean seedlings were transplanted into each pot and grown indoors for 6 weeks by maintaining 80% of the water holding capacity as described in Chapter 4. Beans were harvested for determination of biomass weight and Se concentration analysis. During the growth period, soil pore water was collected twice per week by using Rhizon soil moisture samplers (Eijkelkamp Agrisearch Equipment, Giesbeek, the Netherlands) and analyzed for its total Se concentration in order to evaluate the evolution of the Se release into the soil. pH and TOC of the soil pore water were measured before harvest.

7.2.6 Analytical methods

The method for the determination of total Se in plants (beans and microalga biomass), has been described in Chapters 3 and 4.

pH and EC of the microalgae extract were measured by using a pH (Orion Star A211, Thermo fisher scientific, Waltham, MA, USA) and electrical conductivity meter. For the determination of each element content in the microalgae extract and biomass (in Table 7.1), 2 mL of the microalgae extract or 0.2 g of dry biomass were weighed into a digestion vessel followed by the addition of 8 mL or 10 mL of concentrated ultrapure HNO₃, respectively. The tubes were sonicated for 1 h, then placed in a microwave oven (CEM Mars 6, Matthews, NC, USA) for digestion. The digestion programme was the same as aforementioned for the determination of total Se in plants. The digested samples were measured using inductively coupled plasma-optical emission spectrometry (ICP-OES, iCAP 7000, Thermo Scientific, Waltham, MA, USA) for all elements analysis in Table 7.1, except for Se. Se was determined by ICP-MS.

The pH and TOC of the soil pore water was determined as described in Chapter 4.

Chapter 7

7.2.7 Statistical analysis

Statistical differences were identified with the ANOVA and Duncan's multiple comparison tests at 95% confidence interval level using SPSS 20.0. The Shapiro–Wilk test was used to assess the normality of the distribution of residuals in each data set before conducting ANOVA analysis. The germination percentage (GP) and germination index (GI) were calculated as described in Hernández-Herrera et al. (Hernández-Herrera et al., 2013):

$$GP = \frac{\text{Number of germinated seeds}}{\text{Total number of seeds}} \times 100 \quad (1)$$

$$GI = \sum \left(\frac{G_t}{T_t} \right) \quad (2)$$

Where G_t is the number of seeds germinated on day t and T_t is the number of days.

The mean germination time (MGT) was estimated according to (Ellis & Roberts, 1981; Hernández-Herrera et al., 2013):

$$MGT(d) = \frac{\sum (G_t \times T_t)}{\sum T_t} \quad (3)$$

The seedling vigor index (SVI) was determined by the following formula (Hernández-Herrera et al., 2013; Orchard, 1977):

$$SVI = \text{Seedling length (cm)} \times GP \quad (4)$$

7.3 Results and discussion

7.3.1 Effect of Se-enriched microalgae extract on seed germination and growth of bean seedlings

The germination percentage of bean seeds is shown in Fig. 7.1. Germination occurred in all treatments after 2 days. The Se-enriched microalgae extract showed a slightly stimulatory effect on seed germination at low concentrations, but an inhibitory effect at higher concentrations. Specifically, the maximum GP among all treatments was found at the concentration of 1% microalgae extract. Conversely, 5% of microalgae extract had no significant impact on GP in comparison to the control, except for a significant decrease (by 33%, $P \leq 0.05$) at day 3. However, 10–75% of the microalgae extract

Chapter 7

significantly ($P \leq 0.05$) delayed the bean seed germination, showing a remarkable decline of GP the first 4 days after sowing. It should be noted that the undiluted microalgae extract (100%) treatment significantly dropped off the GP during the entire germination period.

The microalgae extract had a significant ($P \leq 0.05$) effect on the growth of the bean seedlings (Fig. 7.2). Similar to its effect on the GP, the microalgae extracts stimulated the growth of the bean seedlings at low concentrations ($\leq 5\%$). The highest seedling length was observed for the bean seeds presoaked in 1% of the microalgae extract, which was around 2 times higher than that in the control. The addition of Se-enriched microalgae extract at the concentration range of 10–75% had no significant effect on the seedling length. However, 100% of Se-enriched microalgae extract application significantly ($P \leq 0.05$) decreased the seedlings length on day 2 in comparison with the control, although no obvious difference between them was noted on day 4 (Fig. 7.2).




Figure 7.1. Effect of Se-enriched microalgae extracts treatment on the bean seeds germination percentage (GP). Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences between treatments within the same day according to Duncan's multiple comparison tests ($P \leq 0.05$).

Chapter 7




Figure 7.2. Effect of Se-enriched microalgae extracts on the length of bean seedlings. Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences between treatments within the same day according to Duncan's multiple comparison tests ($P \leq 0.05$).

Furthermore, seeds presoaked in 1% of microalgae extract showed the maximum GI and SVI (higher than the control treatment for SVI, $P \leq 0.05$), but the shortest MGT (Fig. 7.3). These results indicate that low concentrations of microalgae extract ($\leq 1\%$) can act as biostimulant for bean seeds germination. Therefore, low concentrations (0.5 and 1%) of microalgae extract were selected for the subsequent pot experiments to further study their effects on the bean plant growth. Besides, taking into account that the function of the microalgae extract might be diluted by the soil or plant after being applied in pot experiments, slightly higher concentrations (5 and 10%) were selected for the subsequent experiments as well.

Previous studies have also evaluated the beneficial effects of microalgae or macroalgae extracts on the germination and growth of different crops (Chiaiese et al., 2018). For instance, Gupta and Shukla (Gupta & Shukla, 1969) studied the effect of *Phormidium* species extracts on the growth of rice seedlings and demonstrated that presoaking rice seeds with algal extracts had a markedly beneficial impact on the development of both roots and shoots. The greatest effect was observed with 1 and 5% of algal extracts through ether and water extraction, respectively. Hernández-Herrera et al. (2014) evidenced that tomato seeds (*Solanum lycopersicum* L.) presoaked in 0.2%

Chapter 7

of *Ulva lactuca* and *Padina gymnospora* (Kützinger) Sonder extracts showed an enhanced germination rate and greater plumule and radicle length. The study of Kumar and Sahoo (Kumar & Sahoo, 2011) also showed that the application of 20% of *Sargassum wightii* Grev. extracts significantly enhanced the germination of wheat seeds (*Triticum aestivum* var. Pusa Gold) and seedling shoot and root growth. Besides, there is still a chance that the positive effect originates from the additional Se. Therefore, performing germination tests upon pure chemical Se exposure can help a bit to understand if the stimulation effect originates from Se or other compounds. To do this, identification of Se species in the microalgae extract is needed in order to provide information on pure chemical Se species to be dosed in such germination test.

The significantly lower GP, GI and SVI observed for the bean seeds presoaked in a high concentration (e.g. 100%) of the microalgae extracts could be explained by salinity stress, which can be deduced from the high EC (6.10 mS/cm) and salt content (e.g., Na^+ , Ca^{2+} , Mg^{2+} and K^+) of the liquid microalgae extract (Table 7.1). The osmotic pressure caused by the high salt content would inhibit the seeds ability to imbibe water (Coppens et al., 2015; Hernández-Herrera et al., 2013), resulting in adverse effects on seeds germination and, eventually, seedling growth. Kaveh et al. (2011) evidenced that increasing salinity levels from 2.5 to 10 mS/cm (EC) delayed the germination percentage and rate, as well as the emergence percentage and rate of all tested tomato species. Likewise, Hernández-Herrera et al. (2013) elucidated that the negative effects of high concentrations of macroalgae extracts on the germination and growth of tomato could be a result of high salinity (around 4.00 mS/cm). Besides, ion toxicity could also explain the detrimental effects of the highly concentrated microalgae extracts on seeds germination and seedling growth. The high concentration of ions in the non-diluted microalgae extracts, such as Na^+ (616 mg/L), K^+ (476 mg/L) and Ca^{2+} (280 mg/L), can be toxic to the embryo and developing seedlings (Benlloch et al., 1994).

Chapter 7




Figure 7.3. Effect of Se-enriched microalgae extracts on the germination parameters of bean seeds: (a) germination index (GI), (b) mean germination time (MGT), (c) seedling vigor index (SVI) at day 2, and (d) SVI at day 4. Values are average $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences between treatments according to Duncan's multiple comparison tests ($P \leq 0.05$).

Chapter 7

7.3.2 Effect of foliar and soil drench application of Se-enriched microalgae extract on the bean plant growth and selenium concentration in the bean plants

7.3.2.1 Bean plant growth

Application as a foliar spray was more effective in influencing the growth of bean plants than the soil drench application (Fig. 7.4). Bean plants treated with the foliar spray at 1% microalgae extract displayed a significant increase of 65 and 29% ($P \leq 0.05$) in the fresh and dry weight of the whole plants, respectively (Figs. 7.4a and 7.4c). An obvious increase in fresh and dry weight of the roots (113 and 51%, respectively) and seeds (364 and 252%, respectively) (Tables S7.2 and S7.3 in supplementary information) was observed ($P \leq 0.05$). Basically, no significant difference in the total fresh and dry weight of the entire plant was recorded when beans received the microalgae extract in the soil drench (Figs. 7.4b and 7.4d). Analysis of the significant difference in each tissue of the beans showed that 1% of the microalgae extract applied as soil drench resulted in a significant increase (approximately 13%, $P \leq 0.05$) in the fresh and dry weight of bean seeds (Tables 7.S2 and 7.S3). It should be noted that the relatively high concentration of microalgae extract (5 and 10%) did not inhibit the growth of beans through the foliar spray and soil drench application (Fig. 7.4).

These results indicated that the microalgae extract exhibits growth-stimulating activities on beans, which is partially consistent with other studies. Kumar and Sahoo (2011) found that 20% of macroalgal seaweed extracts obtained by water boil extraction significantly increased the yield of wheat (*Triticum aestivum* var. Pusa Gold) by 22.86%, measured as dry weight of seeds. . Application of 50% of *Chlorella vulgaris* extract by foliar spray resulted in an obvious increment of the yield in wheat (*Triticum aestivum* L. var. Giza 69) of more than 140% over the control (Shaaban, 2001a). A substantial increase in the yield of eggplants was achieved by foliar spray of commercial Spirufert® fertilizer (*Spirulina platensis*) (Dias et al., 2016). The more effectiveness of foliar application on stimulation plant growth is expected, as liquid microalgae extract being directly applied onto plants via spraying, which avoids the retention of biostimulant by the soil (e.g. binding by soil organic matter), thus improving the microalgae extract utilization by plants.

Chapter 7

The stimulation effects of the microalgae extract on plant growth could be due to the presence of growth-promoting substances such as macro- and microelement nutrients (Table 7.1), amino acids, vitamins and phytohormones (e.g., cytokinins, auxins and gibberellins) that affect cellular physiology (e.g. cell division and cell elongation) in plants, leading to enhanced growth and crop yield (Hernández-Herrera et al., 2013). Another possibility is the presence of polysaccharides (e.g. carboxyated and sulfated polysaccharides or uronic acids) in the microalgae extract (Rachidi et al., 2020), which can improve plant growth in a similar way to hormones (Hernández-Herrera et al., 2013; Rolland et al., 2002).




Figure 7.4. Effect of Se-enriched microalgae extracts applied as foliar spray (a, c) and soil drench (b, d) on fresh weight (a, b) and dry weight (c, d) of bean tissues. Values are average $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences between different treatments according to Duncan's multiple comparison tests ($P \leq 0.05$).

7.3.2.2 Se concentration in the plants and seeds

Chapter 7

Foliar spray and soil drench of the Se-enriched microalgae extract gradually increased the Se content in the bean plant (Fig. 7.5). Generally, foliar spray of Se-enriched microalgae extract provided a higher Se content in the leaves and seeds of the bean plants, while soil drench application resulted in a moderate Se content in the roots and stems of the bean plant. Increasing the application of microalgae extract from 0 to 10% by foliar spray increased the content of Se in the leaves, stems and seeds of beans by 6.2, 2.5 and 1.7 times, respectively (Fig. 7.5a), whereas no significant difference was found in the roots. On the other hand, the application of 10% microalgae extract as soil drench significantly increased the Se content in the roots and stems of the beans by 1.6 and 3.8 times, respectively, in comparison with the control (Fig. 7.5b), whereas, the Se content in the seeds of beans was not significantly different between soil drench application treatments. This indicates a slow translocation of Se from the roots to seeds in bean plants.

It should be noted that the Se concentration in all tissues after the application of Se-enriched microalgae extract at doses below 1% showed no significant difference with the control (except for the leaf with foliar spray) (Fig. 7.5), whereas the highest fresh and dry biomass of beans was observed at 1% of Se-enriched microalgae extract application (Fig. 7.4). This indicates that the dose of microalgae extract and Se concentration in the extract needs to be carefully determined in order to balance out good nutrition (e.g. Se content) and obtain the highest yield of bean seeds.




Figure 7.5. Effect of Se-enriched microalgae extracts applied as (a) foliar spray and (b) soil drench on the Se concentration in bean tissues. Values are average $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences between treatments in the same tissue according to Duncan's multiple comparison tests ($P \leq 0.05$).

7.3.3 Effect of Se-enriched microalgae biomass on the bean plant growth and Se concentration in the plants

7.3.3.1 Bean plant growth

Fig. 7.6 illustrates the fresh and dry weight of the beans grown in soil amended with Se-enriched microalgae biomass. The supplementation of Se-enriched microalgae biomass significantly stimulated the growth of the whole plant, except for the 0.5% Se-

Chapter 7

enriched microalgae amendment. The highest biomass yield was found at 1% Se-enriched microalgae addition, similarly to the foliar application of microalgae extract in the previous experiment. It gave an increase of 64 and 43%, respectively, in fresh and dry biomass of the whole plant. Besides, among all tissues, the roots and seeds of the beans were more sensitive to the Se-enriched microalgae supplementation compared to the leaves and stems, as reflected in the significant increase of biomass in the roots and seeds for the 0.5-5% of Se-enriched microalgae supplementation, but absence of a considerable increase in the leaf and stems biomass (Table 7.S4). Approximately 4 times higher seed yields (both fresh and dry weight) were obtained for the 1% of Se-enriched microalgae amendment compared to the control.

In line with these results, Shaaban (2001b) reported an increase in the dry weight of shoots and roots of maize (*Zea mays* L.) grown in a soil amended with the microalgae *Chlorella vulgaris*. The best treatments were 150 and 200 kg algae/Fed (1 Feddan = 0.42 hectare) (Shaaban, 2001b). As aforementioned in growth-stimulation by the microalgae extracts, the stimulation effects of the microalgae biomass on beans growth was partially a result of the slow release of macro- and micro-nutrients from the microalgae biomass, particularly N and P, which have the same effects on plant growth as inorganic fertilizer (Mulbry et al., 2005). The applied microalgae biomass was composed of 48% protein and 32% carbohydrates and 21% lipids (Chapter 6), thus enhancing the nitrogen and carbon content upon its application to the soil, leading to an increase in the soil microbial activity and potentially promoting plant growth. The presence of plant biostimulants (e.g., amino acids, polysaccharides and phytohormones) (Kumar & Sahoo, 2011) contained in the microalgae biomass could also have contributed to the positive effects on the beans growth.

Chapter 7




Figure 7.6. Effect of Se-enriched microalgae biomass on the (a) fresh and (b) dry weight of beans. Values are average $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences between treatments in the same tissue according to Duncan's multiple comparison tests ($P \leq 0.05$).

7.3.3.2 Se concentration in the bean plants

The Se concentration in the bean plants depended on the amount of Se-enriched microalgae supplemented to the soil (Fig. 7.7). The Se content in all tissues of the bean plant raised gradually with the increasing dosage of Se-enriched microalgae amendment (Fig. 7.7). An increment of Se-enriched microalgae dosage from 0 to 10% increased the Se content of the beans stepwise from 1.05 to 4.15 mg/kg in the root,

Chapter 7

0.12 to 0.34 mg/kg in the leaf, 0.09 to 0.42 mg/kg in the stem and 0.10 to 0.28 mg/kg in the seed.




Figure 7.7 Effect of Se-enriched microalgae biomass on the Se concentration in the different tissues of beans. Values are mean $\pm$ standard deviation ($n=3$). Different letters indicate statistically significant differences between treatments in the same tissue according to Duncan's multiple comparison tests ($P \leq 0.05$).

It should be noted that the Se content in all tissues of beans at 1% of Se-enriched supplement did not significantly differ from that of the control and it was even slightly lower than that of the supplement at 0.5% dosage, which is partially similar to the results observed in foliar and soil drench application of Se-enriched microalgae extract (Fig. 7.5). This may be related to the greatest amount of biomass being produced at 1% of Se-enriched microalgae biomass/extract addition among all dosages (Figs. 7.4 and 7.6), which resulted in a biological dilution of Se in the plant tissues due to the significant biomass increase. This is supported by the obviously higher Se accumulation in beans at 1% of Se-enriched microalgae biomass application in comparison with that of the control among all tissues, except for the stem ($P \leq 0.05$) (Table 7.S5).

Taking into account the biomass (Fig. 7.6) and Se content (Fig. 7.7) of the bean plants, 5% of dried Se-enriched microalgae biomass was selected as a recommended dosage

Chapter 7

in practice, which could not only increase the Se concentration in the seeds of the beans, but also enhance the growth of the bean plants.

7.3.3.3 Evolution of Se content in the soil pore water during the growth period

The Se content in the pore water of the soil amended with Se-enriched microalgae biomass during the entire growth period of the beans is shown in Fig. 7.8 and Table 7.S6. Amendment with Se-enriched microalgae from 0 to 10% gradually and significantly increased the Se content in the soil pore water. Specifically, the addition of 0.5, 1, 5 and 10% of Se-enriched microalgae biomass increased the Se content in the soil pore water to 6.89, 9.20, 21.7, 30.9 $\mu\text{g/L}$ after the first day, which was 3, 4, 9 and 13 times higher than that of the control, respectively. However, after 22 days of growth of the beans, the Se content in the pore water of the soil amended with Se-enriched microalgae at doses below 5% did not show statistically significant difference with the control (Table 7.S6).

For the same amount of Se-enriched microalgae addition, the Se content in the soil pore water significantly declined along with the growth time during the first 22 days and was stable afterwards (Fig. 7.8 and Table 7.S6). Increasing the growth time from 1 to 22 days reduced the Se content in the soil pore water by 74, 79, 83 and 68% for the 0.5, 1, 5 and 10 amendment dose, respectively.

The Se content in the soil can indeed be increased by the application of Se-enriched organic materials, which was also reported in other studies. For instance, Bañuelos et al. (2015) observed that the dose of Se-enriched *Stanleya pinnata* applied was positively correlated to the soluble and bioavailable Se content in soils. The application of Se-enriched wheat (*Triticum aestivum* L.) and raya (*Brassica juncea* L.) straw from 0 to 1% (ratio of straw weight to soil weight) increased the hot water-soluble Se (bioavailable Se) fraction in a sandy-loam soil from 18 $\mu\text{g/kg}$ to 36 and 79 $\mu\text{g/kg}$, respectively (Dhillon et al., 2007). The significant decrease of Se in the soil pore water during the first 22 days of beans growth could be attributed to the fast adsorption of the released Se onto soil or organic matter and to Se uptake by bean plants (Li et al., 2017).

Se-enriched microalgae biomass could be considered as a slow-release Se biofertilizer because only around 3% of Se in the biomass was released to the soil pore water on

Chapter 7

the first day after application of high concentrations ($\geq 5\%$) of biomass (Table 7.S7). The higher Se accumulation in the beans compared to the Se content in the soil pore water (on the first day) also evidenced that more Se was slowly released from the Se-enriched microalgae matrix and gradually supplied for uptake by the beans during the entire growth period (Table 7.S7).




Figure 7.8. Evolution of the Se concentration in the pore water of soil amended with Se-enriched microalgae over the growth period of beans. Values are mean $\pm$ standard deviation ($n=3$).

7.3.3.4 pH and TOC in the soil pore water

The pH and TOC in the pore water of the soil amended with Se-enriched microalgae were measured at the time of harvest (Table 7.3). The addition of 10% of Se-enriched microalgae biomass noticeably increased the pore water pH and TOC ($P \leq 0.05$), while other applications did not result in a remarkable difference in comparison with the control, except for the TOC at 5% microalgae amendment. The highest TOC found at the highest application rate was 149 mg/L, which was 3 times higher than that of the control.

Chapter 7

Table 7.3. pH and TOC content in the pore water of the soil amended with different amounts of Se-enriched microalgae biomass at the time of beans harvesting. Values are mean $\pm$ standard deviation (n=3). Different letters indicate statistically significant differences between treatments according to Duncan's multiple comparison tests ($P \leq 0.05$).

	Control	0.5%	1%	5%	10%
pH	5.2 $\pm$ 0.3 ^c	6.2 $\pm$ 0.4 ^{ab}	6.0 $\pm$ 0.6 ^{abc}	5.3 $\pm$ 0.5 ^{bc}	6.5 $\pm$ 0.7 ^a
TOC (mg/L)	50 $\pm$ 7.7 ^c	47 $\pm$ 12.6 ^c	39 $\pm$ 6.6 ^c	65 $\pm$ 2.8 ^b	149 $\pm$ 8.7 ^a

7.3.4 Practical implications of using Se-enriched microalgae biomass and its extract as biostimulants and Se sources

This study demonstrated that the Se-enriched microalgae biomass and its extract produced from Se-enriched domestic wastewater have the potential to be used as biostimulant and Se biofertilizer for enhancement of beans growth and Se biofortification. However, the potential loading of heavy metals, micropollutants and pathogens onto the biomass is still a main concern for the application of Se-enriched microalgae produced from domestic wastewater. The EU fertilizing products and amending regulation (2019) stipulates that contaminants in an organic fertilizer must not exceed the following limit values (expressed as mg/kg dry matter): Cd 1.5, Cr (VI) 2.0, Hg 1.0, Ni 50, Pb 120 and As 40. The Cu and Zn content must not exceed 300 and 800 mg/kg dry matter, respectively. Pathogen loads must not exceed the following limits: *Salmonella* spp. absence in 25 g or 25 mL and *Escherichia coli* or *Enterococcaceae* 1000 CFU in 1 g or 1 mL.

In this study, the level of all heavy metals in the Se-enriched microalgae biomass was much below the safety limits, with the exception of Cr (Table 7.1). It should be noted that the fertilizer regulation limited the Cr(VI) content instead of total Cr, since Cr(VI) is both toxic and carcinogenic, while other Cr species (e.g. Cr (0) and Cr (III)) are considered not toxic (Kimbrough et al., 1999). In most cases, Cr (III) is the dominating species in the environment and food (Kimbrough et al., 1999). The Cr species in the microalgae biomass in this study is possibly also be dominated by Cr (III), but additional analyses are needed to confirm this.

Chapter 7

Besides, the pathogen load in the Se-enriched microalgae biomass used in this study have been characterized in our previous study (in Chapter 6). It could be confirmed that *Salmonella* spp. and *Escherichia coli* were absent in the freeze-dried biomass, and most microorganisms were reduced after drying processes. Eventhough the heavy metal contents and pathogen loads of the microalgae biomass did not exceed the European regulation of fertilizer products, more research is still needed to confirm environmental safety and assess human health risks related to the use of Se-enriched microalgae as biostimulant in crop production. Besides, further studies should be conducted to assess the effect of the Se-enriched microalgae and their extracts on Se accumulation in crops and growth of these crops under field conditions. Lastly, improvement of the Se extraction efficiency from the Se-enriched microalgae is required, as Se extraction efficiency rom microalgae biomass is relatively low (15.4%). Combination of enzymatic and mechanical techniques would help to disrupt the robust microalgae cell wall and improve the Se release.

7.4 Conclusions

Application of relatively low dosages of Se-enriched microalgae extract was beneficial for seed germination ($\leq 1\%$ dosage) and seedling growth ($\leq 5\%$ dosage) of beans, while high dosages ($> 50\%$) significantly delayed the mean germination time. Foliar application of Se-enriched microalgae extract was more effective to stimulate the bean growth and increase the Se concentration in the seeds compared to soil drench application. 5% dosage of Se-enriched microalgae biomass could be used as a biostimulant enhancing the growth of bean plants, and as an organic slow-release Se biofertilizer significantly improving the Se content in the beans, including the seeds. These results indicate that Se-enriched microalgae biomass and its extracts could potentially be used as an added-value biostimulant and Se biofertilizer replacing conventional mineral Se fertilizers in biofortification schemes. As the microalgae were generated during domestic wastewater treatment, the proposed approach contributes to resource recovery within the framework of the circular economy.

Chapter 7

Supplementary Information




Figure 7.S1 Optic microscope images of the microalgae biomass composition over the experimental period (scale 10 um).

Chapter 7

Table 7.S1. Effect of Se-enriched microalgae extracts on germination parameters of bean seeds: germination index (GI), mean germination time (MGT) and seedling vigor index (SVI). Values are average $\pm$ standard deviation (n=3). Different letters within columns indicate statistically significant differences between treatments according to Duncan's multiple comparison tests ($P \leq 0.05$).

Microalgae extract	Parameter			
	GI	MGT (days)	SVI at 2d	SVI at 4d
Control	16.5 $\pm$ 3.4 ^a	4.15 $\pm$ 0.0 ^d	0.62 $\pm$ 0.2 ^c	1.86 $\pm$ 0.3 ^{bc}
0.5%	16.2 $\pm$ 1.8 ^a	4.21 $\pm$ 0.1 ^{cd}	0.91 $\pm$ 0.3 ^b	2.14 $\pm$ 0.4 ^{ab}
1%	18.2 $\pm$ 0.7 ^a	4.11 $\pm$ 0.0 ^d	1.43 $\pm$ 0.0 ^a	2.58 $\pm$ 0.6 ^a
5%	12.1 $\pm$ 0.2 ^b	4.14 $\pm$ 0.1 ^d	0.72 $\pm$ 0.0 ^{bc}	1.76 $\pm$ 0.6 ^{bcd}
10%	11.5 $\pm$ 0.9 ^b	4.28 $\pm$ 0.1 ^{bcd}	0.52 $\pm$ 0.2 ^{cd}	1.16 $\pm$ 0.2 ^{def}
25%	11.4 $\pm$ 0.0 ^b	4.21 $\pm$ 0.1 ^{cd}	0.31 $\pm$ 0.0 ^{de}	0.93 $\pm$ 0.1 ^{ef}
50%	10.8 $\pm$ 1.7 ^b	4.34 $\pm$ 0.1 ^{bc}	0.25 $\pm$ 0.1 ^{de}	1.31 $\pm$ 0.3 ^{cde}
75%	10.3 $\pm$ 1.3 ^b	4.43 $\pm$ 0.2 ^b	0.18 $\pm$ 0.1 ^f	1.30 $\pm$ 0.1 ^{cde}
100%	5.8 $\pm$ 0.6 ^c	4.60 $\pm$ 0.1 ^a	0.05 $\pm$ 0.0 ^f	0.60 $\pm$ 0.1 ^f

Chapter 7

Table 7.S2. Effect of Se-enriched microalgae extracts applied as foliar spray and soil drench on fresh weight (g) of bean tissues at different doses. Values are average $\pm$ standard deviation (n=3). Different letters within columns indicate statistically significant differences between treatments according to Duncan's multiple comparison tests ($P \leq 0.05$).

Treatment	Foliar spray				Soil drench			
	Root	Leaf	Stem	Seed	Root	Leaf	Stem	Seed
Control	5.9 $\pm$ 1.1 ^b	9.7 $\pm$ 0.9 ^b	10.9 $\pm$ 0.8 ^a	1.8 $\pm$ 1.3 ^b	8.9 $\pm$ 1.4 ^a	5.2 $\pm$ 0.8 ^a	9.5 $\pm$ 0.1 ^a	9.6 $\pm$ 0.2 ^{ab}
0.5%	8.9 $\pm$ 2.2 ^{ab}	10.1 $\pm$ 0.9 ^{ab}	11.5 $\pm$ 0.5 ^a	6.3 $\pm$ 2.1 ^{ab}	9.8 $\pm$ 0.3 ^a	5.7 $\pm$ 3.2 ^a	9.9 $\pm$ 2.5 ^a	9.5 $\pm$ 2.5 ^{ab}
1%	12.6 $\pm$ 2.2 ^a	12.9 $\pm$ 1.1 ^a	13.0 $\pm$ 0.6 ^a	8.2 $\pm$ 4.0 ^a	8.7 $\pm$ 2.1 ^a	8.7 $\pm$ 2.4 ^a	12.1 $\pm$ 2.7 ^a	11.0 $\pm$ 1.7 ^a
5%	6.9 $\pm$ 0.4 ^b	8.5 $\pm$ 1.7 ^b	10.9 $\pm$ 1.6 ^a	1.4 $\pm$ 0.2 ^b	7.6 $\pm$ 4.7 ^a	6.5 $\pm$ 4.9 ^a	10.8 $\pm$ 3.1 ^a	3.4 $\pm$ 0.2 ^c
10%	7.2 $\pm$ 3.3 ^b	7.7 $\pm$ 0.9 ^b	9.9 $\pm$ 2.0 ^a	1.1 $\pm$ 0.8 ^b	7.1 $\pm$ 1.0 ^a	8.5 $\pm$ 1.8 ^a	11.4 $\pm$ 0.5 ^a	5.9 $\pm$ 2.6 ^{bc}

Chapter 7

Table 7.S3. Effect of Se-enriched microalgae extract treatment applied as foliar spray and soil drench on dry weight (g) of bean tissues at different doses. Values are average $\pm$ standard deviation (n=3). Different letters within columns indicate statistically significant differences between treatments according to Duncan's multiple comparison tests ($P \leq 0.05$).

Treatment	Foliar spray				Soil drench			
	Root	Leaf	Stem	Seed	Root	Leaf	Stem	Seed
Control	0.55 $\pm$ 0.0 ^b	2.23 $\pm$ 0.0 ^a	1.72 $\pm$ 0.1 ^a	0.26 $\pm$ 0.1 ^{bc}	0.66 $\pm$ 0.1 ^a	1.78 $\pm$ 0.1 ^a	1.64 $\pm$ 0.1 ^a	0.81 $\pm$ 0.0 ^a
0.5%	0.59 $\pm$ 0.1 ^b	2.15 $\pm$ 0.2 ^a	1.95 $\pm$ 0.2 ^a	0.58 $\pm$ 0.2 ^{ab}	0.73 $\pm$ 0.2 ^a	1.70 $\pm$ 0.1 ^a	1.58 $\pm$ 0.1 ^a	0.82 $\pm$ 0.2 ^a
1%	0.83 $\pm$ 0.2 ^a	2.43 $\pm$ 0.2 ^a	2.09 $\pm$ 0.0 ^a	0.92 $\pm$ 0.4 ^a	0.63 $\pm$ 0.1 ^a	1.87 $\pm$ 0.3 ^a	1.89 $\pm$ 0.2 ^a	0.92 $\pm$ 0.2 ^a
5%	0.60 $\pm$ 0.1 ^b	2.44 $\pm$ 0.2 ^a	2.02 $\pm$ 0.2 ^a	0.20 $\pm$ 0.0 ^{bc}	0.77 $\pm$ 0.4 ^a	1.81 $\pm$ 0.5 ^a	1.70 $\pm$ 0.4 ^a	0.30 $\pm$ 0.0 ^b
10%	0.55 $\pm$ 0.1 ^b	2.19 $\pm$ 0.0 ^a	1.82 $\pm$ 0.1 ^a	0.13 $\pm$ 0.6 ^c	0.54 $\pm$ 0.1 ^a	1.99 $\pm$ 0.4 ^a	1.69 $\pm$ 0.3 ^a	0.50 $\pm$ 0.2 ^b

Chapter 7

Table 7.S4. Effect of Se-enriched microalgae biomass soil application on the fresh and dry weight of beans (g). Values are average $\pm$ standard deviation (n=3). Different letters within columns indicate statistically significant differences between treatments in the same tissue according to Duncan's multiple comparison tests ($P \leq 0.05$).

Treatment	Fresh weight				Dry weight			
	Root	Leaf	Stem	Seed	Root	Leaf	Stem	Seed
Control	4.31 $\pm$ 0.2 ^c	7.11 $\pm$ 0.5 ^b	10.4 $\pm$ 0.9 ^a	3.65 $\pm$ 0.2 ^c	0.30 $\pm$ 0.1 ^b	1.75 $\pm$ 0.2 ^a	1.42 $\pm$ 0.0 ^a	0.25 $\pm$ 0.1 ^c
0.5%	5.64 $\pm$ 0.4 ^c	8.01 $\pm$ 1.4 ^b	9.69 $\pm$ 1.6 ^a	8.28 $\pm$ 0.2 ^b	0.54 $\pm$ 0.1 ^b	1.85 $\pm$ 0.4 ^a	1.52 $\pm$ 0.2 ^a	0.68 $\pm$ 0.0 ^b
1%	9.89 $\pm$ 2.6 ^a	6.44 $\pm$ 1.9 ^b	9.27 $\pm$ 0.6 ^a	14.2 $\pm$ 1.5 ^a	0.86 $\pm$ 0.1 ^a	1.72 $\pm$ 0.1 ^a	1.49 $\pm$ 0.1 ^a	1.10 $\pm$ 0.0 ^a
5%	8.09 $\pm$ 3.1 ^{ab}	8.03 $\pm$ 1.5 ^b	10.8 $\pm$ 1.7 ^a	9.95 $\pm$ 1.3 ^b	0.57 $\pm$ 0.2 ^b	2.04 $\pm$ 0.2 ^a	1.64 $\pm$ 0.2 ^a	0.78 $\pm$ 0.1 ^b
10%	6.35 $\pm$ 1.0 ^{ab}	12.5 $\pm$ 1.4 ^a	11.7 $\pm$ 2.4 ^a	3.67 $\pm$ 1.7 ^c	0.39 $\pm$ 0.1 ^b	2.01 $\pm$ 0.4 ^a	1.46 $\pm$ 0.4 ^a	0.29 $\pm$ 0.1 ^c

Chapter 7

Table 7.S5. Effect of Se-enriched microalgae biomass application to soil on Se accumulation ($\mu\text{g/pot}$) in different tissues of beans. Values are average $\pm$ standard deviation ($n=3$). Different letters within columns indicate statistically significant differences between treatments in the same tissue according to Duncan's multiple comparison tests ($P \leq 0.05$).

Treatment	Se accumulation			
	Root	Leaf	Stem	Seed
Control	0.32 ± 0.1^c	0.21 ± 0.0^d	0.13 ± 0.0^b	0.02 ± 0.0^c
0.5%	0.63 ± 0.2^c	0.32 ± 0.0^c	0.15 ± 0.0^b	0.10 ± 0.0^b
1%	0.99 ± 0.1^b	0.27 ± 0.0^{cd}	0.15 ± 0.0^b	0.15 ± 0.0^a
5%	1.11 ± 0.1^b	0.44 ± 0.1^b	0.31 ± 0.1^b	0.14 ± 0.0^a
10%	1.47 ± 0.2^a	0.65 ± 0.1^a	0.58 ± 0.2^a	0.07 ± 0.0^b

Note: Se accumulation ($\mu\text{g/pot}$) was calculated by multiplying the Se concentration in tissues ($\mu\text{g/g}$) by the dry weight of corresponding tissues (g).

Chapter 7

Table 7.S6. Selenium concentrations ($\mu\text{g/L}$) in the pore water extracted from soil amended with Se-enriched microalgae biomass. Mean $\pm$ standard deviation, $n=3$. Different lowercase letters indicate statistically significant differences between different incubation days according to Duncan's multiple comparison tests ($P < 0.05$). Uppercase indicates significant differences between different doses.

	Day 1	Day 8	Day 15	Day 22	Day 33	Day 42
Control	$2.45 \pm 0.6^{\text{D}}$	$1.69 \pm 0.2^{\text{D}}$	$0.95 \pm 0.6^{\text{C}}$	$0.87 \pm 0.6^{\text{D}}$	$1.33 \pm 0.1^{\text{B}}$	$1.26 \pm 0.1^{\text{B}}$
0.5%	$6.89 \pm 1.1 \text{ a}^{\text{C}}$	$2.42 \pm 0.3 \text{ b}^{\text{CD}}$	$1.77 \pm 0.3 \text{ bc}^{\text{C}}$	$1.82 \pm 0.3 \text{ bc}^{\text{C}}$	$1.44 \pm 0.3 \text{ bc}^{\text{B}}$	$1.13 \pm 0.2 \text{ c}^{\text{B}}$
1%	$9.20 \pm 3.1 \text{ a}^{\text{C}}$	$3.91 \pm 1.3 \text{ b}^{\text{C}}$	$2.18 \pm 0.4 \text{ b}^{\text{C}}$	$1.91 \pm 0.2 \text{ b}^{\text{C}}$	$1.70 \pm 0.1 \text{ b}^{\text{B}}$	$1.55 \pm 0.5 \text{ b}^{\text{B}}$
5%	$21.7 \pm 7.2 \text{ a}^{\text{B}}$	$10.2 \pm 1.3 \text{ b}^{\text{B}}$	$5.37 \pm 0.7 \text{ bc}^{\text{B}}$	$3.74 \pm 0.2 \text{ c}^{\text{B}}$	$2.86 \pm 0.1 \text{ c}^{\text{B}}$	$2.73 \pm 0.8 \text{ c}^{\text{B}}$
10%	$30.9 \pm 0.9 \text{ a}^{\text{A}}$	$18.5 \pm 1.3 \text{ b}^{\text{A}}$	$10.5 \pm 3.3 \text{ c}^{\text{A}}$	$9.73 \pm 0.6 \text{ c}^{\text{A}}$	$10.6 \pm 1.9 \text{ c}^{\text{A}}$	$11.1 \pm 1.7 \text{ c}^{\text{A}}$

Chapter 7

Table 7.S7. Selenium mass balance calculated from the amount of Se applied through Se-enriched microalgae biomass, the Se content in the soil pore water and the Se accumulated in the plant.

Treatment	Applied Se *	Se in pore water ** (first day)	Se in pore water (last day)	Se accumulation by plant***	Se in pore water (first day) / applied Se	Se in plants / applied Se
		[µg/pot]			%	%
0.5%	6.53	0.91	0.15	1.20	9	18
1%	13.1	1.21	0.20	1.56	7	12
5%	65	2.86	0.36	2.00	4	3
10%	130	4.08	1.47	2.76	3	2

Note:

* *Applied Se = Se concentration in the microalgae × weight of microalgae biomass applied in each pot*

** *Se in pore water = Se concentration in the soil pore water × water content of the soil*

***Se accumulation in the plant is the sum of Se accumulation in each plant tissue.

Chapter 8 General discussion, conclusions and future perspectives

Chapter 8

The main objective of this PhD dissertation was to explore the possibility of producing micronutrient-enriched biomaterials on wastewater as feedstock through ecotechnologies, and using these materials as feed supplement (Chapters 3 and 6) or biofertilizer and biostimulant (Chapters 4, 5 and 7). The used production methods are phytoextraction using duckweed and *Azolla*, bioreduction by anaerobic sludge, and microalgae-based wastewater treatment methods.

8.1 Production of micronutrient-enriched bioproducts as potential feed/food supplements

8.1.1 Production and valorization of Se/Zn-enriched duckweed as potential feed supplements

Lemna (named Duckweed later) and *Azolla*, two aquatic plants with a substantial protein content, were selected to evaluate the possibility to produce Se/Zn-enriched dietary proteins and fertilizers while removing Se/Zn from wastewater. The interaction effects occurring between Se (Se(IV) and Se(VI)) and Zn when these micronutrients are taken up simultaneously by *Lemna* and *Azolla* were assessed as well. The results demonstrated that both plant species could accumulate (around 10 times) more Se(IV) than Se(VI). This is in agreement with what was previously observed for other plants, such as sunflower and maize, when being cultivated in hydroponic systems. A synergetic effect between Se and Zn was observed in *Lemna*, but an antagonistic effect in *Azolla*. This was concluded from the significant increase of the removal efficiency of Se and its accumulation in *Lemna* when increasing the Zn dosage, but the opposite being observed in *Azolla*.

A high content of true protein (approximately 17%) in freeze-dried *Lemna* and *Azolla* and high Se/Zn accumulation in the two plant tissues (up to 1664 mg/kg for Se and 3144 mg/kg for Zn) were observed. Besides, the ability of *Lemna* and *Azolla* to take up and transform inorganic Se(IV) in the growth medium into organic Se (e.g., SeMet, SeCys₂, and SeMetSeCys) in their tissues was validated. Organic Se species are considered beneficial in human and animal nutrition. These results together with the fast growth rate of the plants make *Lemna* and *Azolla* potential candidates for the production of Se/Zn-enriched biomass that can be used as crop fertilizers or protein-rich food/feed supplements or ingredients.

Chapter 8

However, it should be noted that the Se/Zn removal efficiency by *Lemna* and *Azolla* may require further improvement. This may be attributed to the small amount of biomass used at the start of the experiments and the limited water surface area to volume ratio applied in this study. The latter is an important parameter as it concerns floating aquatic plants. Therefore, further study should reconsider the biomass used and the wastewater surface area to volume ratio. Besides, the challenge of dewatering aquatic plants between harvest and their use as feed supplements or fertilizers should also be considered from energy- and cost-saving perspectives.

8.1.2 Production and valorization of Se-enriched microalgae as potential feed supplements

Batch lab-scale and continuous pilot-scale (HRAPs) experiments have been conducted to study the production of Se-enriched microalgae biomass as potential Se-enriched feed supplement and biofertilizer (Chapter 6). Based on the results of the batch test, Se(IV) was preferred over Se(VI) to be supplied to the continuous HRAPs systems for the production of Se-enriched microalgae, as microalgae biomass could accumulate remarkably more Se(IV) than Se(VI). Furthermore, the HRAPs fed with domestic wastewater were operated for 3 months under two HRTs (4 days and 8 days). The HRAPs had a good wastewater treatment performance, with an average COD, NH_4^+ -N, and total phosphorus removal efficiency of, respectively, 70%, 93%, and 77%. This is in line with previous studies (Arashiro et al., 2019; Gutierrez et al., 2016). The Se removal efficiency by HRAP systems under the two HRTs was around 44% at the dose of around 50 μg Se/L for both 4 d and 8 d HRT operation.

Nutritional analyses evidenced the potential of the produced Se-enriched microalgae as feed supplements or alternatives for animal protein, as both the content of protein (around 48%) and the occurrence of essential amino acids in the Se-enriched microalgae were comparable to those of conventional plant-based protein sources used in feed (soybeans). Moreover, Se-enriched microalgae biomass was shown to contain a higher content of fatty acids beneficial for human and animal consumption, such as omega-3, omega-6, and EPA. Moreover, the predominance of selenoamino acids (SeMet, accounting for 91% of the total Se) found in the Se-enriched microalgae grown in the HRAP demonstrates the ability of microalgae to upgrade low-value inorganic Se to high-value products. However, *in vitro* digestibility tests indicated that

Chapter 8

only 49% and 63% of the incorporated Se are bioaccessible for animals in raw and ball-milled Se-enriched microalgae, respectively. This may be due to the robustness of the microalgae cell wall, which cannot be easily disrupted, eventually resulting in the low Se bioaccessibility. In fact, this is currently also the bottleneck for the reuse of microalgae for other purposes, e.g. as biofuel. Therefore, future research should also focus on improving disruption of the cell wall of the microalgae in a cost-effective and energy-efficient manner for improved digestion by animals.

8.2 Valorisation of the produced micronutrient-enriched bioproducts as Se/Zn biofertilizers

8.2.1 Agronomic biofortification of *Phaseolus vulgaris* with Se/Zn-enriched duckweed and sludge

Pot experiments using green beans (*Phaseolus vulgaris*) were conducted to assess the potential of two micronutrient-enriched biomaterials (sludge and duckweed) as slow-release Se and Zn biofertilizers (Chapter 4 and 5). The biomaterials were previously generated using single Se or simultaneous Se and Zn-bearing water as feedstock. The results demonstrated that the Se concentration in the soil pore water is much higher when soil is amended with Se-enriched duckweed in comparison with the soil amended with Se-enriched sludge at the beginning of the incubation. However, the Se-enriched sludge offers more bioavailable Se for plant uptake than the Se-enriched duckweed. Accordingly, higher Se and SeMet concentration and a higher SeMet ratio relative to other species have been found in the bean seeds upon Se-enriched sludge application. This is due to (1) the different Se species and organic matter content present in the two Se-enriched biomaterials. Specifically, elemental nano-Se was the predominant Se species in the Se-enriched sludge, which is relatively stable and bioavailable (for nano-Se), while Se(VI) was the main Se form in the Se-enriched duckweed, which is not stable in the environment and (2) the different Se immobilization rate and transformation in soils after Se release of Se from the two biomaterials.

This thesis thus concluded that the micronutrient-enriched sludge is considered as the preferred slow-release Se biofertilizer for Se-deficient areas, in comparison to micronutrient-enriched duckweed. Besides, the main Se aminoacid species (Se-

Chapter 8

methionine, 76–89%) detected in the bean seeds together with the estimated daily intake (EDI) and health risk index (HRI) indicated that the bean seeds produced by biofortification using Se-enriched duckweed/sludge as fertilizer could contribute significantly to achieving the recommended daily Se intake for human diets and would likely not pose a potential risk of excessive Se intake.

On the other hand, Zn released from the biomaterials was not readily transferred from underground to aboveground plant parts, especially to the seeds. Therefore, the biofortification of Zn through the application of Se/Zn-enriched biomaterials as fertilizers in this thesis was not completely successful. Besides, it should be highlighted that higher amounts of duckweed (>11 g in 0.5 kg soil) application resulted in negative effects on plant growth (i.e. lower biomass yield), which is thus not recommended in practice. Some pre-treatment (e.g., composting, pyrolysis, or extracting) of the micronutrient-enriched duckweed may be needed before application as Se/Zn biofertilizer.

8.2.2 Valorization of Se-enriched microalgae and their extract as potential Se biofertilizers and biostimulants

The use of Se-enriched microalgal biomass and their extract as potential Se biofertilizer and biostimulant to simultaneously enhance plant growth and Se uptake was assessed through soil and foliar application via pot experiments (Chapter 7). Green bean (*Phaseolus vulgaris*), a protein-rich crop regularly grown in moderate climates, was targeted. We concluded that the foliar application of Se-enriched microalgae extracts is recommended in practice, in comparison with the application of Se-enriched microalgae extract as soil drench, because the foliar application of Se-enriched microalgae extracts could obviously enhance the plant yield and meanwhile improve the Se content in the tissues of beans (seeds, leaves, and stems). Besides, the results also verified the possibility of Se-enriched microalgal biomass as a potential Se biofertilizer, which was shown by the remarkable increase of plant biomass and Se concentration in bean tissues after application of Se-enriched microalgae biomass (1% and 5%), compared to that of the control. This could lead to a higher market value of the beans. Overall, this thesis has valorized the use of the high value-added Se-enriched microalgae biomass and their extracts generated from domestic wastewater as biostimulant and biofertilizer.

8.3 Comparison of Se utilization rates in duckweed, microalgae and sludge based production systems

To evaluate efficiency of the production systems of Se-enriched biomaterials based on duckweed, microalgae and sludge, we compared their Se uptake when being exposed to the same Se concentrations in the growth medium. The Se utilization rate by crops when using these Se-enriched duckweed, microalgae and sludge as biofertilizer was compared as well (Table 8.1). The results demonstrated that the Se concentration in the duckweed was the highest upon 0.5 mg/L Se(IV) exposure among the three biomaterials, while microalgae had the lowest Se concentration. This might be attributed to a lower ability microalgae to take up Se and the different growth media used for growing the duckweed and microalgae. A nutrient medium was used for duckweed cultivation, while real domestic wastewater was applied for microalgae growth. The higher organic carbon and solids contents in real wastewater may result in higher retention of applied Se in the wastewater and result in lower Se accumulation in the biomass. Additionally, for Se(VI) exposure, the maximum Se concentration was observed in sludge. In terms of the Se utilization efficiency by beans grown in soil amended with the Se-enriched biomaterials, the sludge application resulted in the highest Se utilization efficiency, while the minimum value was found in the microalgae application.

It should be noted that sludge also showed the greatest Se removal efficiency (>90%, Chapter 4) as well as the highest Se tolerance capacity (> 5 mg/L). This supports the conclusion that Se-enriched sludge can be considered as the preferred slow-release Se biofertilizer, in comparison with duckweed and microalgae, meanwhile removing a high amount of Se from the wastewater. The lower Se utilization efficiency by beans grown on soil amended with microalgae biomass or its extract, compared to duckweed, could be attributed to the low Se accumulation in the microalgae biomass, introducing relatively more organic matter when applied in the soil, subsequently resulting in lower Se utilization by the beans. Besides, the harvest of microalgae is much more difficult than that duckweed, involving a higher energy consumption. However, the faster growth rate of microalgae and lower growth space requirement in comparison with duckweed could still counteract this drawback (Table 8.2). Nevertheless, improvement of Se accumulation in microalgae is considered as a requirement towards the valorization of Se-enriched microalgae as biofertilizer.

Chapter 8

Table 8.1 Selenium uptake by duckweed, microalgae and sludge upon 0.5 mg Se/L exposure and Se utilization efficiency by beans grown in soil amended with the Se-enriched biomaterials.

	Se concentration in the biomass upon 0.5 mg Se/L exposure		Se utilization efficiency by beans grown in soil amended with Se-enriched biomaterials **	
	mg Se/kg DW biomass			
	Se(IV)	Se(VI)	%	
Duckweed	171	15.5	5.8% (DW1)	1.7% (DW5)
Microalgae	67	23	3.2% (5% biomass soil application) 4.1% (5% extract foliar application)	2.2% (10% biomass soil application) 2.3% (10% extract foliar application)
Sludge	82.1*	31.4*	5.8% (SL1)	4.5% (SL5)

Note:

* values estimated from the 5.0 mg Se/L exposure instead of 0.5 mg Se/L, as Se removal by sludge upon 0.5 mg Se/L exposure was not conducted.

** Se utilization efficiency = Se accumulation in plants of each pot (g)/ amount of Se applied in each pot (g)

Table 8.2 Comparison of growth conditions between duckweed and microalgae.

	Climate Conditions	Growth rate	Harvest	Resources	Space requirement
Duckweed	Wide variety	High	Easy	Proteins and starch	Floating on water surface, more growth space required
Microalgae	Warm	Very High	Very hard	Proteins and oil	Entire water phase, less space, but need extra energy to mix the biomass

8.4 Risks related to the use of Se-enriched biomaterials generated from wastewater as micronutrient biofertilizer and feed supplement

The main concerns of re-using the Se-enriched biomaterials produced on wastewater are the presence of heavy metals, pathogens, and organic (micro)pollutants. The biomaterials grown on wastewater can transfer these contaminants to the food chain when they are consumed as feed or applied on agricultural land. They may ultimately

Chapter 8

affect animal/human health and the natural environment. A complete risk assessment for use of these biomaterials should be conducted.

Potential heavy metal leaching and transfer to the food chain could occur when adding biomaterials from wastewater on agricultural land. The EU (1986 and 2003) has adopted the following limit values for heavy metals concentration in sludge for use in agriculture (expressed as mg/kg dry matter): Cu 1000-1750, Zn 2500-4000, Ni 300-400, Cr 500, Pb 750-1200, and Cd 20-40 (Cieřlik et al., 2015). The heavy metal concentrations in Se-enriched sludge used in our study were within these limits (Table 4.S1). Accordingly, the EU fertilizing products and amending regulation (EU, 2019) has also stipulated some contaminants in plant biostimulants and organic fertilizers (one of the product function categories of EU fertilizing products), such as heavy metals and pathogen loads (please see the exact values in Chapter 7). The levels of all heavy metals in the Se-enriched duckweed (Table 4.S1) and microalgae (Table 7.1) are well below these safety limits, except Cr in the microalgae (although the regulation stipulates Cr(VI), whereas we measured total Cr in our study). However, it should be re-emphasized that the Se-enriched duckweed and sludge produced in our study were generated from synthetic medium, instead of real wastewater, and thus the heavy metal concentrations are expected to differ if grown in real wastewater. Even for microalgae grown in real domestic wastewater, different types of wastewater would also result in different heavy metal contents. Nevertheless, these regulations and other regulations from worldwide should be always considered when valorizing recycled products from wastewater in agriculture. On the other hand, for feed supplements, there is no complete regulation yet controlling the use of algae as feed supplement in terms of heavy metals content. Currently, only thresholds for Pb, Hg and Cd in food supplements are regulated through the Commission Regulation 333/2007 (EC, 2007). The values of the three heavy metals in the Se-enriched microalgae (cultivation in real wastewater) and duckweed (cultivation in medium) biomass are within the limit levels. However, it should be noted that a feed supplement would be different from a food supplement and thresholds of these elements in food supplement can not completely be adopted to assess the toxic issues. Therefore, in this context, new regulations involving a wide range of heavy metal limitations should be therefore well established. Besides regulations, assessments of the chronic toxicity of these biomaterials as feed supplement are also essential.

Chapter 8

In terms of pathogen risk, strict sterilization could facilitate the reduction of most pathogens. This was demonstrated by the results reported in Chapter 7. However, a broader pathogenic species analysis is necessary to ensure the safety of using these bioproducts as biofertilizer and feed supplement. Organic (micro)pollutants refer to substances such as pharmaceutically active compounds, hormones, personal care products and detergents (Lam et al., 2020). Currently, there are no specific regulations globally limiting these organic pollutants in biomaterials for use as biofertilizer and feed supplement. Therefore, the establishment of new regulations in order to ensure the biosafety of bioproducts for use as fertilizer or feed supplement is urgently needed.

Except for the main contaminants potentially being present in wastewater, externally adding Se into the water for producing Se-enriched biomaterials could also introduce a potential risk for the environment and organisms. Particularly, duckweed and microalgae do not take up all applied Se. If the Se residues enter freshwaters, Se concentrations in the receiving water bodies will be elevated. Toxicity problems for aquatic organisms may be caused if the Se concentration exceeds the current threshold for aquatic life (e.g., 1.5 µg Se/L in lentic waters and 3.1 µg Se/L in lotic waters adopted by the United States Environmental Protection Agency) (USEAP, 2016). This should be carefully addressed and assessed in the future studies. Lastly, except for economic and environmental assessment, social acceptance for using the produced Se-enriched microalgae as biofertilizer and feed supplements still needs to be addressed.

8.5 Future perspectives

8.5.1 Production of micronutrient-rich bioproducts

The Se/Zn-enriched duckweed and sludge used as biofertilizer were generated from a micronutrient-enriched medium mimicking wastewater in this thesis. The physicochemical properties of real Se/Zn-contaminated water being used (partially purified wastewater) can be different from those of the Se/Zn-enriched media prepared in the laboratory. For instance, real wastewaters contain a certain amount of organic carbon and solids, while nutrient medium does not. Organic carbon could probably serve as a reducing agent, resulting in the change of redox potential, leading to the reduction of Se(VI) or Se(IV) in water. This could affect the Se uptake by duckweed. Besides, a high content of solids in wastewater could adsorb or precipitate Se(IV) or

Chapter 8

Se(VI), reducing bioavailable Se for duckweed uptake. High concentration of S, P or N (NO_3^- and NO_2^-) contained in wastewaters would also significantly reduce Se accumulation in duckweed/sludge. These differences could possibly lead to producing bioproducts having different physicochemical properties and different Se uptake. Thus, the production of Se/Zn-enriched duckweed and sludge under real wastewater conditions needs to be further confirmed. In this context, potential environmental and health risks related to the possible presence of heavy metals, pathogen loads, and organic micropollutants in the bioproducts generated from waste should be particularly addressed.

Besides, taking into account the low Se removal efficiency of duckweed and microalgae, the enhancement of Se accumulation by duckweed and microalgae is still needed in future studies. This could be achieved through: (1) optimization of growth conditions (e.g. optimal temperature and light); (2) screening specific duckweed/microalgae species with higher Se accumulation ability; (3) optimization of the wastewater treatment process, such as conducting pretreatment processes aimed to remove other components (e.g. S and P), which could interfere with the Se uptake; (4) inoculation microalgae/duckweed with specific bacteria facilitating Se removal; and (5) genetic manipulation for enhancing Se accumulation or transformation after being taken up by duckweed/microalgae, such as transgenic methods. Apart from the use of real Se/Zn-enriched wastewater, wastewater with low Se/Zn content (e.g. domestic wastewater) could also be considered to act as nutrients source to produce the bioproducts while spiking Se and Zn. Additionally, pilot studies are also recommended for upscaling and verifying the results obtained in our study, in particular for duckweed, as the performance of the plants is expected to be different under different conditions (i.e. different geometry of growth tanks, different biomass density, and different ambient conditions).

In chapter 6, we did not perform replications of the Se-HRAPs due to the limitation of the facility (two HRAPs) and we did not analyze nutritional parameters of biomass produced at different time points. Accordingly, some findings in our study, e.g. Se inducing more omega-3 ($\omega 3$) and omega-6 ($\omega 6$) formation, require additional data collection for further validation. Thus, analysis of more biomass samples or more replications of the HRAPs (if applicable) are suggested for future studies.

Chapter 8

Additionally, although the potential of the Se/Zn-enriched duckweed and microalgae as value-added feed supplements was confirmed by measuring different nutritional parameters (i.e. amino acids, fatty acids, and organic Se species), the digestibility in animals (pig) assessed through *in vitro* studies may still be complemented by *in vivo* studies as the real bioavailability after ingestion may still be different in living animals.

8.5.2 Biofortification

Along with the incubation time, the considerable decrease of Se or Zn concentration in the pore water of soils amended with Se/Zn-enriched duckweed and microalgae was observed in this thesis. However, the explanation of these results is based on crop yield and agronomic parameters, such as absorption, transformation between different Se/Zn species in the soil matrix, Se and Zn uptake and translocation of Se and Zn within the plant tissues. The specific mechanisms affecting the availability of the released Se and Zn from Se/Zn-enriched biomaterials in soils should still be explored and verified in future studies. This could include measurement of Se/Zn in different fractions, e.g., the exchangeable, Fe/Mn oxide-bound, and organic matter-bound fractions of soils fertilized with Se/Zn-enriched biomaterials. This could contribute to understanding the dynamics, fate, and transformation of Se/Zn in the soil after being released from these biomaterials and to assessing the possible factors affecting the plant uptake.

Micronutrient-enriched duckweed was less efficient in increasing the Se/Zn concentrations in beans and even inhibited the plant growth at high application doses, compared to the micronutrient-enriched sludge (Chapter 4 and 5). Moreover, the Se/Zn in the soil pore water after being released from the duckweed was not stable in the first three weeks after the soil application. All these results indicate that stabilization of the produced micronutrient-enriched duckweed is definitely needed before being added to soils as a biofertilizer, contributing to improving the utilization efficiency of the trace elements. Post-treatment of the Se/Zn-enriched duckweed (such as composting and anaerobic digestion) after being harvested from wastewater is also needed to reduce its organic carbon content and phytotoxic substances (i.e. low-molecular weight organic acids), consequently mitigating its effects on plant growth at higher application doses.

Chapter 8

Besides, a long term experiment with continuous planting should be performed in soils amended with micronutrient-enriched bioproducts to further investigate the evolution of Se/Zn in the soil pore water and its effects on the crops and soil environment on the longer term. The potential leaching and retaining of Se/Zn in the soil with micronutrient-enriched biomaterials application should be further investigated on a longer-term. Besides, the effects of the fertilization of Se/Zn-enriched biomaterials on soil quality including microorganism activities are still needed. Finally, a field trial study with supplementation of the micronutrient-enriched bioproducts is recommended to compare and validate the results of the lab study on a larger scale.

In the biostimulant experiment, the extraction efficiency of Se from microalgae biomass was not found very high, only 15.4%. Improvement of the Se extraction efficiency from the Se-enriched microalgae is therefore needed in future studies, e.g. by using an alternative, unharmed mechanical process or organic solvents or enzymes to completely disrupt the microalgae cell wall. The combination of different methods, such as the combination of enzymatic and mechanical techniques, may also be explored.

8.5.3 Feasibility analysis

A thorough analysis of economic, environmental and social impacts, including safety and health issues, for the production and use of the micronutrient-enriched biomaterials with wastewater as feedstock is still needed. Health-related risk from pathogen loads and the potential presence of heavy metals and micropollutants should be carefully evaluated on the long term and compared with the legal frameworks of fertilizer legislation. Conducting a complete life cycle assessment (LCA) would be beneficial. Moreover, a cost-benefit analysis has to be done based on the results of pilot studies to evaluate the profit margin of production (recovery), processing, and use of these micronutrient-enriched bioproducts produced from wastes. The market potential of these new micronutrient-enriched bioproducts should still be assessed.

References

References

- Ajwa, H.A., Bañuelos, G.S., Mayland, H.F. 1998. Selenium uptake by plants from soils amended with inorganic and organic materials. *Journal of Environment Quality*, **27**(5), 1218-1227.
- Antonkiewicz, J., Poplawska, A., Kolodziej, B., Ciarkowska, K., Gambus, F., Bryk, M., Babula, J. 2020. Application of ash and municipal sewage sludge as macronutrient sources in sustainable plant biomass production. *Journal of Environmental Management*, **264**, 110450.
- APHA-AWWA-WEF. 2012. *Standard methods for the examination of water and wastewater*. 22nd ed. American Public Health Association, Washington, D.C.
- Arashiro, L.T., Boto-Ordóñez, M., Van Hulle, S.W.H., Ferrer, I., Garfí, M., Rousseau, D.P.L. 2020a. Natural pigments from microalgae grown in industrial wastewater. *Bioresource Technology*, **303**, 122894.
- Arashiro, L.T., Ferrer, I., Pániker, C.C., Gómez-Pinchetti, J.L., Rousseau, D.P.L., Van Hulle, S.W.H., Garfí, M. 2020b. Natural Pigments and Biogas Recovery from Microalgae Grown in Wastewater. *ACS Sustainable Chemistry & Engineering*, **8**(29), 10691-10701.
- Arashiro, L.T., Ferrer, I., Rousseau, D.P.L., Van Hulle, S.W.H., Garfí, M. 2019. The effect of primary treatment of wastewater in high rate algal pond systems: Biomass and bioenergy recovery. *Bioresource Technology*, **280**, 27-36.
- Arbestain, M.C. 2001. Modeling selenium transfers in straw-amended soils. *Soil Science*, **166**(8), 539-547.
- Aro, A., Alfthan, G., Varo, P. 1995. Effects of supplementation of fertilizers on human selenium status in Finland. *The Analyst*, **120**(3), 841-843.
- Arvy, M.P. 1993. Selenate and selenite uptake and translocation in bean-plants (*Phaseolus vulgaris*). *Journal of Experimental Botany*, **44**(6), 1083-1087.
- Arvy, M.P. 1992. Some aspects of selenium relationships in soils and plants. *Communications in Soil Science and Plant Analysis*, **23**(13-14), 1397-1407.
- Azhar u, d., Huang, J.-C., Gan, X., He, S., Zhou, W. 2020. Interactive effects of earthworm *Eisenia fetida* and bean plant *Phaseolus vulgaris* L on the fate of soil selenium. *Environmental Pollution*, **260**, 114048.

References

- Bañuelos, G.S., Arroyo, I., Pickering, I.J., Yang, S.I., Freeman, J.L. 2015. Selenium biofortification of broccoli and carrots grown in soil amended with Se-enriched hyperaccumulator *Stanleya pinnata*. *Food Chemistry*, **166**, 603-8.
- Bañuelos, G.S., Arroyo, I.S., Dangi, S.R., Zambrano, M.C. 2016. Continued selenium biofortification of carrots and broccoli grown in soils once amended with Se-enriched *S. pinnata*. *Frontiers in Plant Science*, **7**, 1251.
- Bañuelos, G.S., Hanson, B.D. 2010. Use of selenium-enriched mustard and canola seed meals as potential bioherbicides and green fertilizer in strawberry production. *Hortscience*, **45**(10), 1567-1572.
- Bañuelos, G.S., Lin, Z.-Q., Broadley, M. 2017. Selenium biofortification. in: *Selenium in plants: Molecular, Physiological, Ecological and Evolutionary Aspects*, (Eds.) E.A.H. Pilon-Smits, L.H.E. Winkel, Z.-Q. Lin, Springer International Publishing. Cham, pp. 231-255.
- Batish, D.R., Lavanya, K., Singh, H.P., Kohli, R.K. 2007. Phenolic allelochemicals released by *Chenopodium murale* affect the growth, nodulation and macromolecule content in chickpea and pea. *Plant Growth Regulation*, **51**(2), 119-128.
- Becker, E.W. 2013. Microalgae for human and animal nutrition. Second Edition ed. in: *Handbook of Microalgal Culture*, (Eds.) A. Richmond, Emeritus, Q. Hu, John Wiley & Sons, Ltd., pp. 461-503.
- Benlloch, M., Ojeda, M.A., Ramos, J., Rodriguez-Navarro, A. 1994. Salt sensitivity and low discrimination between potassium and sodium in bean plants. *Plant and Soil*, **166**(1), 117-123.
- Bingham, R.L.M.A.L.P.F.T. 1989. Factors affecting selenium accumulation by agricultural crops. in: *Selenium in agriculture and the environment*, pp. 65-94.
- Boldrin, P.F., Faquin, V., Ramos, S.J., Boldrin, K.V.F., Ávila, F.W., Guilherme, L.R.G. 2013. Soil and foliar application of selenium in rice biofortification. *Journal of Food Composition and Analysis*, **31**(2), 238-244.
- Borowitzka, M.A., Moheimani, N.R. 2013. Sustainable biofuels from algae. *Mitigation and Adaptation Strategies for Global Change*, **18**(1), 13-25.
- Broadley, M.R., White, P.J., Bryson, R.J., Meacham, M.C., Bowen, H.C., Johnson, S.E., Hawkesford, M.J., McGrath, S.P., Zhao, F.J., Breward, N., Harriman, M.,

References

- Tucker, M. 2006. Biofortification of UK food crops with selenium. *Proceedings of the Nutrition Society*, **65**(2), 169-81.
- Broadley, M.R., White, P.J., Hammond, J.P., Zelko, I., Lux, A. 2007. Zinc in plants. *New Phytologist*, **173**(4), 677-702.
- Brouwer, P., Nierop, K.G.J., Huijgen, W.J.J., Schluepmann, H. 2019. Aquatic weeds as novel protein sources: Alkaline extraction of tannin-rich *Azolla*. *Biotechnology Reports*, **24**, e00368.
- Brouwer, P., Schluepmann, H., Nierop, K.G., Elderson, J., Bijl, P.K., van der Meer, I., de Visser, W., Reichart, G.J., Smeekens, S., van der Werf, A. 2018. Growing *Azolla* to produce sustainable protein feed: the effect of differing species and CO₂ concentrations on biomass productivity and chemical composition. *Journal of the Science of Food and Agriculture*, **98**(12), 4759-4768.
- Broyer, T.C., Johnson, C.M., Huston, R.P. 1972. Selenium and nutrition of *Astragalus*: I. Effects of selenite or selenate supply on growth and growth selenium content. *Plant and Soil*, **36**(3), 635-649.
- Calvo, P., Nelson, L., Kloepper, J.W. 2014. Agricultural uses of plant biostimulants. *Plant and Soil*, **383**(1-2), 3-41.
- Carvalho, K.M., Martin, D.F. 2001. Removal of aqueous selenium by four aquatic plants. *Journal of Aquatic Plant Management*, **39**, 33-36.
- Čertík, M., Breierová, E., Oláhová, M., Šajbidor, J., Márová, I. 2013. Effect of selenium on lipid alternations in pigment-forming yeasts. *Food Science and Biotechnology*, **22**(1), 45-51.
- CH2MHill. 2010. Review of available technologies for removal of selenium from water. *Final report, prepared for North American Metals Council (NAMC)*.
- Chen, G., Li, X.D., Picot, B., Huang, X.F., He, X.J., He, S.L., Zhou, Q. 2006. Nutrients removal mechanisms in high rate algal pond treating rural domestic sewage in East China. *Water Supply*, **6**(6), 43-50.
- Chen, Z., Shao, S., He, Y., Luo, Q., Zheng, M., Zheng, M., Chen, B., Wang, M. 2020. Nutrients removal from piggery wastewater coupled to lipid production by a newly isolated self-flocculating microalga *Desmodesmus* sp. PW1. *Bioresource Technology*, **302**, 122806.

References

- Chiaiese, P., Corrado, G., Colla, G., Kyriacou, M.C., Rouphael, Y. 2018. Renewable sources of plant biostimulation: Microalgae as a sustainable means to improve crop performance. *Frontiers in Plant Science*, **9**, 1782.
- Cieřlik, B.M., Namieřnik, J., Konieczka, P. 2015. Review of sewage sludge management: standards, regulations and analytical methods. *Journal of Cleaner Production*, **90**, 1-15.
- Combs, G.F., Jr. 2001. Selenium in global food systems. *British Journal of Nutrition*, **85**(5), 517-47.
- Coppens, J., Grunert, O., Van Den Hende, S., Vanhoutte, I., Boon, N., Haesaert, G., De Gelder, L. 2015. The use of microalgae as a high-value organic slow-release fertilizer results in tomatoes with increased carotenoid and sugar levels. *Journal of Applied Phycology*, **28**(4), 2367-2377.
- Coppin, F., Chabroulet, C., Martin-Garin, A., Balesdent, J., Gaudet, J.P. 2006. Methodological approach to assess the effect of soil ageing on selenium behaviour: first results concerning mobility and solid fractionation of selenium. *Biology and Fertility of Soils*, **42**(5), 379-386.
- Council, N.R. 2005. *Mineral Tolerance of Animals: Second Revised Edition, 2005*. The National Academies Press, Washington, DC.
- Darago, A., Sapota, A., Nasiadek, M., Klimczak, M., Kilanowicz, A. 2016. The effect of zinc and selenium supplementation mode on their bioavailability in the rat prostate. Should administration be joint or separate? *Nutrients*, **8**(10).
- de Oliveira, A.P., Nomura, C.S., Naozuka, J. 2017. Evaluation of selenium enrichment of adzuki bean (*Vigna angularis*) sprouts: Translocation, bioaccessibility and Se-protein speciation. *Microchemical Journal*, **134**, 19-26.
- Delgadillo-Mirquez, L., Lopes, F., Taidi, B., Pareau, D. 2016. Nitrogen and phosphate removal from wastewater with a mixed microalgae and bacteria culture. *Biotechnology Report*, **11**, 18-26.
- Deng, X., Zhao, Z., Lv, C., Zhang, Z., Yuan, L., Liu, X. 2021. Effects of sulfur application on selenium uptake and seed selenium speciation in soybean (*Glycine max* L.) grown in different soil types. *Ecotoxicology and Environmental Safety*, **209**, 111790.

References

- Dermauw, V., Yisehak, K., Belay, D., Van Hecke, T., Du Laing, G., Duchateau, L., Janssens, G.P.J. 2013. Mineral deficiency status of ranging zebu (*Bos indicus*) cattle around the Gilgel Gibe catchment, Ethiopia. *Tropical Animal Health and Production*, **45**(5), 1139-1147.
- Dhillon, S.K., Hundal, B.K., Dhillon, K.S. 2007. Bioavailability of selenium to forage crops in a sandy loam soil amended with Se-rich plant materials. *Chemosphere*, **66**(9), 1734-4173.
- Di Tullo, P., Versini, A., Bueno, M., Le Hécho, I., Thiry, Y., Biron, P., Castrec-Rouelle, M., Pannier, F. 2015. Stable isotope tracing: a powerful tool for selenium speciation and metabolic studies in non-hyperaccumulator plants (*ryegrass Lolium perenne* L.). *Analytical and Bioanalytical Chemistry*, **407**(30), 9029-9042.
- Dias, G.A., Rocha, R.H.C., Araújo, J.L., Lima, J.F., Guedes, W.A. 2016. Growth, yield, and postharvest quality in eggplant produced under different foliar fertilizer (*Spirulina platensis*) treatments. *Semina: Ciências Agrárias*, **37**(6), 3893.
- Díaz-Alarcón, J.P., Navarro-Alarcón, M., López-García de la Serrana, H., López-Martínez, M.C. 1996. Determination of selenium in cereals, legumes and dry fruits from southeastern Spain for calculation of daily dietary intake. *Science of The Total Environment*, **184**(3), 183-189.
- Dinh, Q.T., Cui, Z., Huang, J., Tran, T.A.T., Wang, D., Yang, W., Zhou, F., Wang, M., Yu, D., Liang, D. 2018. Selenium distribution in the Chinese environment and its relationship with human health: A review. *Environment International*, **112**, 294-309.
- Dinh, Q.T., Zhou, F., Wang, M., Peng, Q., Wang, M., Qi, M., Tran, T.A.T., Chen, H., Liang, D. 2020. Assessing the potential availability of selenium in the soil-plant system with manure application using diffusive gradients in thin-films technique (DGT) and DOM-Se fractions extracted by selective extractions. *Science of The Total Environment*, 143047.
- Directive, C. 1986. Council Directive 86/278/EEC of 12 June 1986 on the protection of the environment, and in particular of the soil, when sewage sludge is used in agriculture.
- Doughman, S.D., Krupanidhi, S., Sanjeevi, C.B. 2007. Omega-3 fatty acids for nutrition and medicine: considering microalgae oil as a vegetarian source of EPA and DHA. *Current Diabetes Reviews*, **3**(3), 198-203.

References

- Dong, Z., Liu, Y., Dong, G., Wu, H. 2021. Effect of boiling and frying on the selenium content, speciation, and in vitro bioaccessibility of selenium-biofortified potato (*Solanum tuberosum* L.). *Food Chemistry*, **348**, 129150.
- Duan, L.Q., Song, J.M., Li, X.G., Yuan, H.M., Xu, S.S. 2010. Distribution of selenium and its relationship to the eco-environment in Bohai Bay seawater. *Marine Chemistry*, **121**(1-4), 87-99.
- Dubois, M., Gilles, K., Hamilton, J.K., Rebers, P.A., Smith, F. 1951. A Colorimetric Method for the Determination of Sugars. *Nature*, **168**(4265), 167-167.
- Ebrahimi, N., Hartikainen, H., Hajiboland, R., Seppänen, M.M. 2019. Uptake and remobilization of selenium in *Brassica napus* L. plants supplied with selenate or selenium-enriched plant residues. *Journal of Plant Nutrition and Soil Science*, **182**(2), 196-202.
- EC. 2007. Commission Regulation (EC) No 333/2007 of 28 March 2007 laying down the methods of sampling and analysis for the official control of the levels of lead, cadmium, mercury, inorganic tin, 3-MCPD and benzo(a)pyrene in foodstuffs (Text with EEA relevance).
- EC. 2003. Regulation (EC) No 2003/2003 of the European Parliament and of the Council of 13 October 2003 relating to fertilisers (Text with EEA relevance).
- EFSA. Chronic Food Consumption Grams per kilogram of body weight per day (g/kg bw per day)-consumers only.
<https://www.efsa.europa.eu/en/microstrategy/foodex2-level-1>.
- EFSA. 2016. Safety and efficacy of selenium compounds (E8) as feed additives for all animal species: Sodium selenite, based on a dossier submitted by Todini and Co SpA. *EFSA Journal*, **14**(3).
- EFSA. 2014. Scientific opinion on dietary reference values for selenium. *EFSA Journal*, **12**(10), 3846.
- Egene, C.E., Van Poucke, R., Ok, Y.S., Meers, E., Tack, F.M.G. 2018. Impact of organic amendments (biochar, compost and peat) on Cd and Zn mobility and solubility in contaminated soil of the Campine region after three years. *Science of the Total Environment*, **626**, 195-202.

References

- Eich-Greatorex, S., Sogn, T.A., Øgaard, A.F., Aasen, I. 2007. Plant availability of inorganic and organic selenium fertiliser as influenced by soil organic matter content and pH. *Nutrient Cycling in Agroecosystems*, **79**(3), 221-231.
- Eiche, E., Bardelli, F., Nothstein, A.K., Charlet, L., Gottlicher, J., Steininger, R., Dhillon, K.S., Sadana, U.S. 2015. Selenium distribution and speciation in plant parts of wheat (*Triticum aestivum*) and Indian mustard (*Brassica juncea*) from a seleniferous area of Punjab, India. *Science of the Total Environment*, **505**, 952-61.
- El-Ramady, H., Faizy, S.E.D., Abdalla, N., Taha, H., Domokos-Szabolcsy, É., Fari, M., Elsakhawy, T., Omara, A.E.-D., Shalaby, T., Bayoumi, Y., Shehata, S., Geilfus, C.-M., Brevik, E.C. 2020. Selenium and nano-selenium biofortification for human health: Opportunities and challenges. *Soil Systems*, **4**(3), 57.
- Ellis, R.H., Roberts, E.H. 1981. The quantification of aging and survival in orthodox seeds. *Seed Science and Technology*, **9**(2), 373-410.
- Etteieb, S., Magdouli, S., Zolfaghari, M., Brar, S. 2020. Monitoring and analysis of selenium as an emerging contaminant in mining industry: A critical review. *Science of the Total Environment*, **698**, 134339.
- EU. 2019. Regulation (EU) 2019/1009 of the European Parliament and of the Council of 5 June 2019 laying down rules on the making available on the market of EU fertilising products and amending Regulations (EC) No 1069/2009 and (EC) No 1107/2009 and repealing Regulation (EC) No 2003/2003.
- Faheed, F.A., Abd-El Fattah, Z. 2008. Effect of *Chlorella vulgaris* as biofertilizer on growth parameters and metabolic aspects of lettuce plant. *Journal of Agriculture & Social Science*, **4**.
- FAO/WHO. 2001. Human Vitamin and Mineral Requirements. *Report of a joint FAO/WHO expert consultation, Bangkok, Thailand*.
- Feng, R., Wei, C., Tu, S. 2013. The roles of selenium in protecting plants against abiotic stresses. *Environmental and Experimental Botany*, **87**, 58-68.
- Feng, R.W., Wei, C.Y., Tu, S.X., Wu, F.C. 2009. Effects of Se on the uptake of essential elements in *Pteris vittata* L. *Plant and Soil*, **325**(1-2), 123-132.

References

- Fernández-Martínez, A., Charlet, L. 2009. Selenium environmental cycling and bioavailability: a structural chemist point of view. *Reviews in Environmental Science and Bio/Technology*, **8**(1), 81-110.
- Filek, M., Gzyl-Malcher, B., Zembala, M., Bednarska, E., Laggner, P., Kriechbaum, M. 2010. Effect of selenium on characteristics of rape chloroplasts modified by cadmium. *Journal of Plant Physiology*, **167**(1), 28-33.
- Folch, J., Lees, M., Sloane Stanley, G.H. 1957. A simple method for the isolation and purification of total lipides from animal tissues. *The Journal of Biological Chemistry*, **226**(1), 497-509.
- Fontes, R.L.F., Pereira, J.M.N., Neves, J.C.L. 2014. Uptake and translocation of Cd and Zn in two lettuce cultivars. *Anais da Academia Brasileira de Ciências*, **86**(2), 907-922.
- Food, E.S.C.o. 2003. Opinion of the Scientific Committee on Food on the Revision of Reference Values for Nutrition Labelling, (Ed.) C.o.t.E. Communities. Brussels.
- Fordyce, F. 2007. Selenium geochemistry and health. *Ambio*, **36**(1).
- Fordyce, F.M. 2013. Selenium deficiency and toxicity in the environment. in: *Essentials of Medical Geology: Revised Edition*, (Ed.) O. Selinus, Springer Netherlands. Dordrecht, pp. 375-416.
- Fordyce, F.M., Guangdi, Z., Green, K., Xinping, L. 2000. Soil, grain and water chemistry in relation to human selenium-responsive diseases in Enshi District, China. *Applied Geochemistry*, **15**(1), 117-132.
- Gan, X., Huang, J.C., Zhou, C., He, S., Zhou, W. 2019. Relationship between selenium removal efficiency and production of lipid and hydrogen by *Chlorella vulgaris*. *Chemosphere*, **217**, 825-832.
- García, J., Ortiz, A., Álvarez, E., Belohlav, V., García-Galán, M.J., Díez-Montero, R., Álvarez, J.A., Uggetti, E. 2018. Nutrient removal from agricultural run-off in demonstrative full scale tubular photobioreactors for microalgae growth. *Ecological Engineering*, **120**, 513-521.
- Garcia-Gonzalez, J., Sommerfeld, M. 2016. Biofertilizer and biostimulant properties of the microalga *Acutodesmus dimorphus*. *Journal of Applied Phycology*, **28**(2), 1051-1061.

References

- Garousi, F., Veres, S., Kovacs, B. 2016. Comparison of selenium toxicity in sunflower and maize seedlings grown in hydroponic cultures. *Bulletin of Environmental Contamination and Toxicology*, **97**(5), 709-713.
- Gerhardt, M.B., Green, F.B., Newman, R.D., Lundquist, T.J., Tresan, R.B., Oswald, W.J. 1991. Removal of selenium using a novel algal-bacterial process. *Research Journal of the Water Pollution Control Federation*, **63**(5), 799-805.
- Gibson, R.S. 2006. Zinc: the missing link in combating micronutrient malnutrition in developing countries. *The Proceedings of the Nutrition Society*, **65**(1), 51-60.
- Gómez-Jacinto, V., Navarro-Roldán, F., Garbayo-Nores, I., Vílchez-Lobato, C., Borrego, A.A., García-Barrera, T. 2020. In vitro selenium bioaccessibility combined with in vivo bioavailability and bioactivity in Se-enriched microalga (*Chlorella sorokiniana*) to be used as functional food. *Journal of Functional Foods*, **66**, 103817.
- Grzesik, M., Romanowska-Duda, Z. 2014. Improvements in germination, growth, and metabolic activity of corn seedlings by grain conditioning and root application with cyanobacteria and microalgae. *Polish Journal of Environmental Studies*, **23**(4), 1147-1153.
- Guerrero, B., Llugany, M., Palacios, O., Valiente, M. 2014. Dual effects of different selenium species on wheat. *Plant Physiology and Biochemistry*, **83**, 300-307.
- Gupta, A.B., Shukla, A.C. 1969. Effect of algal extracts of *Phormidium* species on growth and development of rice seedlings. *Hydrobiologia*, **34**(1), 77-84.
- Gupta, M., Gupta, S. 2017. An overview of selenium uptake, metabolism, and toxicity in plants. *Frontiers in Plant Science*, **7**(2074).
- Gutierrez, R., Ferrer, I., Gonzalez-Molina, A., Salvado, H., Garcia, J., Uggetti, E. 2016. Microalgae recycling improves biomass recovery from wastewater treatment high rate algal ponds. *Water Research*, **106**, 539-549.
- Han, W., Mao, Y., Wei, Y., Shang, P., Zhou, X. 2020. Bioremediation of aquaculture wastewater with algal-bacterial biofilm combined with the production of selenium rich biofertilizer. *Water*, **12**(7), 2071.
- Handa, N., Bhardwaj, R., Kaur, H., Poonam, Kapoor, D., Rattan, A., Kaur, S., Thukral, A.K., Kaur, S., Arora, S., Kapoor, N. 2016. Chapter 7 - Selenium: An

References

- antioxidative protectant in plants under stress. in: *Plant Metal Interaction*, (Ed.) P. Ahmad, Elsevier, pp. 179-207.
- Hassan, A.M.A., Mostafa, E.M. 2016. Selenium invoked antioxidant defense system in *Azolla caroliniana* plant. *Phyton-International Journal of Experimental Botany*, **85**, 262-269.
- Hatfield, D.L., Tsuji, P.A., Carlson, B.A., Gladyshev, V.N. 2014. Selenium and selenocysteine: roles in cancer, health, and development. *Trends in Biochemical Sciences*, **39**(3), 112-120.
- Haug, A., Graham, R.D., Christophersen, O.A., Lyons, G.H. 2007. How to use the world's scarce selenium resources efficiently to increase the selenium concentration in food. *Microbial Ecology in Health and Disease*, **19**(4), 209-228.
- Hawrylak-Nowak, B., Dresler, S., Wójcik, M. 2014. Selenium affects physiological parameters and phytochelatins accumulation in cucumber (*Cucumis sativus* L.) plants grown under cadmium exposure. *Scientia Horticulturae*, **172**, 10–18.
- He, Z.L., Shentu, J., Yang, X.E. 2010. Manganese and selenium. in: *Trace Elements in Soils*, pp. 481-495.
- Health, M.o. 2018. Kenya national food fortification strategic plan: 2018-2022., (Ed.) R.o.K.-M.o. Health.
- Hernández-Herrera, R.M., Santacruz-Ruvalcaba, F., Ruiz-López, M.A., Norrie, J., Hernández-Carmona, G. 2013. Effect of liquid seaweed extracts on growth of tomato seedlings (*Solanum lycopersicum* L.). *Journal of Applied Phycology*, **26**(1), 619-628.
- Hoffmann, J.P. 1998. Wastewater treatment with suspended and nonsuspended algae. *Journal of Phycology*, **34**, 757-763.
- Hu, T., Li, H., Li, J., Zhao, G., Wu, W., Liu, L., Wang, Q., Guo, Y. 2018. Absorption and bio-transformation of selenium nanoparticles by wheat seedlings (*Triticum aestivum* L.). *Frontiers in Plant Science*, **9**, 597.
- Huang, Y., Wang, Q., Gao, J., Lin, Z., Bañuelos, G.S., Yuan, L., Yin, X. 2013. Daily dietary selenium intake in a high selenium area of Enshi, China. *Nutrients*, **5**(3), 700-710.
- Huang, Y.Z., Hu, Y., Liu, Y.X. 2008. Interactions between sulfur and selenium uptake by corn in solution culture. *Journal of Plant Nutrition*, **31**(1), 43-54.

References

- IRIS, U.-E. 2006. Selenium and Compounds; CASRN 7782-49-2. Intergrated risk information system (IRIS) program. *U.S. Environmental Protection Agency (EPA)*.
- Jin, Z., Shah, T., Zhang, L., Liu, H., Peng, S., Nie, L. 2020. Effect of straw returning on soil organic carbon in rice–wheat rotation system: A review. *Food and Energy Security*, **9**(2).
- Johnson, C.C., Fordyce, F.M., Rayman, M.P. 2009. Symposium on 'Geographical and geological influences on nutrition' Factors controlling the distribution of selenium in the environment and their impact on health and nutrition: Conference on 'Over- and undernutrition: challenges and approaches'. *Proceedings of the Nutrition Society*, **69**(1), 119-132.
- Joy, E.J., Ander, E.L., Young, S.D., Black, C.R., Watts, M.J., Chilimba, A.D., Chilima, B., Siyame, E.W., Kalimira, A.A., Hurst, R., Fairweather-Tait, S.J., Stein, A.J., Gibson, R.S., White, P.J., Broadley, M.R. 2014. Dietary mineral supplies in Africa. *Physiol Plant*, **151**(3), 208-29.
- Jozwiak, W., Politycka, B. 2019. Effect of selenium on alleviating oxidative stress caused by a water deficit in cucumber roots. *Plants (Basel)*, **8**(7), 217.
- Kaveh, H., Nemati, H., Farsi, M., Jartoodeh, S.V. 2011. How salinity affect germination and emergence of tomato lines. *Journal of Biological and Environmental Science*, **5**(15), 159-163.
- Khan, S.A., Sharma, G.K., Malla, F.A., Kumar, A., Rashmi, Gupta, N. 2019. Microalgae based biofertilizers: A biorefinery approach to phycoremediate wastewater and harvest biodiesel and manure. *Journal of Cleaner Production*, **211**, 1412-1419.
- Kieliszek, M., Błażej, S., Bzducha-Wróbel, A., Kot, A.M. 2019. Effect of selenium on lipid and amino acid metabolism in yeast cells. *Biological Trace Element Research*, **187**(1), 316-327.
- Kikkert, J., Berkelaar, E. 2013. Plant uptake and translocation of inorganic and organic forms of selenium. *Archives of Environmental Contamination and Toxicology*, **65**(3), 458-465.
- Kimbrough, D.E., Cohen, Y., Winer, A.M., Creelman, L., Mabuni, C. 1999. A critical assessment of chromium in the environment. *Critical Reviews in Environmental Science and Technology*, **29**(1), 1-46.

References

- Kjeldahl, J. 1883. A new method for the determination of nitrogen in organic matter - Neue Methode zur Bestimmung des Stickstoffs in organischen Körpern. *Zeitschrift für analytische Chemie*, **22**(1), 366-382.
- Kumar, G., Sahoo, D. 2011. Effect of seaweed liquid extract on growth and yield of *Triticum aestivum* var. Pusa Gold. *Journal of Applied Phycology*, **23**(2), 251-255.
- Lam, K.L., Zlatanovic, L., van der Hoek, J.P. 2020. Life cycle assessment of nutrient recycling from wastewater: A critical review. *Water Research*, **173**, 115519.
- Lavu, R.V.S., De Schepper, V., Steppe, K., Majeti, P.N.V., Tack, F., Du Laing, G. 2013. Use of selenium fertilizers for production of Se-enriched Kenaf (*Hibiscus cannabinus*): Effect on Se concentration and plant productivity. *Journal of Plant Nutrition and Soil Science*, **176**(4), 634-639.
- Lavu, R.V.S., Du Laing, G., Van de Wiele, T., Pratti, V.L., Willekens, K., Vandecasteele, B., Tack, F. 2012. Fertilizing soil with selenium fertilizers: impact on concentration, speciation, and bioaccessibility of selenium in leek (*Allium ampeloprasum*). *Journal of Agricultural and Food Chemistry*, **60**(44), 10930-10935.
- Lavu, R.V.S., Van De Wiele, T., Pratti, V.L., Tack, F., Du Laing, G. 2016. Selenium bioaccessibility in stomach, small intestine and colon: Comparison between pure Se compounds, Se-enriched food crops and food supplements. *Food Chemistry*, **197**, 382-387.
- Lemly, A.D. 1997. Environmental implications of excessive selenium: A review. *Biomedical and Environmental Sciences*, **10**(4), 415-435.
- Lenz, M., Hullebusch, E.D.v., Farges, F., Nikitenko, S., Borca, C.N., Grolimund, D., Lens, P.N.L. 2008. Selenium speciation assessed by X-Ray absorption spectroscopy of sequentially extracted anaerobic biofilms. *Environmental Science & Technology*, **42**(20), 7587-7593.
- Lenz, M., Lens, P.N. 2009. The essential toxin: the changing perception of selenium in environmental sciences. *Sci Total Environ*, **407**(12), 3620-33.
- Li, H.F., Lombi, E., Stroud, J.L., McGrath, S.P., Zhao, F.J. 2010. Selenium speciation in soil and rice: influence of water management and Se fertilization. *Journal of Agricultural and Food Chemistry*, **58**(22), 11837-11843.

References

- Li, H.F., McGrath, S.P., Zhao, F.J. 2008. Selenium uptake, translocation and speciation in wheat supplied with selenate or selenite. *New Phytologist*, **178**(1), 92-102.
- Li, J., Ferrer, I., Gijs Du, L., Otero-Gonzalez, L., Loi, G., Lens, P.N.L. 2020a. Selenate and selenite uptake, accumulation and toxicity in *Lemna minuta*. *Water Science and Technology*, **81**(9), 1852-1862.
- Li, J., Lens, P.N.L., Otero-Gonzalez, L., Du Laing, G. 2020b. Production of selenium- and zinc-enriched *Lemna* and *Azolla* as potential micronutrient-enriched bioproducts. *Water Research*, **172**, 115522.
- Li, J., Liang, D., Qin, S., Feng, P., Wu, X. 2015a. Effects of selenite and selenate application on growth and shoot selenium accumulation of pak choi (*Brassica chinensis* L.) during successive planting conditions. *Environment Science and Pollution Research*, **22**(14), 11076-86.
- Li, J., Peng, Q., Liang, D., Liang, S., Chen, J., Sun, H., Li, S., Lei, P. 2016. Effects of aging on the fraction distribution and bioavailability of selenium in three different soils. *Chemosphere*, **144**, 2351-2359.
- Li, X., Li, B.Q., Yang, Y.P. 2018. Effects of foliar selenite on the nutrient components of turnip (*Brassica rapa* var. *rapa* Linn.). *Frontiers in Chemistry*, **6**.
- Li, Z., Liang, D., Peng, Q., Cui, Z., Huang, J., Lin, Z. 2017. Interaction between selenium and soil organic matter and its impact on soil selenium bioavailability: A review. *Geoderma*, **295**, 69-79.
- Li, Z., Man, N., Wang, S., Liang, D., Liu, J. 2015b. Selenite adsorption and desorption in main Chinese soils with their characteristics and physicochemical properties. *Journal of Soils and Sediments*, **15**(5), 1150-1158.
- Li, Z.-Y., Guo, S.-Y., Li, L. 2003. Bioeffects of selenite on the growth of *Spirulina platensis* and its biotransformation. *Bioresource Technology*, **89**(2), 171-176.
- Lim, T.T., Goh, K.H. 2005. Selenium extractability from a contaminated fine soil fraction: implication on soil cleanup. *Chemosphere*, **58**(1), 91-101.
- Liu, F., Huang, J.C., Zhou, C., Gao, W., Xia, S., He, S., Zhou, W. 2019. Development of an algal treatment system for selenium removal: Effects of environmental factors and post-treatment processing of Se-laden algae. *Journal of Hazardous Materials*, **365**, 546-554.

References

- Liu, F., Huang, J.C., Zhou, C., He, S., Zhou, W. 2016. Development of an algal treatment system for Se removal: Effects of light regimes, nutrients, sulfate and hypersalinity. *Chemosphere*, **164**, 372-378.
- Lönnerdal, B. 2002. Phytic acid–trace element (Zn, Cu, Mn) interactions. *International Journal of Food Science & Technology*, **37**(7), 749-758.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall, R.J. 1951. Protein measurement with the Folin phenol reagent. *The Journal of Biological Chemistry*, **193**(1), 265-75.
- Ma, X.-N., Chen, T.-P., Yang, B., Liu, J., Chen, F. 2016. Lipid Production from *Nannochloropsis*. *Marine Drugs*, **14**(4), 61.
- Mal, J., Nancharaiah, Y.V., van Hullebusch, E.D., Lens, P.N.L. 2016. Effect of heavy metal co-contaminants on selenite bioreduction by anaerobic granular sludge. *Bioresource Technology*, **206**, 1-8.
- Malagoli, M., Schiavon, M., dall'Acqua, S., Pilon-Smits, E.A. 2015. Effects of selenium biofortification on crop nutritional quality. *Frontiers in Plant Science*, **6**, 280.
- Malik, J.A., Goel, S., Kaur, N., Sharma, S., Singh, I., Nayyar, H. 2012. Selenium antagonises the toxic effects of arsenic on mungbean (*Phaseolus aureus* Roxb.) plants by restricting its uptake and enhancing the antioxidative and detoxification mechanisms. *Environmental and Experimental Botany*, **77**, 242-248.
- Mao, H., Wang, J., Wang, Z., Zan, Y., Lyons, G., Zou, C. 2014. Using agronomic biofortification to boost zinc, selenium, and iodine concentrations of food crops grown on the loess plateau in China. *Journal of Soil Science and Plant Nutrition*, **14**(2), 459-470.
- Maqbool, M.A., Beshir, A., Lübberstedt, T. 2019. Zinc biofortification of maize (*Zea mays* L.): Status and challenges. *Plant Breeding*, **138**(1), 1-28.
- Markou, G., Wang, L., Ye, J., Unc, A. 2018. Using agro-industrial wastes for the cultivation of microalgae and duckweeds: Contamination risks and biomass safety concerns. *Biotechnology Advance*, **36**(4), 1238-1254.
- Matassa, S., Verstraete, W., Pikaar, I., Boon, N. 2016. Autotrophic nitrogen assimilation and carbon capture for microbial protein production by a novel enrichment of hydrogen-oxidizing bacteria. *Water Research*, **101**, 137-146.

References

- Mechora, S., Stibilj, V., Germ, M. 2015. Response of duckweed to various concentrations of selenite. *Environmental Science and Pollution Research*, **22**(4), 2416-2422.
- Michalak, I., Chojnacka, K., Dmytryk, A., Wilk, R., Gramza, M., Roj, E. 2016. Evaluation of supercritical extracts of algae as biostimulants of plant growth in field trials. *Frontiers in Plant Science*, **7**, 1591.
- Michiels, J., Tagliabue, M.M., Akbarian, A., Owyn, A., De Smet, S. 2014. Oxidative status, meat quality and fatty acid profile of broiler chickens reared under free-range and severely feed-restricted conditions compared with conventional indoor rearing. *Avian Biology Research*, **7**(2), 74-82.
- Miranda, A.F., Biswas, B., Ramkumar, N., Singh, R., Kumar, J., James, A., Roddick, F., Lal, B., Subudhi, S., Bhaskar, T., Mouradov, A. 2016. Aquatic plant *Azolla* as the universal feedstock for biofuel production. *Biotechnology for Biofuels*, **9**(1), 221.
- Mishra, V. 2014. Biosorption of zinc ion: a deep comprehension. *Applied Water Science*, **4**(4), 311-332.
- Moheimani, N.R., Vadiveloo, A., Ayre, J.M., Pluske, J.R. 2018. Nutritional profile and in vitro digestibility of microalgae grown in anaerobically digested piggery effluent. *Algal Research*, **35**, 362-369.
- Mohsenpour, S.F., Hennige, S., Willoughby, N., Adeloye, A., Gutierrez, T. 2021. Integrating micro-algae into wastewater treatment: A review. *Science of the Total Environment*, **752**, 142168.
- Montalvo, D., Degryse, F., da Silva, R.C., Baird, R., McLaughlin, M.J. 2016. Agronomic effectiveness of Zinc sources as micronutrient Fertilizer. *Advance in Agronomy*, **139**, 215-267.
- Montville, T.J., Matthews, K.R. 2008. *Food Microbiology*. John Wiley & Sons, Limited.
- Moreno-Martin, G., Sanz-Landaluze, J., Leon-Gonzalez, M.E., Madrid, Y. 2020. Insights into the accumulation and transformation of Ch-SeNPs by *Raphanus sativus* and *Brassica juncea*: Effect on essential elements uptake. *Science of the Total Environment*, **725**, 138453.

References

- Morrison, S.J., Metzler, D.R., Dwyer, B.P. 2002. Removal of As, Mn, Mo, Se, U, V and Zn from groundwater by zero-valent iron in a passive treatment cell: reaction progress modeling. *Journal of Contaminant Hydrology*, **56**(1-2), 99-116.
- Mulbry, W., Westhead, E.K., Pizarro, C., Sikora, L. 2005. Recycling of manure nutrients: use of algal biomass from dairy manure treatment as a slow release fertilizer. *Bioresource Technology*, **96**(4), 451-8.
- Mushtaq, W., Mehdizade, M., Badruzzaman Siddiqui, M., Ozturk, M., Jabran, K., Altay, V. 2020. Phytotoxicity of above - ground weed residue against some crops and weeds. *Pakistan Journal of Botany*, **52**(3).
- N. Terry, A. M. Zayed, M. P. de Souza, a., Tarun, A.S. 2000. SELENIUM IN HIGHER PLANTS. *Annual Review of Plant Physiology and Plant Molecular Biology*, **51**(1), 401-432.
- Naramabuye, F.X., Haynes, R.J. 2006. Effect of organic amendment on soil pH and Al solubility and use of laboratory indices to predict their liming effect. *Soil Science*, **171**(10), 754-763.
- Ngigi, P.B. 2019. Assessment and optimization of dietary selenium intake in Kenya: exploration of biofortification as a solution to the hidden hunger. *PhD dissertation*, Faculty of Bioscience Engineering, Ghent University, Ghent, Belgium.
- Ngigi, P.B., Lachat, C., Masinde, P.W., Du Laing, G. 2019. Agronomic biofortification of maize and beans in Kenya through selenium fertilization. *Environmental Geochemistry and Health*, **41**(6), 2577-2591.
- Ohlbaum, M., Wadgaonkar, S.L., van Bruggen, J.J.A., Nancharaiah, Y.V., Lens, P.N.L. 2018. Phytoremediation of seleniferous soil leachate using the aquatic plants *Lemna minor* and *Egeria densa*. *Ecological Engineering*, **120**, 321-328.
- Opala, P.A., Okalebo, J.R., Othieno, C.O. 2012. Effects of organic and inorganic materials on soil acidity and phosphorus availability in a soil incubation study. *ISRN Agronomy*, **2012**, 597216.
- Orchard, T. 1977. Estimating the parameters of plant seedling emergence. *Seed Science and Technology*, **5**, 61-69.
- Otleş, S., Pire, R. 2001. Fatty acid composition of *Chlorella* and *Spirulina* microalgae species. *J AOAC Int*, **84**(6), 1708-14.

References

- Palmgren, M.G., Clemens, S., Williams, L.E., Kramer, U., Borg, S., Schjorring, J.K., Sanders, D. 2008. Zinc biofortification of cereals: problems and solutions. *Trends in Plant Science*, **13**(9), 464-73.
- Peng, Q., Wang, D., Wang, M., Zhou, F., Yang, W., Liu, Y., Liang, D. 2019. Prediction of selenium uptake by pak choi in several agricultural soils based on diffusive gradients in thin-films technique and single extraction. *Environmental Pollution*, 113414.
- Poblaciones, M.J., Rengel, Z. 2017. Combined foliar selenium and zinc biofortification in field pea (*Pisum sativum*): accumulation and bioavailability in raw and cooked grains. *Crop and Pasture Science*, **68**(3), 265.
- Prado, R.d.M., Cruz, F.J.R., Ferreira, R.L.d.C. 2017. Selenium biofortification and the problem of its safety. in: *Superfood and Functional Food - An Overview of Their Processing and Utilization*, (Ed.) V.Y. Waisundara, INTECH.
- Rachidi, F., Benhima, R., Sbabou, L., El Arroussi, H. 2020. Microalgae polysaccharides bio-stimulating effect on tomato plants: Growth and metabolic distribution. *Biotechnology Reports*, **25**, e00426.
- Rasouli, Z., Valverde-Pérez, B., D'Este, M., De Francisci, D., Angelidaki, I. 2018. Nutrient recovery from industrial wastewater as single cell protein by a co-culture of green microalgae and methanotrophs. *Biochemical Engineering Journal*, **134**, 129-135.
- Rayman, M.P. 2008. Food-chain selenium and human health: emphasis on intake. *British Journal of Nutrition*, **100**(2), 254-68.
- Rayman, M.P. 2000. The importance of selenium to human health. *The Lancet*, **356**(9225), 233-241.
- Rayman, M.P. 2012. Selenium and human health. *The Lancet*, **379**(9822), 1256-1268.
- Rayman, M.P. 2004. The use of high-selenium yeast to raise selenium status: how does it measure up? *British Journal of Nutrition*, **92**(4), 557-73.
- Reunova, Y.A., Aizdaicher, N.A., Khristoforova, N.K., Reunov, A.A. 2007. Effects of selenium on growth and ultrastructure of the marine unicellular alga *Dunaliella salina* (Chlorophyta). *Russian Journal of Marine Biology*, **33**(2), 125-132.
- Reyes, L.H., Encinar, J.R., Marchante-Gayon, J.M., Alonso, J.I., Sanz-Medel, A. 2006. Selenium bioaccessibility assessment in selenized yeast after "in vitro"

References

- gastrointestinal digestion using two-dimensional chromatography and mass spectrometry. *Journal of Chromatography A*, **1110**(1-2), 108-16.
- Roekens, E.J., Robberecht, H.J., Deelstra, H.A. 1986. Dietary selenium intake in Belgium for different population groups at risk for deficiency. *Diätetische Selen-Aufnahme in Belgien bei verschiedenen Bevölkerungsgruppen mit den Risiken eines Defizits*, **182**(1), 8-13.
- Rofkar, J.R., Dwyer, D.F., Bobak, D.M. 2014. Uptake and toxicity of arsenic, copper, and silicon in *Azolla Caroliniana* and *Lemna Minor*. *International Journal of Phytoremediation*, **16**(2), 155-166.
- Rolland, F., Moore, B., Sheen, J. 2002. Sugar sensing and signaling in plants. *Plant Cell*, **14 Suppl**(Suppl), S185-205.
- Ronga, D., Biazzi, E., Parati, K., Carminati, D., Carminati, E., Tava, A. 2019. Microalgal biostimulants and biofertilisers in crop productions. *Agronomy*, **9**(192), 1-22.
- Roohani, N., Hurrell, R., Kelishadi, R., Schulin, R. 2013. Zinc and its importance for human health: An integrative review. *Journal of research in medical sciences : the official journal of Isfahan University of Medical Sciences*, **18**(2), 144-157.
- Ros, G.H., van Rotterdam, A.M.D., Bussink, D.W., Bindraban, P.S. 2016. Selenium fertilization strategies for bio-fortification of food: an agro-ecosystem approach. *Plant and Soil*, **404**(1), 99-112.
- Ruz, M., Codoceo, J., Galgani, J., Muñoz, L., Gras, N., Muzzo, S., Leiva, L., Bosco, C. 1999. Single and multiple selenium-zinc-iodine deficiencies affect rat thyroid metabolism and ultrastructure. *The Journal of Nutrition*, **129**(1), 174-180.
- Salt, D.E., Prince, R.C., Pickering, I.J. 2002. Chemical speciation of accumulated metals in plants: evidence from X-ray absorption spectroscopy. *Microchemical Journal*, **71**(2-3), 255-259.
- Sánchez, E., SidaArreola, J., vilaQuezada, G., OjedaBarrios, D., SotoParra, J., PreciadoRangel, P. 2017. Can biofortification of zinc improve the antioxidant capacity and nutritional quality of beans? *Emirates Journal of Food and Agriculture*, **29**(3), 237.
- Sarathchandra, S., Watkinson, J. 1981. Oxidation of elemental selenium to selenite by *Bacillus megaterium*. *Science*, **211**(4482), 600-601.

References

- Sarwar, N., Akhtar, M., Kamran, M.A., Imran, M., Riaz, M.A., Kamran, K., Hussain, S. 2020. Selenium biofortification in food crops: Key mechanisms and future perspectives. *Journal of Food Composition and Analysis*, **93**, 103615.
- Sasmaz, M., Arslan Topal, E.I., Obek, E., Sasmaz, A. 2015. The potential of *Lemna gibba* L. and *Lemna minor* L. to remove Cu, Pb, Zn, and As in gallery water in a mining area in Keban, Turkey. *Journal of Environmental Management*, **163**, 246-53.
- Sazawal, S., Dhingra, U., Dhingra, P., Dutta, A., Deb, S., Kumar, J., Devi, P., Prakash, A. 2018. Efficacy of high zinc biofortified wheat in improvement of micronutrient status, and prevention of morbidity among preschool children and women - a double masked, randomized, controlled trial. *Nutr J*, **17**(1), 86.
- Schiavon, M., Ertani, A., Parrasia, S., Vecchia, F.D. 2017a. Selenium accumulation and metabolism in algae. *Aquatatic Toxicology*, **189**, 1-8.
- Schiavon, M., Lima, L.W., Jiang, Y., Hawkesford, M.J. 2017b. Effects of selenium on plant metabolism and implications for crops and consumers. in: *Selenium in plants: Molecular, Physiological, Ecological and Evolutionary Aspects*, (Eds.) E.A.H. Pilon-Smits, L.H.E. Winkel, Z.-Q. Lin, Springer International Publishing. Cham, pp. 257-275.
- Schiavon, M., Pilon-Smits, E.A., Citta, A., Folda, A., Rigobello, M.P., Dalla Vecchia, F. 2016. Comparative effects of selenate and selenite on selenium accumulation, morphophysiology, and glutathione synthesis in *Ulva australis*. *Environmental Science and Pollution Research*, **23**(15), 15023-32.
- Schrauzer, G.N. 2003. The nutritional significance, metabolism and toxicology of selenomethionine. in: *Advances in Food and Nutrition Research*, Vol. 47, Academic Press, pp. 73-112.
- Schrauzer, G.N. 2000. Selenomethionine: A review of its nutritional significance, metabolism and toxicity. *The Journal of Nutrition*, **130**(7), 1653-1656.
- Schumann, A.W., Sumner, M.E. 2004. Formulation of environmentally sound waste mixtures for land application. *Water, Air, and Soil Pollution*, **152**(1), 195-217.
- Sciences, N.A.o. 2000. Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids, (Ed.) N.A.o. Sciences, The National Academies Press. Washington, DC, pp. 528.

References

- Shaaban, M.M. 2001a. Green microalgae water extract as foliar feeding to wheat plants. *Pakistan Journal of Biological Sciences*, **4**(6), 628-632.
- Shaaban, M.M. 2001b. Nutritional status and growth of maize plants as affected by green microalgae as soil additives. *Journal of Biological Sciences*, **1**(6), 475-479.
- Sihlahla, M., Mouri, H., Nomngongo, P.N. 2019. Uptake of trace elements by vegetable plants grown on agricultural soils: Evaluation of trace metal accumulation and potential health risk. *Journal of African Earth Sciences*, **160**, 103635.
- Silambarasan, S., Logeswari, P., Sivaramakrishnan, R., Incharoensakdi, A., Cornejo, P., Kamaraj, B., Chi, N.T.L. 2021. Removal of nutrients from domestic wastewater by microalgae coupled to lipid augmentation for biodiesel production and influence of deoiled algal biomass as biofertilizer for *Solanum lycopersicum* cultivation. *Chemosphere*, **268**, 129323.
- Simmons, D.B., Wallschlager, D. 2011. Release of reduced inorganic selenium species into waters by the green fresh water algae *Chlorella vulgaris*. *Environmental Science & Technology*, **45**(6), 2165-71.
- Singh, H.P., Batish, D.R., Pandher, J.K., Kohli, R.K. 2003. Assessment of allelopathic properties of *Parthenium hysterophorus* residues. *Agriculture, Ecosystems & Environment*, **95**(2-3), 537-541.
- Singh, M., Singh, N. 1978. Selenium toxicity in plants and its detoxication by phosphorus. *Soil Science*, **126**(5), 255-262.
- Solórzano, L. 1969. Determination of ammonia in natural seawater by the phenol-hypochlorite method. *Limnology and Oceanography*, **14**(5), 799-801.
- Staicu, L.C., van Hullebusch, E.D., Lens, P.N.L. 2015a. Production, recovery and reuse of biogenic elemental selenium. *Environmental Chemistry Letters*, **13**(1), 89-96.
- Staicu, L.C., van Hullebusch, E.D., Oturan, M.A., Ackerson, C.J., Lens, P.N. 2015b. Removal of colloidal biogenic selenium from wastewater. *Chemosphere*, **125**, 130-138.
- Stavridou, E., Thorup-Kristensen, K., Young, S.D. 2011. Assessment of selenium mineralization and availability from catch crops. *Soil Use and Management*, no-no.

References

- Stone, R. 2009. Diseases. A medical mystery in middle China. *Science*, **324**(5933), 1378-81.
- Stonehouse, G.C., McCarron, B.J., Guignardi, Z.S., El Mehdawi, A.F., Lima, L.W., Fakra, S.C., Pilon-Smits, E.A.H. 2020. Selenium metabolism in Hemp (*Cannabis sativa* L.)-potential for phytoremediation and biofortification. *Environment Science & Technology*, **54**(7), 4221-4230.
- Subramanyam, K., Du Laing, G., Van Damme, E.J.M. 2019. Sodium selenate treatment using a combination of seed priming and foliar spray alleviates salinity stress in rice. *Frontiers in Plant Science*, **10**(116).
- Sun, X., Zhong, Y., Huang, Z., Yang, Y. 2014. Selenium accumulation in unicellular green alga *Chlorella vulgaris* and its effects on antioxidant enzymes and content of photosynthetic pigments. *PLoS One*, **9**(11), e112270.
- Supraja, K.V., Behera, B., Balasubramanian, P. 2020. Efficacy of microalgal extracts as biostimulants through seed treatment and foliar spray for tomato cultivation. *Industrial Crops and Products*, **151**, 112453.
- Supriatin, S., Weng, L., Comans, R.N.J. 2016. Selenium-rich dissolved organic matter determines selenium uptake in wheat grown on low-selenium arable land soils. *Plant and Soil*, **408**(1-2), 73-94.
- Tan, J.a., Zhu, W., Wang, W., Li, R., Hou, S., Wang, D., Yang, L. 2002. Selenium in soil and endemic diseases in China. *Science of The Total Environment*, **284**(1), 227-235.
- Tan, L.C., Nanchaiah, Y.V., van Hullebusch, E.D., Lens, P.N.L. 2016. Selenium: environmental significance, pollution, and biological treatment technologies. *Biotechnology Advance*, **34**(5), 886-907.
- Tan, X.-B., Zhao, X.-C., Yang, L.-B. 2019. Strategies for enhanced biomass and lipid production by *Chlorella pyrenoidosa* culture in starch processing wastewater. *Journal of Cleaner Production*, **236**, 117671.
- Terry, N., Zayed, A.M., de Souza, M.P., Tarun, A.S. 2000. Selenium in higher plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, **51**(1), 401-432.
- Thavarajah, D., Ruszkowski, J., Vandenberg, A. 2008. High potential for selenium biofortification of lentils (*Lens culinaris* L.). *Journal of Agricultural and Food Chemistry*, **56**(22), 10747-10753.

References

- Thomson, C.D. 2004. Selenium and iodine intakes and status in New Zealand and Australia. *British Journal of Nutrition*, **91**(5), 661-672.
- Tsuji, N., Hirayanagi, N., Okada, M., Miyasaka, H., Hirata, K., Zenk, M.H., Miyamoto, K. 2002. Enhancement of tolerance to heavy metals and oxidative stress in *Dunaliella tertiolecta* by Zn-induced phytochelatin synthesis. *Biochemical and Biophysical Research Communications*, **293**(1), 653-659.
- Umysova, D., Vitova, M., Douskova, I., Bisova, K., Hlavova, M., Cizkova, M., Machat, J., Doucha, J., Zachleder, V. 2009. Bioaccumulation and toxicity of selenium compounds in the green alga *Scenedesmus quadricauda*. *BMC Plant Biology*, **9**, 58.
- USEAP. 2016. Aquatic life ambient water quality criterion for selenium - Freshwater 2016. United States Environmental Protection Agency (EPA-822-R-16-006).
- Van Hoewyk, D. 2013. A tale of two toxicities: malformed selenoproteins and oxidative stress both contribute to selenium stress in plants. *Annals of Botany*, **112**(6), 965-72.
- Van Ranst, E., Verloo, M., Demeyer, A. 1999. *Manual for the soil chemistry and fertility laboratory: analytical methods for soils and plants equipment and management of consumables*. Ghent : UGent. Faculty of agricultural and applied biological sciences. Department of applied analytical and physical chemistry.
- van Zelst, M., Hesta, M., Gray, K., Staunton, R., Du Laing, G., Janssens, G.P.J. 2016. Biomarkers of selenium status in dogs. *Bmc Veterinary Research*, **12**.
- Versini, A., Di Tullo, P., Aubry, E., Bueno, M., Thiry, Y., Pannier, F., Castrec-Rouelle, M. 2016. Influence of Se concentrations and species in hydroponic cultures on Se uptake, translocation and assimilation in non-accumulator ryegrass. *Plant Physiology and Biochemistry*, **108**, 372-380.
- Vitova, M., Bisova, K., Hlavova, M., Zachleder, V., Rucki, M., Cizkova, M. 2011. Glutathione peroxidase activity in the selenium-treated alga *Scenedesmus quadricauda*. *Aquatic Toxicology*, **102**(1-2), 87-94.
- Vriens, B., Behra, R., Voegelin, A., Zupanic, A., Winkel, L.H. 2016. Selenium Uptake and Methylation by the Microalga *Chlamydomonas reinhardtii*. *Environmental Science and Technology*, **50**(2), 711-20.

References

- Vu, D.L., Saurav, K., Mylenko, M., Ranglova, K., Kuta, J., Ewe, D., Masojidek, J., Hrouzek, P. 2019. In vitro bioaccessibility of selenoamino acids from selenium (Se)-enriched *Chlorella vulgaris* biomass in comparison to selenized yeast; a Se-enriched food supplement; and Se-rich foods. *Food Chemistry*, **279**, 12-19.
- Wake, H. 2005. Oil refineries: a review of their ecological impacts on the aquatic environment. *Estuarine, Coastal and Shelf Science*, **62**(1-2), 131-140.
- Wallenberg, M., Olm, E., Hebert, C., Bjornstedt, M., Fernandes, A.P. 2010. Selenium compounds are substrates for glutaredoxins: a novel pathway for selenium metabolism and a potential mechanism for selenium-mediated cytotoxicity. *Biochemical Journal*, **429**, 85-93.
- Wan, Y.N., Yu, Y., Wang, Q., Qiao, Y.H., Li, H.F. 2016. Cadmium uptake dynamics and translocation in rice seedling: Influence of different forms of selenium. *Ecotoxicology and Environmental Safety*, **133**, 127-134.
- Wang, D., Dinh, Q.T., Anh Thu, T.T., Zhou, F., Yang, W., Wang, M., Song, W., Liang, D. 2018. Effect of selenium-enriched organic material amendment on selenium fraction transformation and bioavailability in soil. *Chemosphere*, **199**, 417-426.
- Wang, M., Ali, F., Wang, M., Dinh, Q.T., Zhou, F., Bañuelos, G.S., Liang, D. 2020. Understanding boosting selenium accumulation in Wheat (*Triticum aestivum* L.) following foliar selenium application at different stages, forms, and doses. *Environmental Science and Pollution Research*, **27**(1), 717-728.
- Wang, M.K., Peng, Q., Zhou, F., Yang, W.X., Dinh, Q.T., Liang, D.L. 2019. Uptake kinetics and interaction of selenium species in tomato (*Solanum lycopersicum* L.) seedlings. *Environmental Science and Pollution Research*, **26**(10), 9730-9738.
- Wang, Q., Yu, Y., Li, J., Wan, Y., Huang, Q., Guo, Y., Li, H. 2017. Effects of different forms of selenium fertilizers on Se accumulation, distribution, and residual effect in winter wheat-summer maize rotation system. *Journal of Agricultural and Food Chemistry*, **65**(6), 1116-1123.
- Wang, Z., Gao, Y. 2001. Biogeochemical cycling of selenium in Chinese environments. *Applied Geochemistry*, **16**(11), 1345-1351.

References

- Weng, L., Vega, F.A., Supriatin, S., Bussink, W., Riemsdijk, W.H.V. 2011. Speciation of Se and DOC in soil solution and their relation to Se bioavailability. *Environmental Science & Technology*, **45**(1), 262-267.
- White, P.J. 2016. Selenium accumulation by plants. *Annals of Botany*, **117**(2), 217-235.
- White, P.J., Broadley, M.R. 2009. Biofortification of crops with seven mineral elements often lacking in human diets – iron, zinc, copper, calcium, magnesium, selenium and iodine. *New Phytologist*, **182**(1), 49-84.
- White, P.J., Broadley, M.R. 2005. Biofortifying crops with essential mineral elements. *Trends in Plant Science*, **10**(12), 586-593.
- Winkel, L.H., Johnson, C.A., Lenz, M., Grundl, T., Leupin, O.X., Amini, M., Charlet, L. 2012. Environmental selenium research: from microscopic processes to global understanding. *Environmental Science & Technology*, **46**(2), 571-9.
- Winkel, L.H., Vriens, B., Jones, G.D., Schneider, L.S., Pilon-Smits, E., Banuelos, G.S. 2015. Selenium cycling across soil-plant-atmosphere interfaces: a critical review. *Nutrients*, **7**(6), 4199-239.
- Witt, C., Cassman, K.G., Olk, D.C., Biker, U., Liboon, S.P., Samson, M.I., Ottow, J.C.G. 2000. Crop rotation and residue management effects on carbon sequestration, nitrogen cycling and productivity of irrigated rice systems. *Plant and Soil*, **225**(1), 263-278.
- Wolf, E., Kollonitsch, V., Kline, C.H. 1963. Selenium supplementation, survey of selenium treatment in livestock production. *Journal of Agricultural and Food Chemistry*, **11**(4), 355-360.
- Wu, Z., Bañuelos, G.S., Lin, Z.-Q., Liu, Y., Yuan, L., Yin, X., Li, M. 2015. Biofortification and phytoremediation of selenium in China. *Frontiers in Plant Science*, **6**(136).
- Yadav, S. 2010. Heavy metals toxicity in plants: An overview on the role of glutathione and phytochelatins in heavy metal stress tolerance of plants. *South African Journal of Botany*, **76**, 167-179.
- Yawata, A., Oishi, Y., Anan, Y., Ogra, Y. 2010. Comparison of selenium metabolism in three *Brassicaceae* plants. *Journal of Health Science*, **56**(6), 699-704.
- Zahedi, S.M., Hosseini, M.S., Daneshvar Hakimi Meybodi, N., Teixeira da Silva, J.A. 2019. Foliar application of selenium and nano-selenium affects pomegranate

References

- (*Punica granatum* cv. *Malase Saveh*) fruit yield and quality. *South African Journal of Botany*, **124**, 350-358.
- Zaman, Q.u., Aslam, Z., Yaseen, M., Ihsan, M.Z., Khaliq, A., Fahad, S., Bashir, S., Ramzani, P.M.A., Naeem, M. 2018. Zinc biofortification in rice: leveraging agriculture to moderate hidden hunger in developing countries. *Archives of Agronomy and Soil Science*, **64**(2), 147-161.
- Zhan, X., Wang, M., Zhao, R., Li, W., Xu, Z. 2007. Effects of different selenium source on selenium distribution, loin quality and antioxidant status in finishing pigs. *Animal Feed Science and Technology*, **132**(3), 202-211.
- Zhang, H., Zhao, Z., Zhang, X., Zhang, W., Huang, L., Zhang, Z., Yuan, L., Liu, X. 2019. Effects of foliar application of selenate and selenite at different growth stages on selenium accumulation and speciation in potato (*Solanum tuberosum* L.). *Food Chemistry*, **286**, 550-556.
- Zhang, Y.L., Pan, G.X., Chen, J., Hu, Q.H. 2003. Uptake and transport of selenite and selenate by soybean seedlings of two genotypes. *Plant and Soil*, **253**(2), 437-443.
- Zhao, Y., Fang, Y., Jin, Y., Huang, J., Bao, S., Fu, T., He, Z., Wang, F., Zhao, H. 2014. Potential of duckweed in the conversion of wastewater nutrients to valuable biomass: a pilot-scale comparison with water hyacinth. *Bioresource Technology*, **163**, 82-91.
- Zhong, Y., Cheng, J. 2016. Effects of selenium on biological and physiological properties of the duckweed *Landoltia punctata*. *Plant Biology*, **18**(5), 797-804.
- Zhong, Y., Cheng, J.J. 2017. Effects of selenite on unicellular green microalga *Chlorella pyrenoidosa*: Bioaccumulation of selenium, enhancement of photosynthetic pigments, and amino acid production. *Journal of Agricultural and Food Chemistry*, **65**(50), 10875-10883.
- Zhong, Y., Li, Y., Cheng, J.J. 2016. Effects of selenite on chlorophyll fluorescence, starch content and fatty acid in the duckweed *Landoltia punctata*. *Journal of Plant Research*, **129**(5), 997-1004.
- Zhou, F., Peng, Q., Wang, M., Liu, N., Dinh, Q.T., Zhai, H., Xue, M., Liang, D. 2021. Influence of processing methods and exogenous selenium species on the

References

- content and in vitro bioaccessibility of selenium in *Pleurotus eryngii*. *Food Chemistry*, **338**, 127661.
- Zhu, L.X., Wang, P., Zhang, W.J., Hui, F., Chen, X.X. 2017. Effects of selenium application on nutrient uptake and nutritional quality of *Codonopsis lanceolata*. *Scientia Horticulturae*, **225**, 574-580.

Acknowledgement

Acknowledgement

The thought of writing this acknowledgment seemed something very far and even impossible to imagine at beginning of my PhD - four years ago. I still cannot believe that I am gonna close the four years PhD chapter soon. Looking back on the four years journey, it was full of challenges, detours, valleys, and even mistakes. But finally, I almost reach the top of PhD mountain today. I appreciate many people for helping and supporting me to achieve the doctoral degree.

First of all, I would like to express appreciation to my supervisors Prof. Gijs Du Laing, Prof. Piet Lens and Prof. Ivet Ferrer for their guidance during these years. Prof. Gijs Du Laing, thank you for giving me the opportunity to carry out this doctoral research and teach me critical thinking during the research journey. I still remember the first skype interview. You clearly introduced your research/Ecochem lab/current PhD students to me, and even checked some practical details for me as an international student, such as the official application procedure and the China platform in Gent University. Thank you a lot for introducing me into SuPER-W, such a nice project, in which I had chances to meet many excellent early-stage researchers and receive a lot of professional training for improving my technical and complementary skills. Prof. Piet Lens, I am grateful that you accommodated me in IHE, Delft for 6 months as start-up for my PhD lab/research life. Thank you for patiently correcting my manuscripts over and over again and tracking my progress regularly. Thank you for teaching me that a PhD student should not only work in the lab. Prof. Ivet, it was super nice to have 6 months' experience in the GEMMA group, Barcelona with you. Because of your countless help and support, I could successfully complete the secondment. Thanks for all your understanding and consideration for my PhD during these years.

I would also like to thank all the members of Examination Committee, Prof. Stefaan De Smet, Prof. Diederik Rousseau, Prof. Dongli Liang, Prof. Gary Bañuelos and Dr. Rubén Díez-Montero for detailed and insightful comments, which helped me a lot in improving this thesis.

It would not have been possible to perform and finalize the lab work without the precious help of Joachim Neri, Roseline Blanckaert, Melgü Kizilmese, Katty Sabo and

Acknowledgement

Wim Goeteyn from the Ecochem lab of Ghent University, and all technicians from IHE and GEMMA group of Universitat Politècnica de Catalunya. My sincere thanks go also to the previous master thesis students, Ana María Murillo Abril and Maria Josep for the contribution to our work.

I would like to express my gratitude to all my colleagues and friends from Ecochem for their sharing, encouragement, and help throughout this PhD journey. They are with many, as the group was growing fast over the past few years. I would also like to thank the colleagues and friends in IHE and GEMMA group, who I ever met.

Last but not least, I would like to express my warmest thanks to my family. 感谢父母一如既往且坚定的支持和鼓励，因为有你们的支持我才能顺利的走到今天，才能在异国他乡安心的完成博士论文。同时十分感谢长兄及长嫂多年替我默默陪伴在父母身旁。 Shulin, you certainly deserve my highest appreciation for this achievement.

Jun Li

Ghent, May 2021.

Curriculum Vitae

Curriculum Vitae

Personal Information

Name: Jun Li

Nationality: China

E-mail: jun.li.nwsuaf@gmail.com

Corresponding address: Coupure Links 653, 9000 Gent, Belgium

Education

UGent University	PhD student	2016.10 - Now
• Faculty of Bioscience Engineering		
Northwest A&F University	Master of Engineering	2013.09-2016.06
• Faculty of Resource and Environment		
Northwest A&F University	Bachelor of Science	2009.09-2013.06
• Faculty of Resource and Environment		

Experience

Avecom Co., Belgium	2020.01-2020.03
• Production of Se-enriched single cell protein	
Universitat Politècnica de Catalunya, Barcelona	2019.01-2019.08
• Se removal from wastewater and production of Se-enriched microalgae	
UNESCO-IHE, The Netherlands	2018.02-2018.10
• Se/Zn removal from wastewater by anaerobic digestion and phytoremediation	
Gent University, Belgium	2016.10-2020.10
• Valorization of Se/Zn-enriched biomaterials as feed supplements and micronutrient fertilizers	

Awards

Curriculum Vitae

- | | |
|----------------------------------|-----------|
| • BOF-CSC Scholarship | 2017-2019 |
| • Outstanding Master Thesis | 2016 |
| • Graduate National Scholarship | 2015 |
| • Excellent Postgraduate Student | 2015 |

Publications

Li, J., Ferrer, I., Lens, P.N.L., Du Laing, G., 2021 Evaluation of selenium-enriched microalgae produced on domestic wastewater as biostimulant and biofertilizer for growth of selenium-enriched crops. *Journal of Applied Phycology* (In press).

Li, J., Du Laing, G., Ferrer, I., Lens, P.N.L. 2021. Selenium biofortification for human and animal nutrition, In: Lens, P.N.L., Pakshirajan, K., (Eds.) *Environmental Technologies to Treat Selenium Pollution - Principles and Engineering*. IWA Publishing (In press).

Li, J., Otero-Gonzalez, L., Michiels, J., Lens, P.N.L., Du Laing, G., Ferrer, I., 2021. Production of Se-enriched microalgae as potential feed supplement in high rate algae ponds treating domestic wastewater. *Bioresource Technology*. 333, 125239.

Li, J., Otero-Gonzalez, L., Parao, A., Tack, P., Folens, K., Ferrer, I., Lens, P.N.L., Du Laing, G. 2021. Valorization of Se-enriched sludge and duckweed generated from wastewater as micronutrient biofertilizers. *Chemosphere*, 281, 130767.

Li, J., Loi, G., Otero-Gonzalez, L., Du Laing, G., Ferrer, I., Lens, P.N.L. 2020. Selenate and selenite uptake, accumulation and toxicity in *Lemna minuta*. *Water Science and Technology*, 81(9), 1852-1862.

Li, J., Lens, P.N.L., Otero-Gonzalez, L., Du Laing, G. 2020. Production of selenium- and zinc-enriched *Lemna* and *Azolla* as potential micronutrient-enriched bioproducts. *Water Research*, 172, 115522.

Li, J., Peng, Q., Liang, D., Liang, S., Chen, J., Sun, H., Li, S., Lei, P. 2016. Effects of aging on the fraction distribution and bioavailability of selenium in three different soils. *Chemosphere*, 144, 2351-2359.

Li, J., Liang, D., Qin, S., Feng, P., Wu, X. 2015. Effects of selenite and selenate application on growth and shoot selenium accumulation of pak choi (*Brassica chinensis* L.) during

Curriculum Vitae

successive planting conditions. *Environmental Science Pollution Research*, 22(14), 11076-11086.

Li, J., Qin, S., Feng, P., Man, N., Liang, D., 2015. The fraction of selenium in cumulated irrigated soil in Gansu Province, China: effect of aging. *Global Advances in Selenium Research from Theory to Application*, 31-32.

Li, J., Otero-Gonzalez, L., Du Laing, G., Lens, P.N.L., Ferrer, I. 2019. Exploration of Se-rich bioproducts generated from (waste)water as fertilizers. In book: *Selenium Research for Environment and Human Health: Perspectives, Technologies and Advancements*. DOI: 10.1201/9780429423482-49.

Conferences and Workshops

Conferences

The 11th International Symposium on Selenium in Biology and Medicine and The 5th International conference on Selenium in the Environment and Human Health. August 2017

- Stockholm, Sweden (Poster presentation)

The 15th International Conference on the Biogeochemistry of Trace element. May 2019

- Nanjing, China (Oral presentation)

3rd IWA Resource Recovery Conference 2019. September 2019

- Venice, Italy (Poster presentation)

The 6th International Conference on Selenium in the Environment and Human Health. November 2019

- Xi'an, China (Oral presentation, award one of the best presentations)

Workshops

Algal biotechnology techniques and opportunities for the sustainable bioeconomy

- Stuttgart, Germany

Workshop on business case development in the field of resource recovery, energy and product from wastewater.

- Prague, Czech Republic

Workshop on sustainability, societal aspects, policy and stakeholder involvement

- Delft, The Netherlands

Curriculum Vitae

Workshop on modelling tools

- Barcelona, Spain

Workshop on creative problem-solving in resource recovery technology

- Aachen, Germany
-