

Nancy Vegas ORCID iD: 0000-0002-8293-2383

Brian Hon-Yin Chung ORCID iD: 0000-0002-7044-5916

florence petit ORCID iD: 0000-0002-1368-1023

Further delineation of Auriculocondylar syndrome based on 14 novel cases and reassessment of 25 published cases

Nancy Vegas^{1*}, Zeynep Demir^{1,2*}, Christopher T. Gordon¹, Sylvain Breton³, Vanessa Romanelli Tavares⁴, Hugo Moisset¹, Roseli Zechi-Ceide⁵, Nancy M. Kokitsu-Nakata⁵, Yasuhiro Kido⁶, Sandrine Marlin^{1,7}, Souad Gherbi Halem⁷, Ilse Meerschaut⁸, Bert Callewaert⁸, Brian Chung⁹, Nicole Revencu¹⁰, Daphné Lehalle^{11,12}, Florence Petit¹³, Evan J. Propst¹⁴, Blake C. Papsin¹⁴, John H. Phillips¹⁴, Linda Jakobsen¹⁵, Pauline Le Tanno¹⁶, Julien Thévenon¹⁶, Julie McGaughran¹⁷, Erica H. Gerkes¹⁸, Chiara Leoni¹⁹, Peter Kroisel²⁰, Tiong Yang Tan²¹, Alex Henderson²², Paulien Terhal²³, Lina Basel-Salmon^{24, 25, 26}, Adila Alkindy²⁷, Susan M. White²¹, Maria Rita Passos Bueno⁴, Véronique Pingault^{1,28}, Loïc De Pontual^{1,29}, Jeanne Amiel^{1,28}

¹Laboratory of Embryology and Genetics of Malformations, Institut National de la Santé et de la Recherche Médicale (INSERM) UMR 1163, Université de Paris, Institut *Imagine*, Paris, France.

²Unité d'hépatologie pédiatrie et transplantation, Hôpital Necker-Enfants Malades, AP-HP, Paris, France.

³Service d'imagerie pédiatrie, Hôpital Necker-Enfants Malades, AP-HP, Paris, France

⁴Centro de Pesquisas do Genoma Humano e Celulas Tronco, Departamento de Genetica e Biología Evolutiva, Instituto de Biociencias, Universidade de Sao Paulo, Sao Paulo, Brazil

⁵Department of Clinical Genetics, Hospital for Rehabilitation of Craniofacial Anomalies, University of Sao Paulo, Bauru, Brazil

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/humu.24349.

This article is protected by copyright. All rights reserved.

⁶Department of Pediatrics, Dokkyo Medical University Koshigaya Hospital, Saitama, Japan

⁷Reference center for genetic hearing loss, Fédération de Génétique et de Médecine Génomique, Hôpital Necker, APHP.CUP, Paris, France.

⁸Center for Medical Genetics, Ghent University Hospital, and Department of Biomolecular Medicine, Ghent University, Belgium.

⁹Department of Paediatrics and Adolescent Medicine, The University of Hong Kong, Hong Kong

¹⁰Center for Human Genetics, Cliniques universitaires Saint Luc, Université catholique de Louvain, Brussels, Belgium.

¹¹Centre de génétique- centre de référence des maladies rares, anomalies du développement et syndrome malformatifs, Centre Hospitalo-Universitaire de Dijon, Bourgogne, France.

¹²UF de Génétique Médicale, Département de Génétique, Groupe Hospitalier Pitié-Salpêtrière, APHP Sorbonne Université, Paris, France.

¹³ CHU Lille, clinique de Génétique Guy Fontaine, F-59000 Lille, France

¹⁴Department of Otolaryngology-Head and Neck Surgery, The Hospital for Sick Children, University of Toronto, Canada.

¹⁵Department of Plastic Surgery, Copenhagen University Hospital, Herlev, Denmark

¹⁶Service de Génétique et Université Grenoble-Alpes, Grenoble, France

¹⁷Genetic Health Queensland, Royal Brisbane and Women's Hospital, Herston and the University of Queensland, St Lucia, Brisbane, Australia

¹⁸Department of Genetics, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

¹⁹Center for Rare Diseases and Birth Defects, Department of Woman and Child Health and Public Health, Fondazione Policlinico A. Gemelli, IRCCS, Italy

²⁰Institute of Human Genetics, Medical University of Graz, Graz, Austria

²¹Victorian Clinical Genetics Services, Murdoch Children's Research Institute, Royal Children's Hospital, and Department of Paediatrics, University of Melbourne, Melbourne, Victoria, Australia

²²Northern Genetics Service, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle Upon Tyne, UK

²³Department of Medical Genetics, University Medical Centre Utrecht, Utrecht, The Netherlands

²⁴Pediatric Genetics, Schneider Children's Medical Center of Israel and Raphael Recanati Genetics Institute, Rabin Medical Center, Beilinson Campus, Petah Tikva,

Israel

²⁵Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

²⁶Felsenstein Medical Research Center, Rabin Medical Center, Petah Tikva, Israel

²⁷Department of Genetics, Sultan Qaboos University Hospital, Muscat, Oman

²⁸Fédération de Génétique et de Médecine Génomique, Hôpital Necker, APHP.CUP, Paris, France.

²⁹Service de pédiatrie, Hôpital Jean Verdier, Bondy, France

* Equal first author

Corresponding authors:

Jeanne Amiel, Chris Gordon

INSERM U1163

Institut *Imagine*,

24 bd du Montparnasse

75015, Paris, France

Phone: (33) 1 44 49 56 48

Fax: (33) 1 44 49 51 50

jeanne.amiel@inserm.fr, chris.gordon@inserm.fr

Grant numbers:

USPC PRES (project SPC/JFG/2013-031),

ANR (project CranioRespiro),

MSD-Avenir (project Devo-Decode),

AXA (project Tête et Cœur)

Abstract

Auriculocondylar syndrome (ACS) is a rare craniofacial disorder characterized by mandibular hypoplasia and an auricular defect at the junction between the lobe and helix, known as a "Question Mark Ear" (QME). Several additional features,

originating from the first and second branchial arches and other tissues, have also been reported. ACS is genetically heterogeneous with autosomal dominant and recessive modes of inheritance. The mutations identified to date are presumed to dysregulate the endothelin 1 signalling pathway. Here we describe 14 novel cases and reassess 25 published cases of ACS through a questionnaire for systematic data collection. All patients harbour mutation(s) in *PLCB4*, *GNAI3* or *EDN1*. This series of patients contributes to the characterization of additional features occasionally associated with ACS such as respiratory, costal, neurodevelopmental and genital anomalies, and provides management and monitoring recommendations.

Key words: Craniofacial anomalies - Auriculocondylar syndrome - Question mark ear - EDN1 - PLCB4 - GNAI3

Introduction

Auriculocondylar syndrome (ACS; MIM#s 602483, 614669, 615706) is a rare craniofacial disorder affecting the first and second branchial arches. It is mainly characterized by variable micrognathia, hypoplasia of the condyle of the mandible and an auricular defect at the lobe-helix junction, known as a "Question Mark Ear" (QME) (Cosman, 1984; Cosman, Bellin, & Crikelair, 1970). Intra and inter familial phenotypic variation in ACS has previously been reported (Gerkes, van Ravenswaaij, & van Essen, 2008; Gordon, Petit, et al., 2013; Gordon, Vuillot, et al., 2013; Kokitsu-Nakata et al., 2012; Masotti et al., 2008; Rieder et al., 2012). The mildest end of the spectrum is an isolated QME (IQME; MIM# 612798). Several studies have described other frequently associated features including microstomia, full cheeks, facial asymmetry, cleft palate, hearing loss and lingual anomalies (Gerkes et al., 2008; Gordon, Petit, et al., 2013; Gordon, Vuillot, et al., 2013; Kokitsu-Nakata et al., 2012;

Masotti et al., 2008; Rieder et al., 2012). Extra-craniofacial manifestations include central sleep apnea, gastro-intestinal disorders and genital anomalies (Gordon, Vuillot, et al., 2013; Kido et al., 2013; Leoni et al., 2016).

ACS is genetically heterogeneous with both autosomal dominant and recessive modes of inheritance. Mutations in the Phospholipase C beta 4 gene (*PLCB4*; MIM# 600810) located on 20p12.3-p12.2 and in the Guanine Nucleotide Binding protein alpha inhibiting activity polypeptide 3 gene (*GNAI3*; MIM# 139370) located on 1p13.3 were identified by whole exome sequencing and confirmatory studies in affected individuals (Gordon, Vuillot, et al., 2013; Rieder et al., 2012). The Endothelin 1 gene (*EDN1*; MIM# 131240) was subsequently identified as a third locus at 6p24.1 in ACS and IQME patients (Gordon, Petit, et al., 2013). Studies in animal models have highlighted the key role of the EDN1-Endothelin Receptor type A (*EDNRA*; MIM# 131243) signalling pathway in the development of the jaw (Clouthier et al., 2013; Walker, Miller, Swartz, Eberhart, & Kimmel, 2007). Mice with targeted deletion of *Edn1* or *Ednra* present with severe defects in jaw development and a homeotic transformation of the lower jaw into an upper jaw (Clouthier, Garcia, & Schilling, 2010; Clouthier et al., 2013; Ozeki, Kurihara, Tonami, Watatani, & Kurihara, 2004; Ruest, Xiang, Lim, Levi, & Clouthier, 2004). Mutations in *GNAI3* and *PLCB4* are thought to interfere with cytoplasmic signalling in the EDN1-EDNRA pathway (Marivin et al., 2016). This is supported by the observation of genetic interaction between *edn1* and *plcb3* in zebrafish during craniofacial development (Walker et al., 2007) and by the fusion and hypoplasia of branchial arch cartilage elements observed in zebrafish with *plcb3* missense mutations (Walker et al., 2007).

We herein describe 14 novel patients with ACS/IQME from 10 families harbouring *PLCB4*, *GNAI3* or *EDNI* mutations. We analysed clinical features of these and previously reported patients in an attempt to establish correlations with the identified disease-causing mutations and to propose recommendations aiming at improving the management and monitoring of patients with ACS.

Patients and methods

PLCB4, *GNAI3* or *EDNI* Sanger sequencing was performed in most ACS or QME patients, according to standard procedures (Romanelli Tavares et al., 2017). Patients 8 and 9 were submitted to next-generation sequencing of a panel of neurocristopathy-related genes, including the three ACS genes (Gordon et al., 2018). Whole exome sequencing was performed for patients 23, 29 and 39. For patient 23 the exome protocol has been published (Thevenon et al., 2016). For patient 29, exome enrichment (Agilent SureSelectXT Human All Exon 50Mb), exome sequencing (Illumina HiSeq), read alignment (BWA) and variant calling (GATK) were performed at BGI-Europe (Denmark), and variant annotation and prioritizing were performed by the Department of Human Genetics, Radboud University Medical Center. For patient 39, exome enrichment (Agilent Sureselect QXT CREv1 kit), exome sequencing (Illumina instrument), read alignment and variant calling were performed by the Clinical genomics laboratory, Victorian Clinical Genetics Services, Melbourne, Australia. Molecular data are summarised in Table 1.

Clinical data of patients with ACS were collected through a standardized questionnaire sent to each referring physician requesting personal and familial medical history of affected individuals, clinical and radiological features, treatments and outcomes. Table 2 indicates the frequency of the major clinical features while

Supp. Table S1 provides detailed phenotypic information for each patient. Information about some clinical features (full cheeks, microstomia, post auricular tags) was obtained from photographs. Available radiological exams were reviewed by a paediatric radiologist (Dr Breton) with expertise in craniofacial imaging at the Necker-Enfants Malades Hospital.

The 14 previously unreported patients are described as follows.

Patient 8 from Family 7, A male was born at term from healthy non-consanguineous parents after an uneventful pregnancy with birth parameters in the normal range. At birth, bilateral QME with hypoplastic antihelix and prominent antitragus, micrognathia, and a hypomobile tongue were noted. Swallowing difficulties and poor suck required nasogastric tube feeding for the first 5 months of life. He snored during sleep but polysomnographic recording was not performed. Speech was delayed with dysarthria, due in part to hearing loss and hypomobile tongue. Growth parameters were normal at last examination at 7 years of age. Cardiovascular examination and skeletal x-rays were normal. CT-scan demonstrated mandibular hypoplasia, dysplasia of the condyles and temporomandibular joints (TMJs) and normal inner ears. A right mandibular osteotomy and mandibular distraction were performed at 2.5 years of age and a second mandibular distraction 33 months later. We identified a *de novo* c.1861C>T; p.(Arg621Cys) mutation in *PLCB4* (RefSeq NM_000933.4) in this patient.

Family 8 includes the proband, a male (**patient 9**), his mother (**patient 10**) and maternal grandmother (**patient 11**). ACS was diagnosed in the proband at 3 years of age based on the association of micro-retrognathia, microstomia and unilateral QME. He was term-born from non-consanguineous parents after an uneventful pregnancy

with birth parameters in the normal range. He had neither feeding nor breathing difficulties but presented with neonatal hypotonia followed by delayed motor and speech milestones. Craniofacial CT-scan and polysomnographic recordings were not performed. Hearing was normal and he had no surgical interventions.

His mother was diagnosed with bilateral QMEs. Since craniofacial CT-scan showed dysplasia of the condyles, she was classified as ACS. She presented with ptosis, hypermetropia and learning difficulties. Clinical examination of the maternal grandmother was normal. The proband, his mother and maternal grandmother were heterozygous for the *PLCB4* mutation c.1862G>A; p.(Arg621His).

Family 10 consists of an affected female (**patient 14**) and her mother (**patient 15**). The diagnosis of ACS was made at 5 years of age in patient 14. She was born full-term after an uneventful pregnancy with growth parameters in the low-normal range. She presented with bilateral QMEs, a bifid uvula and protrusion of the tongue. She had no feeding or breathing difficulties. Growth and motor development were normal. Later, orofacial dyspraxia and snoring were noted but polysomnography was not performed. Craniofacial CT-scan was not available.

Her mother presented with a mild unilateral QME, diabetes mellitus, a single kidney and a bifid uterus. Both harboured a heterozygous c.1705A>T; p.(Ile569Phe) mutation in *PLCB4*.

Family 11 consists of a father, **patient 16**, and his son, **patient 17**. The father was diagnosed with ACS in his fifties based on the association of microretrognathia and bilateral QME (Figure 1: a-c). He snored and experienced obstructive sleep apnea. He has had bilateral mandibular osteotomies and auricular reconstruction. Cardiovascular,

ophthalmological and hearing examinations have been normal. Speech was delayed without hearing loss in childhood and intelligence is within the normal range. A craniofacial CT-scan showed a hypoplastic mandible and hypogenesis or agenesis of the TMJs and condyles.

His son, **patient 17**, presented with bilateral IQMEs (Figure 1: d, e, f) and scoliosis. Speech had been delayed without hearing loss and intelligence was within the normal range. Craniofacial CT scan was not performed. The father and son have no facial hair in a proximal region along the horizontal branch of the mandible reflecting the putative mandibular to maxillary homeotic transformation underlying ACS (Figure 1: a-f) (Gordon et al., 2014). Both were heterozygous for a *PLCB4* c.1015G>A; p.(Gly339Arg) mutation.

Patient 18, a female, was term-born after an uneventful pregnancy from healthy consanguineous parents originating from North Africa. She presented with bilateral QME, severe microretrognathia and microstomia. Dental anomalies including agenesis and misalignment were subsequently observed. Cardiovascular and hearing tests have been normal. She snores and sleeps in a sitting position. A sleep study performed before distraction at 7 years of age demonstrated both obstructive and central apnea (apnea-hypopnea index (AHI) of 86 events per hour) with hypercapnia. Unfortunately, polysomnography was not repeated after mandibular distraction. Growth and psychomotor development were normal. CT-scan confirmed mandibular and condylar dysplasia. Inner ears appeared normal. She harbours the heterozygous mutation c.1861C>T; p.(Arg621Cys) in *PLCB4*.

Patient 19, a female, was term-born in Algeria after an uneventful pregnancy with normal birth parameters. Her parents are not consanguineous and have no relevant

medical history. She presented with bilateral QME and retrognathia (Figure 1: g-i). No feeding or breathing difficulties were noted. She developed post-natal growth retardation (weight far below the 3rd centile and height below the 5th centile at 15 years old), delayed puberty, intellectual deficiency and motor dyspraxia. Bilateral otoplasty was performed at 13 years of age and the diagnosis of ACS was made at that time. She reported neither sleeping difficulties nor snoring. Cardiovascular, ophthalmologic and hearing examinations were normal. She harbours a *de novo* heterozygous c.1862G>A; p.(Arg621His) mutation in *PLCB4*.

Patient 23, a female, was born from consanguineous parents originating from Turkey and has three healthy siblings. A club foot and unilateral ear dysplasia were noted on ultrasound at 23 weeks of gestation (WG) and 36 WG, respectively. She was term-born with growth parameters in the normal range. She presented with a right QME and a slight notch on the left ear and mild retrognathia without palate or tongue anomalies (Figure 1: j-l). She had severe mixed sleep apnea on polysomnography at two months of age, requiring nocturnal non-invasive ventilation. Feeding difficulties necessitated tube feeding for 3 months. She was hypotonic and had several acute episodes of hypertonia, dystonia, swallowing difficulties, irregular breathing and nystagmus. External genitalia were normal. She walked unaided at 2.5 years of age and speaks in short sentences at 4.5 years of age but is still not toilet trained. She integrated into kindergarten with a school life assistant. She has a homozygous c.1620dup; p.(Thr541Hisfs*5) mutation in *PLCB4* identified through exome sequencing. The heterozygous parents do not have craniofacial anomalies. The father has unilateral hearing loss of unknown origin, the mother has bilateral iris coloboma and a sister has unilateral iris coloboma.

Patient 28, a male, was term-born after an uneventful pregnancy with normal birth parameters. Parents had no relevant family history. The diagnosis of ACS was made at birth with the association of QME, severe micrognathia, full cheeks and microstomia requiring gastrostomy and tracheostomy. He presented with failure to thrive, but with normal height. Psychomotor development was in the normal range at 12 months of age. Hearing was normal. Craniofacial CT-scan showed mandibular hypoplasia, abnormal condyles and TMJs and unspecified inner ear anomalies. He harbours a *de novo* heterozygous c.133G>A; p.(Gly45Ser) mutation in *GNAI3* (RefSeq NM_006496.4).

Patient 29, a female, is the only child of non-consanguineous parents originating from China. Retrognathia was noted on ultrasound at 33 WG. She was born at 35 WG with parameters in the normal range. She presented with severe ACS, characterized by bilateral QME with thick, overfolded and dysplastic helices, severe microretrognathia, full cheeks, microstomia and cleft palate (Figure 1: m-o) leading to severe upper airway obstruction and feeding difficulties. A patent ductus arteriosus closed on the 9th day of life. She underwent tracheostomy at birth and gastrostomy at 16 months of age. Craniofacial CT imaging demonstrated a small mandible and bilateral TMJ ankylosis that benefited from arthroplasty and costochondral grafts at 4 years of age, followed by reconstructive surgery of the ears at 5 years of age and external mandibular distraction at 6 years of age. Psychomotor development and growth were normal with OFC on the 3rd centile. She had profound bilateral hearing loss, a speech disorder and orofacial dyspraxia. The tongue was normal and gum hypertrophy was noted. She harbours the heterozygous mutation c.140G>A; p.(Ser47Asn) in *GNAI3*. The mutation was inherited from her mother who presented with mild retrognathia and no ear anomalies.

Patient 39, a male, was term-born with normal birth parameters from healthy nonconsanguineous parents originating from Australia. Unilateral hydronephrosis had been noted on ultrasound at 32 WG, and a scarred and small left kidney was noted on subsequent postnatal imaging. At birth, he presented with a right QME with a post-auricular tag (Figure 1: p-r), microstomia, submucous cleft palate and mild microretrognathia. Growth and psychomotor development were normal. Trigger thumbs were noted. Conductive hearing loss necessitated tympanostomy tube insertion. Sleep apnea with snoring were treated by tonsillectomy at 2 years of age. Craniofacial CT-scan was not available. He harbours a heterozygous c.3G>C; p.? mutation in *EDNI* (RefSeq NM_001955.5), inherited from his father, who has normal ears and no micrognathia or abnormal distribution of his facial hair.

Results

Of 44 questionnaires sent to the referring doctors, 39 patients from 27 families with a completed questionnaire were included, comprising 33 ACS and six IQME cases.

A mutation in one of the three known genes was previously reported or identified in this study in all patients except patient 31, an affected member of a family harbouring an *EDNI* mutation (Table 1). We identified six previously undescribed mutations; heterozygous p.(Ile569Phe) and p.(Gly339Arg) in *PLCB4* (families 10 and 11, respectively), homozygous p.(Thr541Hisfs*5) in *PLCB4* (patient 23), heterozygous p.(Gly45Ser) and p.(Ser47Asn) in *GNAI3* (patients 28 and 29, respectively) and heterozygous p.? in *EDNI* (patient 39). When considering index cases from each family, *PLCB4* is the most common gene associated with ACS (16/27, 59%). Of the 16 families with ACS-associated *PLCB4* mutations, 13 harboured heterozygous

missense variants clustered in the catalytic domain of the protein with one hotspot at codon Arg621, which is replaced by a histidine, a cysteine or a leucine in 3, 4 and 1 cases respectively (Table 1). Gly339 and Ile569 are conserved amino acids of the *PLCB4* catalytic X and Y domains respectively, mutations of which are not reported in the gnomAD database. In the remaining three families with *PLCB4* mutations, the variants were homozygous truncating or compound heterozygous affecting essential splice sites (patients 20-23). *GNAI3* missense heterozygous mutations were less frequent (6/27, 22%). All mutations clustered in the G1 box of the catalytic domain except the previously published mutation p.(Asn269Tyr) (located in the G4 box; patient 25). Concerning *GNAI3* codons Gly45 and Ser47 in the G1 motif, mutations at these positions have already been reported (Gordon, Vuillot, et al., 2013; Romanelli Tavares et al., 2015) but the amino acid substitutions are different for the two novel cases we report (patients 28 and 29). The novel *EDN1* mutation we identified (c.3G>C, p.?, patient 39) is not found in gnomAD and is predicted to alter the initiation codon of translation and result in loss-of-function, considering that methionine residues at position 6 and 59 of *EDN1* are not in a consensus Kozak configuration.

As previously mentioned in the literature, incomplete penetrance and variable expressivity is observed within families segregating heterozygous mutations in either *PLCB4* (families 8, 10 and 11) or *GNAI3* (family 18; (Romanelli Tavares et al., 2015)). Variable expressivity among unrelated patients harbouring *PLCB4* recurrent heterozygous mutations is also observed (for example, patients 4, 5, 8 and 18, each with p.(Arg621Cys)).

The core craniofacial features of the 39 patients are described below, followed by extra-craniofacial features.

Craniofacial anomalies

A small mandible was noted at examination in 31 patients (31/35, 88%). Craniofacial imaging, which was available for 26/39 (67%) patients, demonstrated mandibular hypoplasia in 19/26 (73%) and condyle anomalies in 15/26 (58%), the latter consisting of TMJ dysplasia (14/15, 93%) with or without hypoplasia (12/15, 80%) (Figure 2). Interestingly, we observed hypoplasia of the hyoid bone in two patients (patients 5 and 36, Figure 2), while medial and lateral pterygoid muscles could not be distinguished from one another on imaging in patient 5. From photographs, full cheeks were observed in 23/31 (74%) patients and microstomia in 16/39 (41%). Tongue anomalies were noted in 12/34 (35%), the majority being glossoptosis, while orthodontic anomalies were noted in 12/36 (33%).

External ear anomalies ranged from normal to severe microtia (Figure 3), and were frequently asymmetric. QMEs were present in most cases (36/37, 97%) and were predominantly bilateral (23/30, 77%). Post auricular tags were reported in five patients with ACS (6/37, 16%), three of them harbouring a *PLCB4* mutation at codon Arg621 (patients 3, 4 and 5), one being homozygous for a frameshift mutation of *PLCB4* (patient 22) and one carrying a heterozygous mutation of *EDNI* (patient 39; Figure 1). As the presence of ear tags was not specifically elucidated in the questionnaire, the frequency of this feature may be underestimated.

Hearing loss

Hearing loss was noted in eight patients with ACS. Conductive hearing loss (CHL) was reported in five patients (patient 3 with a heterozygous *PLCB4* mutation, patients 26 and 27 with a *GNAI3* mutation, patient 36 with a homozygous *EDNI* mutation and patient 39 with a heterozygous *EDNI* mutation). The type of hearing loss was unspecified in three patients (patients 8 with a heterozygous *PLCB4* mutation and 25 and 29 with a *GNAI3* mutation). Auditory canal stenosis was reported once in patient 36.

Costal anomalies

Rib defects were reported in two patients harboring a *GNAI3* mutation (patients 24 and 26; Figure 4). Case 24 had hypoplastic first ribs and case 26 had a unilateral fusion of the first and second ribs.

Vascular anomalies

Cerebrovascular malformations were detected by brain MRI in two individuals with no neurological symptoms. Patient 35, carrying a homozygous *EDNI* gene mutation, has an aneurysmal dilatation of the vein of Galen associated with two arteriovenous fistulae (Gordon, Petit, et al., 2013; Gordon, Vuillot, et al., 2013). Patient 26, carrying a *GNAI3* mutation, presented with a left parieto-occipital arteriovenous malformation.

A patent ductus arteriosus, closed by fluid restriction by day 9 (patient 29), is the only congenital heart defect identified upon echocardiography performed in 17 individuals.

Neurological anomalies

Psychomotor developmental delay was reported in 13/27 (44%) patients, and includes neonatal hypotonia, walking unaided after 18 months of age and speech delay. However intellectual disability was rare (3/23 patients, 13%) and was associated with autistic spectrum disorder in one individual. Among the 13 patients with psychomotor delay, seven harbored heterozygous *PLCB4* mutations, three had biallelic *PLCB4* mutations, one had a *GNAI3* mutation and two had homozygous *EDNI* mutations. Neuroimaging revealed non-specific abnormalities in two patients (22 and 38).

Finally, sleep disorders were common (18/21, 86%), with many reporting snoring (23/36, 64%), obstructive (8/21 patients, 38%) or mixed apnea (8/21 patients, 38%). Feeding difficulties and/or gastroesophageal reflux (15/38, 48%) were also frequent.

Discussion

This study describes 39 patients with the phenotypic spectrum of IQME-ACS, and has identified several novel variants in the three causal genes.

PLCB4 and *GNAI3* mutations are associated with an ACS phenotype. With respect to *PLCB4*, two modes of inheritance (autosomal dominant or recessive) are observed. All heterozygous missense mutations affect the catalytic domain of the protein and are predicted to act via a dominant negative mechanism. We observed incomplete penetrance and/or variable expression within families. Conversely, while individuals with heterozygous loss-of-function mutations in *PLCB4* are asymptomatic, craniofacial phenotypes are highly penetrant in patients with homozygous or compound heterozygous loss-of-function alleles and they can present with additional features, including central sleep apnea, constipation and genitourinary anomalies (Leoni et al., 2016). With respect to *EDNI*, we previously observed that patients

heterozygous for predicted loss-of-function mutations present with IQME, however patient 39 reported here, with a heterozygous *EDNI* start site mutation, has QME plus mild mandibular hypoplasia. Predicted hypomorphic *EDNI* mutations result in an ACS phenotype in homozygotes and no phenotype in heterozygotes (Gordon, Petit, et al., 2013). With respect to *GNAI3*, only heterozygous mutations have been identified thus far. Expressivity is highly variable within families and craniofacial anomalies can be severe, requiring maxillofacial surgery, tracheostomy and gastrostomy.

It remains unclear whether some associated features of ACS are suggestive of mutation in a specific gene. Postauricular tags have been observed in a few patients with *PLCB4* mutations, but were also observed here in patient 39 with an *EDNI* mutation. Genital anomalies can occur in association with a biallelic mutation in *PLCB4* (Leoni et al., 2016) and rib anomalies may suggest a *GNAI3* mutation (described here for patients 24 and 26). Interestingly, rib and vertebral fusions are observed in *Gnai3* null mice (Plummer et al., 2012). The possibility of vertebral fusion in patients with ACS remains inconclusive and we recommend cervico-thoraco-lumbar x-rays in all patients with a *GNAI3* mutation. However, all of the above proposed associations are based on relatively few patients, and a larger number of patients will be required to confirm these associations.

Previous studies have suggested a homeotic transformation of the mandible into an upper jaw-like structure in patients with ACS and *PLCB4/GNAI3* mutations (Clouthier et al., 2013; Rieder et al., 2012). Other clinical manifestations of a homeotic transformation within the first branchial arch are palate-like soft tissue on the floor of the mouth in some patients (Clouthier et al., 2013; Rieder et al., 2012) and an abnormal pattern of facial hair in adult males ((Gordon et al., 2014) and family 11

reported here). Possible homeotic transformation of the ossicles as well as permanent teeth remains to be investigated. Considering second and third branchial arch development, we report hyoid bone anomalies in patients 5 and 36. Hyoid defects have previously been described in *Edn1*^{-/-} and *Ednra*^{-/-} mice (Clouthier et al., 1998; Ozeki et al., 2004).

Hearing loss in ACS patients has not been thoroughly described. As expected, conductive hearing loss seems to be predominant and has been attributed to fusion of the malleus and incus in a patient with a *GNAI3* mutation (patient 26) (Propst et al., 2013). Interestingly, *Gnai3*^{-/-} mice have cochlear hair cell anomalies (Ezan et al., 2013) and *Gai* isoforms have been shown to play essential and redundant roles in the development and maturation of various cell types of the cochlea, auditory nerve and central auditory neurons (Beer-Hammer et al., 2018). Attention needs to be paid to middle and inner ear function by way of an audiogram in all patients with ACS to evaluate for hearing loss.

Central sleep apnea has previously been reported in patients with ACS. Although it could be the consequence of severe obstructive apnea (Eckert, Malhotra, & Jordan, 2009) it persisted after treatment of the obstruction by distraction surgery for some patients (Gordon, Vuillot, et al., 2013). Therefore monitoring for both obstructive and central apnea should be considered in all patients with ACS. Of note, the Endothelin 1 signaling pathway is involved in the regulation of ventilation in mice (Gaultier, 2004; Gaultier et al., 2004).

Motor and intellectual development is in the normal range in the vast majority of patients with ACS. Although speech delay was reported in several patients, it is unclear to what extent this phenotype may be due to oral motor issues. However,

patients with biallelic *PLCB4* mutations may be at higher risk of developmental delay. Recent reports have highlighted that heterozygous myocyte enhancer factor 2C (*MEF2C*; MIM# 600662) loss-of-function mutations should be considered as a possible cause of QME associated with severe intellectual deficiency and/or epilepsy (Gordon et al., 2018; Le Meur et al., 2010). *Mef2c* is essential for the development of the craniofacial skeleton in mice and zebrafish (Miller et al., 2007; Verzi et al., 2007), and is a transcriptional target of endothelin signalling in cranial neural crest cells (Hu et al., 2015). Finally, brain MRI may be worthwhile, considering the possible risk of neurovascular malformations, even if developmental milestones are within the normal range.

In conclusion, through systematic analysis of a large cohort of patients with ACS, we have highlighted how the core craniofacial features can be associated with rare extra-branchial arch manifestations such as ventilatory, neurodevelopmental, vascular, genital and skeletal anomalies. Additional anomalies of the middle ear, facial musculature and teeth still need to be better elucidated. Taking into account the frequent and rare features of this condition, this study allows us to make recommendations for the assessment and follow-up of patients with ACS.

Acknowledgments

We thank the patients and their physicians for their participation in the study. This work was supported by a Université Sorbonne Paris-Cité Pôle de recherche et d'enseignement supérieur grant (project number SPC/JFG/2013-031), the Agence Nationale de la Recherche (CranioRespiro), an E-Rare CRANIRARE grant, MSD-Avenir (project Devo-Decode) and AXA (project Tête et Cœur).

Footnotes

The authors have no conflicts of interest to declare.

Ethical approval

Informed consent for genetic testing and for publication of images was obtained for all patients. Human genetic research in this study was performed with approval from the institutional review board "Comité de Protection des Personnes Ile-de-France II" (Necker Hospital); approval received 10th June 2015.

Data Availability Statement

All available clinical and sequencing data are provided in the manuscript and supplementary information. Previously unreported variants have been submitted to <http://www.lovd.nl>.

Web Resources

OMIM: <https://www.omim.org>

GnomAD: <https://gnomad.broadinstitute.org>

LOVD: <http://www.lovd.nl>

Mutalyser: <https://mutalyzer.nl/>

References

- Beer-Hammer, S., Lee, S. C., Mauriac, S. A., Leiss, V., Groh, I. A. M., Novakovic, A., ... Rüttiger, L. (2018). Gai Proteins are Indispensable for Hearing. *Cellular Physiology and Biochemistry: International Journal of Experimental Cellular Physiology, Biochemistry, and Pharmacology*, 47(4), 1509-1532. doi: 10.1159/000490867
- Clouthier, D. E., Hosoda, K., Richardson, J. A., Williams, S. C., Yanagisawa, H., Kuwaki, T., ... Yanagisawa, M. (1998). Cranial and cardiac neural crest defects in endothelin-A receptor-deficient mice. *Development (Cambridge, England)*, 125(5), 813-824.
- Clouthier, D.E., Garcia, E., & Schilling, T. F. (2010). Regulation of facial morphogenesis by endothelin signaling: Insights from mice and fish. *American Journal of Medical Genetics. Part A*, 152A(12), 2962-2973. doi: 10.1002/ajmg.a.33568
- Clouthier, D.E., Passos-Bueno, M. R., Tavares, A. L. P., Lyonnet, S., Amiel, J., & Gordon, C. T. (2013). Understanding the basis of auriculocondylar syndrome: Insights from human, mouse and zebrafish genetic studies. *American Journal of Medical Genetics. Part C, Seminars in Medical Genetics*, 163C(4), 306-317. doi: 10.1002/ajmg.c.31376
- Cosman, B. (1984). The question mark ear: An unappreciated major anomaly of the auricle. *Plastic and Reconstructive Surgery*, 73(4), 572-576. doi: 10.1097/00006534-198404000-00009

- Cosman, B., Bellin, H., & Crikelair, G. F. (1970). The Question Mark ear. *Plastic and Reconstructive Surgery*, *46*(5), 454-457. doi: 10.1097/00006534-197011000-00006
- Eckert, D. J., Malhotra, A., & Jordan, A. S. (2009). Mechanisms of apnea. *Progress in Cardiovascular Diseases*, *51*(4), 313-323. doi: 10.1016/j.pcad.2008.02.003
- Ezan, J., Lasvaux, L., Gezer, A., Novakovic, A., May-Simera, H., Belotti, E., ... Montcouquiol, M. (2013). Primary cilium migration depends on G-protein signalling control of subapical cytoskeleton. *Nature Cell Biology*, *15*(9), 1107-1115. doi: 10.1038/ncb2819
- Gaultier, C. (2004). Genes and genetics in respiratory control. *Paediatric Respiratory Reviews*, *5*(3), 166-172. doi: 10.1016/j.prrv.2004.04.012
- Gaultier, C., Amiel, J., Dager, S., Trang, H., Lyonnet, S., Gallego, J., & Simonneau, M. (2004). Genetics and early disturbances of breathing control. *Pediatric Research*, *55*(5), 729-733. doi: 10.1203/01.PDR.0000115677.78759.C5
- Gerkes, E. H., van Ravenswaaij, C. M. A., & van Essen, A. J. (2008). Question mark ears and post-auricular tags. *European Journal of Medical Genetics*, *51*(3), 264-267. doi: 10.1016/j.ejmg.2008.01.002
- Gordon, C. T., Petit, F., Kroisel, P. M., Jakobsen, L., Zechi-Ceide, R. M., Oufadem, M., ... Amiel, J. (2013). Mutations in endothelin 1 cause recessive auriculocondylar syndrome and dominant isolated question-mark ears. *American Journal of Human Genetics*, *93*(6), 1118-1125. doi: 10.1016/j.ajhg.2013.10.023

- Gordon, C. T., Vuillot, A., Marlin, S., Gerkes, E., Henderson, A., AlKindy, A., ... Amiel, J. (2013). Heterogeneity of mutational mechanisms and modes of inheritance in auriculocondylar syndrome. *Journal of Medical Genetics*, *50*(3), 174-186. doi: 10.1136/jmedgenet-2012-101331
- Gordon, Cunniff, C. M., Green, G. E., Zechi-Ceide, R. M., Johnson, J. M., Henderson, A., ... Amiel, J. (2014). Clinical evidence for a mandibular to maxillary transformation in Auriculocondylar syndrome. *American Journal of Medical Genetics. Part A*, *164A*(7), 1850-1853. doi: 10.1002/ajmg.a.36505
- Gordon, Tessier, A., Demir, Z., Goldenberg, A., Oufadem, M., Voisin, N., ... Amiel, J. (2018). The association of severe encephalopathy and question mark ear is highly suggestive of loss of MEF2C function. *Clinical Genetics*, *93*(2), 356-359. doi: 10.1111/cge.13046
- Hu, J., Verzi, M. P., Robinson, A. S., Tang, P. L.-F., Hua, L. L., Xu, S.-M., ... Black, B. L. (2015). Endothelin signaling activates Mef2c expression in the neural crest through a MEF2C-dependent positive-feedback transcriptional pathway. *Development (Cambridge, England)*, *142*(16), 2775-2780. doi: 10.1242/dev.126391
- Kido, Y., Gordon, C. T., Sakazume, S., Ben Bdira, E., Dattani, M., Wilson, L. C., ... Nagai, T. (2013). Further characterization of atypical features in auriculocondylar syndrome caused by recessive PLCB4 mutations. *American Journal of Medical Genetics. Part A*, *161A*(9), 2339-2346. doi: 10.1002/ajmg.a.36066

- Kokitsu-Nakata, N. M., Zechi-Ceide, R. M., Vendramini-Pittoli, S., Romanelli Tavares, V. L., Passos-Bueno, M. R., & Guion-Almeida, M. L. (2012). Auriculo-condylar syndrome. Confronting a diagnostic challenge. *American Journal of Medical Genetics. Part A*, *158A*(1), 59-65. doi: 10.1002/ajmg.a.34337
- Le Meur, N., Holder-Espinasse, M., Jaillard, S., Goldenberg, A., Joriot, S., Amati-Bonneau, P., ... Bonneau, D. (2010). MEF2C haploinsufficiency caused by either microdeletion of the 5q14.3 region or mutation is responsible for severe mental retardation with stereotypic movements, epilepsy and/or cerebral malformations. *Journal of Medical Genetics*, *47*(1), 22-29. doi: 10.1136/jmg.2009.069732
- Leoni, C., Gordon, C. T., Della Marca, G., Giorgio, V., Onesimo, R., Perrino, F., ... Zampino, G. (2016). Respiratory and gastrointestinal dysfunctions associated with auriculo-condylar syndrome and a homozygous PLCB4 loss-of-function mutation. *American Journal of Medical Genetics. Part A*, *170*(6), 1471-1478. doi: 10.1002/ajmg.a.37625
- Marivin, A., Leyme, A., Parag-Sharma, K., DiGiacomo, V., Cheung, A. Y., Nguyen, L. T., ... Garcia-Marcos, M. (2016). Dominant-negative G α subunits are a mechanism of dysregulated heterotrimeric G protein signaling in human disease. *Science Signaling*, *9*(423), ra37. doi: 10.1126/scisignal.aad2429
- Masotti, C., Oliveira, K. G., Poerner, F., Splendore, A., Souza, J., Freitas, R. da S., ... Passos-Bueno, M. R. (2008). Auriculo-condylar syndrome: Mapping of a first

locus and evidence for genetic heterogeneity. *European Journal of Human Genetics: EJHG*, 16(2), 145-152. doi: 10.1038/sj.ejhg.5201955

Miller, C. T., Swartz, M. E., Khuu, P. A., Walker, M. B., Eberhart, J. K., & Kimmel, C. B. (2007). Mef2ca is required in cranial neural crest to effect Endothelin1 signaling in zebrafish. *Developmental Biology*, 308(1), 144-157. doi: 10.1016/j.ydbio.2007.05.018

Ozeki, H., Kurihara, Y., Tonami, K., Watatani, S., & Kurihara, H. (2004). Endothelin-1 regulates the dorsoventral branchial arch patterning in mice. *Mechanisms of Development*, 121(4), 387-395. doi: 10.1016/j.mod.2004.02.002

Plummer, N. W., Spicher, K., Malphurs, J., Akiyama, H., Abramowitz, J., Nürnberg, B., & Birnbaumer, L. (2012). Development of the mammalian axial skeleton requires signaling through the G α (i) subfamily of heterotrimeric G proteins. *Proceedings of the National Academy of Sciences of the United States of America*, 109(52), 21366-21371. doi: 10.1073/pnas.1219810110

Propst, E. J., Ngan, B. Y., Mount, R. J., Martin-Munoz, D., Blaser, S., Harrison, R. V., ... Papsin, B. C. (2013). Ossicular fusion and cholesteatoma in auriculocondylar syndrome: In vivo evidence of arrest of embryogenesis. *The Laryngoscope*, 123(2), 528-532. doi: 10.1002/lary.23492

Rieder, M. J., Green, G. E., Park, S. S., Stamper, B. D., Gordon, C. T., Johnson, J. M., ... Cunningham, M. L. (2012). A human homeotic transformation resulting from mutations in PLCB4 and GNAI3 causes auriculocondylar syndrome. *American Journal of Human Genetics*, 90(5), 907-914. doi: 10.1016/j.ajhg.2012.04.002

- Romanelli Tavares, V. L., Gordon, C. T., Zechi-Ceide, R. M., Kokitsu-Nakata, N. M., Voisin, N., Tan, T. Y., ... Passos-Bueno, M. R. (2015). Novel variants in GNAI3 associated with auriculocondylar syndrome strengthen a common dominant negative effect. *European Journal of Human Genetics: EJHG*, 23(4), 481-485. doi: 10.1038/ejhg.2014.132
- Romanelli Tavares, V. L., Zechi-Ceide, R. M., Bertola, D. R., Gordon, C. T., Ferreira, S. G., Hsia, G. S. P., ... Passos-Bueno, M. R. (2017). Targeted molecular investigation in patients within the clinical spectrum of Auriculocondylar syndrome. *American Journal of Medical Genetics. Part A*, 173(4), 938-945. doi: 10.1002/ajmg.a.38101
- Ruest, L.-B., Xiang, X., Lim, K.-C., Levi, G., & Clouthier, D. E. (2004). Endothelin-A receptor-dependent and -independent signaling pathways in establishing mandibular identity. *Development (Cambridge, England)*, 131(18), 4413-4423. doi: 10.1242/dev.01291
- Thevenon, J., Duffourd, Y., Masurel-Paulet, A., Lefebvre, M., Feillet, F., El Chehadeh-Djebbar, S., ... Rivière, J. B. (2016). Diagnostic odyssey in severe neurodevelopmental disorders: Toward clinical whole-exome sequencing as a first-line diagnostic test. *Clinical Genetics*, 89(6), 700-707. doi: 10.1111/cge.12732
- Verzi, M. P., Agarwal, P., Brown, C., McCulley, D. J., Schwarz, J. J., & Black, B. L. (2007). The transcription factor MEF2C is required for craniofacial development. *Developmental Cell*, 12(4), 645-652. doi: 10.1016/j.devcel.2007.03.007

Walker, M. B., Miller, C. T., Swartz, M. E., Eberhart, J. K., & Kimmel, C. B. (2007). Phospholipase C, beta 3 is required for Endothelin1 regulation of pharyngeal arch patterning in zebrafish. *Developmental Biology*, 304(1), 194-207. doi: 10.1016/j.ydbio.2006.12.027

Tables

Table 1: Molecular data of the 39 patients. Variants are ordered from 5' to 3', in separate blocks for each gene and mode of inheritance. Patient 31, although not tested, is the affected brother of patient 32, harbouring a heterozygous *EDN1* mutation. ACS, Auriculocondylar syndrome; HMZ, homozygous; HTZ, heterozygous; IQME, Isolated question mark ear.

Gene	Type of mutation	cDNA modification	Protein change	Inheritance	Diagnosis	Familial	Patient number	Publication
PLCB4	HTZ	c.983 A>G	p.(His328Arg)	AD	ACS	Yes	7	Kokitsu-Nakata et al, 2012 (case 2). R. Tavares et al., 2017
		c.1015G>A	p.(Gly339Arg)	AD	ACS	Yes	16	0
		c.1015G>A	p.(Gly339Arg)	AD	IQME	Yes	17	0
		c.1072G>C	p.(Glu358Gln)	AD	ACS	No	6	Kokitsu-Nakata et al, 2012 (case 1). R. Tavares et al., 2017
		c.1073A>T	p.(Glu358Val)	AD	ACS	Yes	12	Shkalim et al, 2008, (case III:2). Gordon et al, 2013, (Case 1-III:2)
		c.1073A>T	p.(Glu358Val)	AD	ACS	Yes	13	Shkalim et al, 2008, (case II:3). Gordon et al, 2013, (Case 1-II:1)

	c.170 5A>T	p.(Ile56 9Phe)	AD	ACS	Yes	14	0
	c.170 5A>T	p.(Ile56 9Phe)	AD	ACS	Yes	15	0
	c.186 1C>T	p.(Arg6 21Cys)	AD	ACS	No	4	Gerkes et al, 2008. Gordon et al, 2013, (Case 4)
	c.186 1C>T	p.(Arg6 21Cys)	AD	ACS	Yes	5	Rieder et al, 2012, (case M003). Gordon et al, 2013, (Case 9)
	c.186 1C>T	p.(Arg6 21Cys)	AD	ACS	No	8	0
	c.186 1C>T	p.(Arg6 21Cys)	AD	ACS	No	18	0
	c.186 2G> A	p.(Arg6 21His)	AD	ACS	No	1	Gordon et al, 2013 (case 6)
	c.186 2G> A	p.(Arg6 21His)	AD	ACS	Yes	9	0
	c.186 2G> A	p.(Arg6 21His)	AD	ACS	Yes	10	0
	c.186 2G> A	p.(Arg6 21His)	AD	ACS	Yes	11	0
	c.186 2G>	p.(Arg6 21His)	AD	ACS	No	19	0

		A						
		c.186 2 G>T	p.(Arg6 21Leu)	AD	ACS	Yes	2	Gordon et al, 2013 (case 5, I:1)
		c.186 2 G>T	p.(Arg6 21Leu)	AD	ACS	Yes	3	Gordon et al, 2013 (case 5, II:1)
	HM Z	c.624 delG	p.(Lys2 08Asnfs *5)	AR	ACS	No	22	Leoni et al, 2016
		c.162 0dup	p.(Thr5 41Hisfs *5)	AR	ACS	No	23	0
	HTZ com poun d	c.854 - 1G> A, c.123 8+1G >C		AR	ACS	Yes	20	Kido et al, 2013 (case 1)
		c.854 - 1G> A, c.123 8+1G >C		AR	ACS	Yes	21	Kido et al, 2013 (case 2)
	HTZ	c.133 G>A	p.(Gly4 5Ser)	AD	ACS	No	28	0
		c.134 G>T	p.(Gly4 5Val)	AD	ACS	No	26	Propst et al, 2013. Romanelli Tavares et al, 2015 (case Sp1)
		c.140	p.(Ser4	AD	ACS	Yes	29	0

		G>A	7Asn)						
		c.141 C>A	p.(Ser4 7Arg)	AD	ACS	Yes	24	Gordon et al, 2013 (case 7)	
		c.143 C>A	p.(Thr4 8Asn)	AD	ACS	No	27	Romanelli Tavares et al, 2015 (case Sp2)	
		c.805 A>T	p.(Asn2 69Tyr)	AD	ACS	Yes	25	Guion-Almeida et al. 2002 (Case 1). Masotti et al., 2008 (F2: III-20). Romanelli Tavares et al., 2015 (F1: III-20)	
	EDNI	HTZ	c.3G >C	p.?	AD	ACS	No	39	0
			c.191 T>A	p.(Val6 4Asp)	AD	IQM E	Yes	33	Gordon et al, 2013 (Case 11). Gordon et al, 2013 (Case F2-I:2)
			c.191 T>A	p.(Val6 4Asp)	AD	IQM E	Yes	34	Gordon et al, 2013 (Case 11). Gordon et al, 2013 (Case F2-II:2)
			c.249 T>G	p.(Tyr8 3*)	AD	IQM E	Yes	30	Gordon et al, 2013 (Case 12). Gordon et al, 2013 (Case F3-III:2)
			c.249 T>G	p.(Tyr8 3*)	AD	IQM E	Yes	32	Gordon et al, 2013 (Case 12). Gordon et al, 2013 (Case F3-IV:2)
	HM Z	c.230 C>A	p.(Pro7 7His)	AR	ACS	Yes	38	Guion-Almeida et al, 2002 (Case 2). Gordon et al, 2013 (Case S1- II:1)	
			c.271 A>G	p.(Lys9 1Glu)	AR	ACS	Yes	35	Gordon et al, 2013 (Case 10). Gordon et al, 2013

								(Case F1-II:1)
		c.271 A>G	p.(Lys9 1Glu)	AR	ACS	Yes	36	Gordon et al, 2013 (Case 10). Gordon et al, 2013 (Case F1-II:3)
		c.271 A>G	p.(Lys9 1Glu)	AR	ACS	Yes	37	Gordon et al, 2013 (Case F1-II:4)
	(not teste d)	(not teste d)	(not teste d)	AD	IQM E	Yes	31	Gordon et al, 2013 (Case 12). Gordon et al, 2013 (Case F3-IV:1)

Table 2: Frequencies of major phenotypes in the 39 patients. ID, intellectual deficiency; TMJ, temporomandibular joint.

Question mark ear	total	36/37	97%
	bilateral	23/30	77%
	unilateral	8/30	27%
Post auricular tag		6/37	16%
Hearing loss		7/36	19%
Mandibular hypoplasia (clinical diagnosis)		31/35	89%
Full Cheeks		23/31	74%
Microstomia		16/39	41%
Dental anomalies		12/36	33%

Eye anomalies		5/36	14%
Neuro-developmental defects		15/39	38%
Intellectual Disability		3/23	13%
Maxillo-facial imaging abnormalities		20/21	95%
Mandibular hypoplasia (radiology)		19/26	73%
Condyle anomalies		15/26	58%
Condyle hypoplasia		12/15	80%
TMJ dysplasia		14/15	93%
Bone abnormalities		6/32	19%
Sleep disorder		18/21	86%
Snoring		23/36	64%
Obstructive apnea		8/21	38%
Mixed apnea		8/22	38%
Gastro-intestinal disorder		15/38	48%
Feeding difficulties		12/38	44%

Figures

Figure 1: Photographs of novel ACS cases. In family 11, the father (patient 16) presents ACS (a, b, c) and his son (patient 17) presents IQME (d, e, f). Note the facial hair growth pattern in both patients (arrows), showing no hair along the proximal region of the horizontal branch of the mandible. Patient 19 presents retrognathia, bilateral QME with dysplastic and protruding antihelix and ear lobe hypoplasia (g, h, i; the photographs were taken after otoplasty). Patient 23 presents bilateral QME, retrognathia and full cheeks (j-l). Patient 29 presents severe ACS with micro-retrognathia, full cheeks, microstomia and QMEs with dysplastic and over folded helices (m-o). Patient 39 presents a unilateral QME (p, q) with post auricular tag (r).



Figure 2: **Craniofacial CT images demonstrating lower jaw anomalies in ACS.** For patient 5 (1.a and 2.a) and a sex- and age-matched control (1.b, 2.b), three-dimensional surface reconstructions (1.a-b) and oblique sagittal images through the temporomandibular joint (2.a-b) are shown. In panel 1.a, full arrow indicates mandibular dysplasia (micrognathia with hypoplasia and shortening of the ascending branch), and arrow head indicates dysplasia of the hyoid bone, with horizontalization and hypoplasia of its major branches compared to control (1b). Double full arrow in 2.a indicates increased distance between external auditory canal and glenoid fossa while double dotted arrow indicates dysplastic tympanic bone (oblique and thickened). Asterisk shows dysplastic temporomandibular joint (condylar articular facet flattened and tilted backwards) compare to control (2b).




Figure 3: **Question mark ears with increasing severity of malformation.** a) protruding antihelix without notch of the pinna, b) notch of the pinna, c) notch of the pinna, over-folded helix and post auricular tag, d) cleft between lobe and helix, protruding antihelix and hypoplastic helix, e) cleft between lobe and helix, and Darwin tubercle, f) microtia with thick and overfolded pinna.




Figure 4: **Antero-posterior X-ray of cervical and thoracic spine of patients 24 and 26.** Arrows show the hypoplastic first ribs in patient 24 (a), dashed arrow shows the fusion of the first and second ribs on the right in case 26 (b).

